

CS 3.307: Intro to Stochastic Processes

Tejas Bodas

Assistant Professor, IIIT Hyderabad

Recap

Recap

Introduction to Stochastic processes

Introduction to Stochastic processes

- ▶ Stochastic process $\{X(t), t \in T\}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a collection of random variables defined such that for every $t \in T$ we have $X(t) : \Omega \rightarrow \mathcal{S}$.

Introduction to Stochastic processes

- ▶ Stochastic process $\{X(t), t \in T\}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a collection of random variables defined such that for every $t \in T$ we have $X(t) : \Omega \rightarrow \mathcal{S}$.
- ▶ T is the parameter space (often resembles time) and \mathcal{S} is the state space.

Introduction to Stochastic processes

- ▶ Stochastic process $\{X(t), t \in T\}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a collection of random variables defined such that for every $t \in T$ we have $X(t) : \Omega \rightarrow \mathcal{S}$.
- ▶ T is the parameter space (often resembles time) and \mathcal{S} is the state space.
- ▶ Random variable $X(t)$ is often denoted by $X(\omega, t)$.

Introduction to Stochastic processes

- ▶ Stochastic process $\{X(t), t \in T\}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is a collection of random variables defined such that for every $t \in T$ we have $X(t) : \Omega \rightarrow \mathcal{S}$.
- ▶ T is the parameter space (often resembles time) and \mathcal{S} is the state space.
- ▶ Random variable $X(t)$ is often denoted by $X(\omega, t)$.
- ▶ When t is fixed and ω is the only variable, we have a random variable $X(\cdot, t)$. When ω is fixed and t is the variable, we have a $X(\omega, \cdot)$ as a function of time. This is also called as a realization or sample path of a stochastic process.

Introduction to Stochastic processes

- ▶ When T is countable, we have a discrete time process.

Introduction to Stochastic processes

- ▶ When T is countable, we have a discrete time process.
- ▶ If T is a subset of real line, we have a continuous time process.

Introduction to Stochastic processes

- ▶ When T is countable, we have a discrete time process.
- ▶ If T is a subset of real line, we have a continuous time process.
- ▶ State space could be integers or real numbers

Introduction to Stochastic processes

- ▶ When T is countable, we have a discrete time process.
- ▶ If T is a subset of real line, we have a continuous time process.
- ▶ State space could be integers or real numbers
- ▶ State space could be \mathbb{R}^n or \mathbb{Z}^n valued

Elementary Examples

Elementary Examples

- ▶ The process of rolling a dice 6 times.

Elementary Examples

- ▶ The process of rolling a dice 6 times.
- ▶ Your bank balance over a week.

Elementary Examples

- ▶ The process of rolling a dice 6 times.
- ▶ Your bank balance over a week.
- ▶ Temperature fluctuations in a 1hr window.

Elementary Examples

- ▶ The process of rolling a dice 6 times.
- ▶ Your bank balance over a week.
- ▶ Temperature fluctuations in a 1hr window.
- ▶ Number of customers in IKEA every day.

Introduction to Stochastic processes

A c.t.s.p. is called an *independent increment process* if for any choice of parameters $t_0 < t_1 < \dots < t_n$, the n increment random variables $X(t_1) - X(t_0), X(t_2) - X(t_1), \dots, X(t_n) - X(t_{n-1})$ are independent.

Introduction to Stochastic processes

A c.t.s.p. is called an *independent increment process* if for any choice of parameters $t_0 < t_1 < \dots < t_n$, the n increment random variables $X(t_1) - X(t_0), X(t_2) - X(t_1), \dots, X(t_n) - X(t_{n-1})$ are independent.

The c.t.m.p. is said to have *stationary increments* if in addition $X(t_2 + s) - X(t_1 + s)$ has the same distribution as $X(t_2) - X(t_1)$ for all $t_1, t_2 \in T$ and any $s > 0$.

Examples

- ▶ Sequence of i.i.d random variables.

Examples

- ▶ Sequence of i.i.d random variables.
- ▶ General random walk: If X_1, X_2, \dots is a sequence i.i.d of random variables, then $S_n = \sum_{i=1}^n X_i$ is a random walk.

Examples

- ▶ Sequence of i.i.d random variables.
- ▶ General random walk: If X_1, X_2, \dots is a sequence i.i.d of random variables, then $S_n = \sum_{i=1}^n X_i$ is a random walk.
- ▶ Weiner process: $\{X(t), t \geq 0\}$ is a Weiner process if
 1. $X(0) = 0$
 2. $\{X(t), t \geq 0\}$ has stationary and independent increments
 3. for every $t > 0$, $X(t)$ is normally distributed with mean 0 and variance t .

Examples

- ▶ Sequence of i.i.d random variables.
- ▶ General random walk: If X_1, X_2, \dots is a sequence i.i.d of random variables, then $S_n = \sum_{i=1}^n X_i$ is a random walk.
- ▶ Wiener process: $\{X(t), t \geq 0\}$ is a Wiener process if
 1. $X(0) = 0$
 2. $\{X(t), t \geq 0\}$ has stationary and independent increments
 3. for every $t > 0$, $X(t)$ is normally distributed with mean 0 and variance t .
- ▶ $\{X(t), t \geq 0\}$ is a Markov process if for $t_1 < t_2 < \dots t_n < t$ we have
$$P(X(t) \leq x | X(t_1) = x_1, \dots, X(t_n) = x_n) = P(X(t) \leq x | X(t_n) = x_n)$$

Examples

- ▶ Sequence of i.i.d random variables.
- ▶ General random walk: If X_1, X_2, \dots is a sequence i.i.d of random variables, then $S_n = \sum_{i=1}^n X_i$ is a random walk.
- ▶ Wiener process: $\{X(t), t \geq 0\}$ is a Wiener process if
 1. $X(0) = 0$
 2. $\{X(t), t \geq 0\}$ has stationary and independent increments
 3. for every $t > 0$, $X(t)$ is normally distributed with mean 0 and variance t .
- ▶ $\{X(t), t \geq 0\}$ is a Markov process if for $t_1 < t_2 < \dots t_n < t$ we have
$$P(X(t) \leq x | X(t_1) = x_1, \dots, X(t_n) = x_n) = P(X(t) \leq x | X(t_n) = x_n)$$
- ▶ Random walk and Wiener process are examples of Markov processes.

Bernoulli/Binomial process

- ▶ Bernoulli(p) random variable

Bernoulli/Binomial process

- ▶ Bernoulli(p) random variable
- ▶ Bernoulli process is a sequence of independent r.v.'s $\{X_i, i = 1, 2, \dots\}$ where each X_i is a Bernoulli(p) random variable.

Bernoulli/Binomial process

- ▶ Bernoulli(p) random variable
- ▶ Bernoulli process is a sequence of independent r.v.'s $\{X_i, i = 1, 2, \dots\}$ where each X_i is a Bernoulli(p) random variable.
- ▶ Binomial random variable S_n counts the sum of n independent Bernoulli(p) variables

Bernoulli/Binomial process

- ▶ Bernoulli(p) random variable
- ▶ Bernoulli process is a sequence of independent r.v.'s $\{X_i, i = 1, 2, \dots\}$ where each X_i is a Bernoulli(p) random variable.
- ▶ Binomial random variable S_n counts the sum of n independent Bernoulli(p) variables
- ▶ let X_i denote the associated Bernoulli variable for toss i , $i = 1, \dots, n$. Then $S_n = \sum_{i=1}^n X_i$ denotes the number of heads/event and $P(S_n = k) = \binom{n}{k} p^k (1 - p)^{n-k}$.

Bernoulli/Binomial process

- ▶ Bernoulli(p) random variable
- ▶ Bernoulli process is a sequence of independent r.v.'s $\{X_i, i = 1, 2, \dots\}$ where each X_i is a Bernoulli(p) random variable.
- ▶ Binomial random variable S_n counts the sum of n independent Bernoulli(p) variables
- ▶ let X_i denote the associated Bernoulli variable for toss i , $i = 1, \dots, n$. Then $S_n = \sum_{i=1}^n X_i$ denotes the number of heads/event and $P(S_n = k) = \binom{n}{k} p^k (1 - p)^{n-k}$.
- ▶ $E[S_n]$? $\text{Var}(S_n)$?

Bernoulli/Binomial process

- ▶ $\{S_n = \sum_{i=1}^n X_i, n = 1, 2, \dots\}$ is called as a Binomial process.

Bernoulli/Binomial process

- ▶ $\{S_n = \sum_{i=1}^n X_i, n = 1, 2, \dots\}$ is called as a Binomial process.
- ▶ Let $T := \{\text{smallest } n : S_n > 0.\}$.

Bernoulli/Binomial process

- ▶ $\{S_n = \sum_{i=1}^n X_i, n = 1, 2, \dots\}$ is called as a Binomial process.
- ▶ Let $T := \{\text{smallest } n : S_n > 0.\}$.
- ▶ T is a geometric random variable with parameter p , i.e.,
 $P(T = n_1) = p(1 - p)^{(n_1-1)}$.

Bernoulli/Binomial process

- ▶ $\{S_n = \sum_{i=1}^n X_i, n = 1, 2, \dots\}$ is called as a Binomial process.
- ▶ Let $T := \{\text{smallest } n : S_n > 0.\}$.
- ▶ T is a geometric random variable with parameter p , i.e.,
 $P(T = n_1) = p(1 - p)^{(n_1-1)}$.
- ▶ Memoryless property: $P(T > m + n | T > n) = P(T > m)$.

Counting process

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

It satisfies the following

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

It satisfies the following

- ▶ $N(t) \geq 0$ and is integer valued

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

It satisfies the following

- ▶ $N(t) \geq 0$ and is integer valued
- ▶ For $s \leq t$, we have $N(s) \leq N(t)$. $N(t) - N(s)$ denotes the number of events in the interval (t, s)

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

It satisfies the following

- ▶ $N(t) \geq 0$ and is integer valued
- ▶ For $s \leq t$, we have $N(s) \leq N(t)$. $N(t) - N(s)$ denotes the number of events in the interval (t, s)
- ▶ $N(t)$ can have independent increments

Counting process

Stochastic process $\{N(t), t \geq 0\}$ is a counting process if it represents the total number of events upto time t .

It satisfies the following

- ▶ $N(t) \geq 0$ and is integer valued
- ▶ For $s \leq t$, we have $N(s) \leq N(t)$. $N(t) - N(s)$ denotes the number of events in the interval (t, s)
- ▶ $N(t)$ can have independent increments
- ▶ $N(t)$ can have stationary increments

Poisson process

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments
- ▶ Number of events in an interval of length t is a Poisson distribution with mean λt . (Hence stationary increments)

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments
- ▶ Number of events in an interval of length t is a Poisson distribution with mean λt . (Hence stationary increments)
- ▶ $E[N(t + s) - N(t)] = \lambda s$

Poisson process

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments
- ▶ Number of events in an interval of length t is a Poisson distribution with mean λt . (Hence stationary increments)
- ▶ $E[N(t + s) - N(t)] = \lambda s$

Condition 3 is difficult to verify ! Hence ...

Poisson process – Alternative definition

A function f is said to be $o(h)$ if $\lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$.

Poisson process – Alternative definition

A function f is said to be $o(h)$ if $\lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$.

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

Poisson process – Alternative definition

A function f is said to be $o(h)$ if $\lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$.

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$

Poisson process – Alternative definition

A function f is said to be $o(h)$ if $\lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$.

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments

Poisson process – Alternative definition

A function f is said to be $o(h)$ if $\lim_{h \rightarrow 0} \frac{f(h)}{h} = 0$.

A Poisson process with rate λ , $\lambda \geq 0$ is a counting process $\{N(t), t \geq 0\}$ with the following properties

- ▶ $N(0) = 0$
- ▶ $N(t)$ has independent and stationary increments
- ▶ $P\{N(h) = 1\} = \lambda h + o(h)$
- ▶ $P\{N(h) \geq 2\} = o(h)$

Poisson process

Poisson process

Lemma

Definition 1 \implies *Definition 2*

Proof on board.

Poisson process

Lemma

Definition 1 \implies Definition 2

Proof on board.

Lemma

Definition 2 \implies Definition 1

Poisson process

Lemma

Definition 1 \implies Definition 2

Proof on board.

Lemma

Definition 2 \implies Definition 1

Self Study: Refer Sheldon Ross, Stochastic processes, Theorem 2.1.1