

# BASICS OF PROGRAMMING

## ASSIGNMENT - 2

LAKSHMI GAYATHRI GUDIPUDI - SM21MTECH11001

### CHAPTER III EX-IV Q-5

Find the condition that the lines

$$y + t_i x = 2at_i + at_i^3$$

where  $i=1,2,3$ , are concurrent.

$$t_1 x + y = 2at_1 + at_1^3$$

$$t_2 x + y = 2at_2 + at_2^3$$

$$t_3 x + y = 2at_3 + at_3^3$$

### SOLUTION

Considering coefficients of three lines in matrix form :

$$(t_1 \ 1) x = 2at_1 + at_1^3 \quad (1)$$

$$(t_2 \ 1) x = 2at_2 + at_2^3 \quad (2)$$

$$(t_3 \ 1) x = 2at_3 + at_3^3 \quad (3)$$

The above equations form a matrix equation as below:

$$\begin{pmatrix} t_1 & 1 \\ t_2 & 1 \\ t_3 & 1 \end{pmatrix} x = \begin{pmatrix} 2at_1 + at_1^3 \\ 2at_2 + at_2^3 \\ 2at_3 + at_3^3 \end{pmatrix} \quad (4)$$

Given the lines are concurrent, so considering above equations are consistent and are reduced to augmented form as below to find the condition for lines to be concurrent:

$$\begin{pmatrix} t_1 & 1 & -2at_1 - at_1^3 \\ t_2 & 1 & -2at_2 - at_2^3 \\ t_3 & 1 & -2at_3 - at_3^3 \end{pmatrix} \quad (5)$$

Considering augmented form of matrix, we get:

$$\begin{pmatrix} t_1 & t_2 & t_3 \\ 1 & 1 & 1 \\ -2at_1 - at_1^3 & -2at_2 - at_2^3 & -2at_3 - at_3^3 \end{pmatrix}$$

Performing row operations on the above augmented matrix as follows:

$$\begin{pmatrix} t_1 & t_2 & t_3 \\ 1 & 1 & 1 \\ -2at_1 - at_1^3 & -2at_2 - at_2^3 & -2at_3 - at_3^3 \end{pmatrix} \xrightarrow{R_3 \leftarrow R_3 + 2aR_1} \begin{pmatrix} t_1 & t_2 & t_3 \\ 1 & 1 & 1 \\ -at_1^3 & -at_2^3 & -at_3^3 \end{pmatrix} \xrightarrow{R_3 \leftarrow R_3 / (-a)} \begin{pmatrix} t_1 & t_2 & t_3 \\ 1 & 1 & 1 \\ t_1^3 & t_2^3 & t_3^3 \end{pmatrix} \quad (6)$$

Since system of equations are considered consistent, we get

$$t_1(t_3^3 - t_2^3) - t_2(t_3^3 - t_1^3) + t_3(t_2^3 - t_1^3) = 0 \quad (7)$$

$$t_1(t_2^3 - t_3^3) + t_2(t_3^3 - t_1^3) + t_3(t_1^3 - t_2^3) = 0 \quad (8)$$

$$t_1 t_2^3 - t_1 t_3^3 + t_2 t_3^3 - t_2 t_1^3 + t_3 t_1^3 - t_3 t_2^3 = 0 \quad (9)$$

$$t_1 t_2(t_2^2 - t_1^2) + t_2 t_3(t_2^2 - t_1^2) + t_3 t_1(t_1^2 - t_3^2) = 0 \quad (10)$$

Simplifying further Equation (10) we get

$$(t_1 t_2(t_2 - t_1)(t_2 + t_1)) + (t_2 t_3(t_3 - t_2)(t_3 + t_2)) + (t_3 t_1(t_1 - t_3)(t_1 + t_3)) = 0 \quad (11)$$

Therefore Equation (11) represents the condition for the three lines to be concurrent.