

Machine Learning

CSE 6363 (Fall 2019)



Lecture 10 Support Vector Machine

Dajiang Zhu, Ph.D.

Department of Computer Science and Engineering




History of SVM

- SVM is related to statistical learning theory [3]
- SVM was first introduced in 1992 [1]
- SVM becomes popular because of its success in handwritten digit recognition
 - 1.1% test error rate for SVM. This is the same as the error rates of a carefully constructed neural network
- SVM is now regarded as an important example of “kernel methods”, one of the key area in machine learning



- [1] B.E. Boser *et al.* A Training Algorithm for Optimal Margin Classifiers. Proceedings of the Fifth Annual Workshop on Computational Learning Theory 5 144-152, Pittsburgh, 1992.
- [2] L. Bottou *et al.* Comparison of classifier methods: a case study in handwritten digit recognition. Proceedings of the 12th IAPR International Conference on Pattern Recognition, vol. 2, pp. 77-82.
- [3] V. Vapnik. The Nature of Statistical Learning Theory. 2nd edition, Springer, 1999.

What is a good Decision Boundary?

- Consider a two-class, linearly separable classification problem
- Many decision boundaries!
 - The Perceptron algorithm can be used to find such a boundary
 - Different algorithms have been proposed
- Are all decision boundaries equally good?




Examples of Bad Decision Boundaries



Large-margin Decision Boundary

- The decision boundary should be as far away from the data of both classes as possible
 - We should maximize the margin, m
 - Distance between the origin and the line $\mathbf{w}^T \mathbf{x} = k$ is $k/||\mathbf{w}||$



$$m = \frac{(x^+ - x^-) \cdot \mathbf{w}}{|\mathbf{w}|} = \frac{2}{|\mathbf{w}|}$$

Finding the Decision Boundary

- Let $\{x_1, \dots, x_n\}$ be our data set and let $y_i \in \{1, -1\}$ be the class label of x_i
- The decision boundary should classify all points correctly
 $\Rightarrow y_i(\mathbf{w}^T \mathbf{x}_i + b) \geq 1, \quad \forall i$
- The decision boundary can be found by solving the following constrained optimization problem

$$\begin{aligned} &\text{Minimize } \frac{1}{2} \|\mathbf{w}\|^2 \\ &\text{subject to } y_i(\mathbf{w}^T \mathbf{x}_i + b) \geq 1 \quad \forall i \end{aligned}$$

- This is a constrained optimization problem. Solving it requires some new tools
 - Feel free to ignore the following several slides; what is important is the constrained optimization problem above

Recap of Constrained Optimization

- Suppose we want to: minimize $f(\mathbf{x})$ subject to $g(\mathbf{x}) = 0$
- A necessary condition for \mathbf{x}_0 to be a solution:

$$\begin{cases} \frac{\partial}{\partial \mathbf{x}}(f(\mathbf{x}) + \alpha g(\mathbf{x})) \Big|_{\mathbf{x}=\mathbf{x}_0} = 0 \\ g(\mathbf{x}) = 0 \end{cases}$$

- α : the Lagrange multiplier
- For multiple constraints $g_i(\mathbf{x}) = 0$, $i=1, \dots, m$, we need a Lagrange multiplier α_i for each of the constraints

$$\begin{cases} \frac{\partial}{\partial \mathbf{x}}(f(\mathbf{x}) + \sum_{i=1}^n \alpha_i g_i(\mathbf{x})) \Big|_{\mathbf{x}=\mathbf{x}_0} = 0 \\ g_i(\mathbf{x}) = 0 \quad \text{for } i = 1, \dots, m \end{cases}$$

Recap of Constrained Optimization

- The case for inequality constraint $g_i(\mathbf{x}) \leq 0$ is similar, except that the Lagrange multiplier α_i should be positive
- If \mathbf{x}_0 is a solution to the constrained optimization problem

$$\min_{\mathbf{x}} f(\mathbf{x}) \quad \text{subject to} \quad g_i(\mathbf{x}) \leq 0 \quad \text{for } i = 1, \dots, m$$

- There must exist $\alpha_i \geq 0$ for $i=1, \dots, m$ such that \mathbf{x}_0 satisfy

$$\begin{cases} \left. \frac{\partial}{\partial \mathbf{x}} \left(f(\mathbf{x}) + \sum_i \alpha_i g_i(\mathbf{x}) \right) \right|_{\mathbf{x}=\mathbf{x}_0} = 0 \\ g_i(\mathbf{x}) \leq 0 \quad \text{for } i = 1, \dots, m \end{cases}$$

- The function $f(\mathbf{x}) + \sum_i \alpha_i g_i(\mathbf{x})$ is also known as the Lagrangian; we want to set its gradient to **0**

Back to the Original Problem

$$\text{Minimize } \frac{1}{2} \|\mathbf{w}\|^2$$

subject to $1 - y_i(\mathbf{w}^T \mathbf{x}_i + b) \leq 0$ for $i = 1, \dots, n$

■ The Lagrangian is

$$\mathcal{L} = \frac{1}{2} \mathbf{w}^T \mathbf{w} + \sum_{i=1}^n \alpha_i (1 - y_i(\mathbf{w}^T \mathbf{x}_i + b))$$

■ Note that $\|\mathbf{w}\|^2 = \mathbf{w}^T \mathbf{w}$

■ Setting the gradient of \mathcal{L} w.r.t. \mathbf{w} and b to zero, we have

$$\mathbf{w} + \sum_{i=1}^n \alpha_i (-y_i) \mathbf{x}_i = \mathbf{0} \quad \Rightarrow \quad \mathbf{w} = \sum_{i=1}^n \alpha_i y_i \mathbf{x}_i$$

$$\sum_{i=1}^n \alpha_i y_i = 0$$

The Dual Problem

- If we substitute $\mathbf{w} = \sum_{i=1}^n \alpha_i y_i \mathbf{x}_i$ to \mathcal{L} , we have

$$\begin{aligned}\mathcal{L} &= \frac{1}{2} \sum_{i=1}^n \alpha_i y_i \mathbf{x}_i^T \sum_{j=1}^n \alpha_j y_j \mathbf{x}_j + \sum_{i=1}^n \alpha_i \left(1 - y_i \left(\sum_{j=1}^n \alpha_j y_j \mathbf{x}_j^T \mathbf{x}_i + b \right) \right) \\ &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j + \sum_{i=1}^n \alpha_i - \sum_{i=1}^n \alpha_i y_i \sum_{j=1}^n \alpha_j y_j \mathbf{x}_j^T \mathbf{x}_i - b \sum_{i=1}^n \alpha_i y_i \\ &= -\frac{1}{2} \sum_{i=1}^n \sum_{j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j + \sum_{i=1}^n \alpha_i\end{aligned}$$

- Note that $\sum_{i=1}^n \alpha_i y_i = 0$
- This is a function of α_i only

The Dual Problem

- The new objective function is in terms of α_i only
- It is known as the dual problem: if we know \mathbf{w} , we know all α_i ; if we know all α_i , we know \mathbf{w}
- The original problem is known as the primal problem
- The objective function of the dual problem needs to be maximized!
- The dual problem is therefore:

$$\max. W(\boldsymbol{\alpha}) = \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j$$

$$\text{subject to } \alpha_i \geq 0, \quad \sum_{i=1}^n \alpha_i y_i = 0$$

Properties of α_i when we introduce the Lagrange multipliers

The result when we differentiate the original Lagrangian w.r.t. b



The Dual Problem

$$\begin{aligned} \max. \quad W(\boldsymbol{\alpha}) &= \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j \\ \text{subject to } \alpha_i &\geq 0, \sum_{i=1}^n \alpha_i y_i = 0 \end{aligned}$$

- This is a quadratic programming (QP) problem
 - A global maximum of α_i can always be found

- \mathbf{w} can be recovered by
$$\mathbf{w} = \sum_{i=1}^n \alpha_i y_i \mathbf{x}_i$$



Characteristics of the Solution


- Many of the α_i are zero
 - \mathbf{w} is a linear combination of a small number of data points
- \mathbf{x}_i with non-zero α_i are called support vectors (SV)
 - The decision boundary is determined only by the SV
 - Let t_j ($j=1, \dots, s$) be the indices of the s support vectors.
We can write $\mathbf{w} = \sum_{j=1}^s \alpha_{t_j} y_{t_j} \mathbf{x}_{t_j}$
- For testing with a new data \mathbf{z}
 - Compute $\mathbf{w}^T \mathbf{z} + b = \sum_{j=1}^s \alpha_{t_j} y_{t_j} (\mathbf{x}_{t_j}^T \mathbf{z}) + b$ and classify \mathbf{z} as class 1 if the sum is positive, and class 2 otherwise
 - Note: \mathbf{w} need not be formed explicitly



The Quadratic Programming Problem

- Many approaches have been proposed
 - Loqo, cplex, etc. (see <http://www.numerical.rl.ac.uk/qp/qp.html>)
- Most are “interior-point” methods
 - Start with an initial solution that can violate the constraints
 - Improve this solution by optimizing the objective function and/or reducing the amount of constraint violation
- For SVM, sequential minimal optimization (SMO) seems to be the most popular
 - A QP with two variables is trivial to solve
 - Each iteration of SMO picks a pair of (α_i, α_j) and solve the QP with these two variables; repeat until convergence
- In practice, we can just regard the QP solver as a “black-box” without bothering how it works


A Geometrical Interpretation



Dataset with Noise

• denotes +1


◦ denotes -1




- **Hard Margin:** So far we require all data points be classified correctly
 - No training error
- **What if the training set is noisy?**
 - **Solution 1:** use very powerful kernels

OVERFITTING!

Robustness of Soft vs Hard Margin SVMs




Soft Margin SVN



Hard Margin SVN

Non-linearly Separable Problems

- We allow “error” ξ_i in classification; it is based on the output of the discriminant function $\mathbf{w}^T \mathbf{x} + b$
- ξ_i approximates the number of misclassified samples



Soft Margin Hyperplane

- If we minimize $\sum_i \xi_i$, ξ_i can be computed by

$$\begin{cases} \mathbf{w}^T \mathbf{x}_i + b \geq 1 - \xi_i & y_i = 1 \\ \mathbf{w}^T \mathbf{x}_i + b \leq -1 + \xi_i & y_i = -1 \\ \xi_i \geq 0 & \forall i \end{cases}$$

- ξ_i are “slack variables” in optimization
- Note that $\xi_i=0$ if there is no error for \mathbf{x}_i
- ξ_i is an upper bound of the number of errors
- We want to minimize $\frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^n \xi_i$
 - C : tradeoff parameter between error and margin
- The optimization problem becomes

$$\text{Minimize } \frac{1}{2} \|\mathbf{w}\|^2 + C \sum_{i=1}^n \xi_i$$

$$\text{subject to } y_i(\mathbf{w}^T \mathbf{x}_i + b) \geq 1 - \xi_i, \quad \xi_i \geq 0$$



The Optimization Problem

- The dual of this new constrained optimization problem is

$$\max. \quad W(\alpha) = \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j$$

See Bishop:
7.22-7.32

$$\text{subject to } C \geq \alpha_i \geq 0, \quad \sum_{i=1}^n \alpha_i y_i = 0$$



- \mathbf{w} is recovered as $\mathbf{w} = \sum_{j=1}^s \alpha_{t_j} y_{t_j} \mathbf{x}_{t_j}$
- This is very similar to the optimization problem in the linear separable case, except that there is an upper bound C on α_i now
- Once again, a QP solver can be used to find α_i




Extension to Non-linear Decision Boundary

- So far, we have only considered large-margin classifier with a linear decision boundary
- How to generalize it to become nonlinear?
- Key idea: transform \mathbf{x}_i to a higher dimensional space to “make life easier”
 - Input space: the space the point \mathbf{x}_i are located
 - Feature space: the space of $\phi(\mathbf{x}_i)$ after transformation
- Why transform?
 - Linear operation in the feature space is equivalent to non-linear operation in input space
 - Classification can become easier with a proper transformation. In the XOR problem, for example, adding a new feature of x_1x_2 make the problem linearly separable

Transforming the Data



$\phi(\cdot)$




Note: feature space is of higher dimension than the input space in practice

- Computation in the feature space can be costly because it is high dimensional
 - The feature space is typically infinite-dimensional!
- The kernel trick comes to rescue

The Kernel Trick

- Recall the SVM optimization problem

$$\begin{aligned} \max. \quad W(\alpha) &= \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j \\ \text{subject to } C &\geq \alpha_i \geq 0, \quad \sum_{i=1}^n \alpha_i y_i = 0 \end{aligned}$$


- The data points only appear as **inner product**
- As long as we can calculate the inner product in the feature space, we do not need the mapping explicitly
- Many common geometric operations (angles, distances) can be expressed by inner products
- Define the kernel function K by

$$K(\mathbf{x}_i, \mathbf{x}_j) = \phi(\mathbf{x}_i)^T \phi(\mathbf{x}_j)$$

An Example for $\phi(\cdot)$ and $K(\cdot, \cdot)$

- Suppose $\phi(\cdot)$ is given as follows

$$\phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right) = (1, \sqrt{2}x_1, \sqrt{2}x_2, x_1^2, x_2^2, \sqrt{2}x_1x_2)$$

- An inner product in the feature space is

$$\langle \phi\left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}\right), \phi\left(\begin{bmatrix} y_1 \\ y_2 \end{bmatrix}\right) \rangle = (1 + x_1y_1 + x_2y_2)^2$$

- So, if we define the kernel function as follows, there is no need to carry out $\phi(\cdot)$ explicitly

$$K(\mathbf{x}, \mathbf{y}) = (1 + x_1y_1 + x_2y_2)^2$$

- This use of kernel function to avoid carrying out $\phi(\cdot)$ explicitly is known as the **kernel trick**



Kernel Functions

- In practical use of SVM, the user specifies the kernel function; the transformation $\phi(\cdot)$ is not explicitly stated
- Given a kernel function $K(\mathbf{x}_i, \mathbf{x}_j)$, the transformation $\phi(\cdot)$ is given by its eigenfunctions (a concept in functional analysis)
 - Eigenfunctions can be difficult to construct explicitly
 - This is why people only specify the kernel function without worrying about the exact transformation
- Another view: kernel function, being an inner product, is really a similarity measure between the objects



Examples of Kernel Functions

- Polynomial kernel with degree d

$$K(\mathbf{x}, \mathbf{y}) = (\mathbf{x}^T \mathbf{y} + 1)^d$$

- Radial basis function kernel with width σ

$$K(\mathbf{x}, \mathbf{y}) = \exp(-\|\mathbf{x} - \mathbf{y}\|^2 / (2\sigma^2))$$

- Closely related to radial basis function neural networks
- The feature space is infinite-dimensional

- Sigmoid with parameter κ and θ

$$K(\mathbf{x}, \mathbf{y}) = \tanh(\kappa \mathbf{x}^T \mathbf{y} + \theta)$$

- It does not satisfy the Mercer condition on all κ and θ

Modification Due to Kernel Function

- Change all inner products to kernel functions
- For training,

Original

$$\max. W(\alpha) = \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j \mathbf{x}_i^T \mathbf{x}_j$$

$$\text{subject to } C \geq \alpha_i \geq 0, \sum_{i=1}^n \alpha_i y_i = 0$$

With kernel
function

$$\max. W(\alpha) = \sum_{i=1}^n \alpha_i - \frac{1}{2} \sum_{i=1, j=1}^n \alpha_i \alpha_j y_i y_j K(\mathbf{x}_i, \mathbf{x}_j)$$

$$\text{subject to } C \geq \alpha_i \geq 0, \sum_{i=1}^n \alpha_i y_i = 0$$

Modification Due to Kernel Function

- For testing, the new data \mathbf{z} is classified as class 1 if $f \geq 0$, and as class 2 if $f < 0$

Original

$$\mathbf{w} = \sum_{j=1}^s \alpha_{t_j} y_{t_j} \mathbf{x}_{t_j}$$
$$f = \mathbf{w}^T \mathbf{z} + b = \sum_{j=1}^s \alpha_{t_j} y_{t_j} \mathbf{x}_{t_j}^T \mathbf{z} + b$$

With kernel
function

$$\mathbf{w} = \sum_{j=1}^s \alpha_{t_j} y_{t_j} \phi(\mathbf{x}_{t_j})$$
$$f = \langle \mathbf{w}, \phi(\mathbf{z}) \rangle + b = \sum_{j=1}^s \alpha_{t_j} y_{t_j} K(\mathbf{x}_{t_j}, \mathbf{z}) + b$$


More on Kernel Functions

- Since the training of SVM only requires the value of $K(\mathbf{x}_i, \mathbf{x}_j)$, there is no restriction of the form of \mathbf{x}_i and \mathbf{x}_j
 - \mathbf{x}_i can be a sequence or a tree, instead of a feature vector
- $K(\mathbf{x}_i, \mathbf{x}_j)$ is just a similarity measure comparing \mathbf{x}_i and \mathbf{x}_j
- For a test object \mathbf{z} , the discriminant function essentially is a weighted sum of the similarity between \mathbf{z} and a pre-selected set of objects (the support vectors)


$$f(\mathbf{z}) = \sum_{\mathbf{x}_i \in S} \alpha_i y_i K(\mathbf{z}, \mathbf{x}_i) + b$$

S : the set of support vectors


SVM Examples




linear



2^{nd} order polynomial



4^{th} order polynomial



8^{th} order polynomial

Example

- Suppose we have 5 1D data points
 - $x_1=1, x_2=2, x_3=4, x_4=5, x_5=6$, with 1, 2, 6 as class 1 and 4, 5 as class 2 $\Rightarrow y_1=1, y_2=1, y_3=-1, y_4=-1, y_5=1$
- We use the polynomial kernel of degree 2
 - $K(x,y) = (xy+1)^2$
 - C is set to 100
- We first find α_i ($i=1, \dots, 5$) by

$$\begin{aligned} \max. \quad & \sum_{i=1}^5 \alpha_i - \frac{1}{2} \sum_{i=1}^5 \sum_{j=1}^5 \alpha_i \alpha_j y_i y_j (x_i x_j + 1)^2 \\ \text{subject to} \quad & 100 \geq \alpha_i \geq 0, \sum_{i=1}^5 \alpha_i y_i = 0 \end{aligned}$$

Example

- By using a QP solver, we get
 - $\alpha_1=0, \alpha_2=2.5, \alpha_3=0, \alpha_4=7.333, \alpha_5=4.833$
 - Note that the constraints are indeed satisfied
 - The support vectors are $\{x_2=2, x_4=5, x_5=6\}$


- The discriminant function is

$$\begin{aligned} f(z) &= 2.5(1)(2z + 1)^2 + 7.333(-1)(5z + 1)^2 + 4.833(1)(6z + 1)^2 + b \\ &= 0.6667z^2 - 5.333z + b \end{aligned}$$

Diagrammatic annotations: Red arrows point from α_5 to the coefficient 4.833, from y_5 to the constant term 1, and from $K(z, x_5)$ to the polynomial $(6z + 1)^2$.

- b is recovered by solving $f(2)=1$ or by $f(5)=-1$ or by $f(6)=1$, as x_2 and x_5 lie on the line $\phi(w)^T \phi(x) + b = 1$ and x_4 lies on the line $\phi(w)^T \phi(x) + b = -1$
- All three give $b=9 \rightarrow f(z) = 0.6667z^2 - 5.333z + 9$

Example





Choosing the Kernel Function

- Probably the most tricky part of using SVM.
 - The kernel function is important because it creates the kernel matrix, which summarizes all the data
 - Many principles have been proposed (diffusion kernel, Fisher kernel, string kernel, ...)
 - There is even research to estimate the kernel matrix from available information
-
- In practice, a low degree polynomial kernel or RBF kernel with a reasonable width is a good initial try
 - Note that SVM with RBF kernel is closely related to RBF neural networks, with the centers of the radial basis functions automatically chosen for SVM



Strengths and Weaknesses of SVM

■ Strengths

- Training is relatively easy
 - No local optimal, unlike in neural networks
- It scales relatively well to high dimensional data
- Tradeoff between classifier complexity and error can be controlled explicitly
- Non-traditional data like strings and trees can be used as input to SVM, instead of feature vectors

■ Weaknesses

- Need to choose a “good” kernel function.