



MOHAMMED VI
POLYTECHNIC
UNIVERSITY

Ingénierie des Systèmes Complexes et Systèmes Humains

Master Modélisation Hybride Avancée et Calcul Scientifique

M9: Méthodes Numériques

Notes de cours sur la Méthode des Éléments Finis

Années 2019-2020

Abdellah El Kacimi

1 erreur9 avertissements

Contents

Contents	2
Introduction	6
1 The Exact Boundary controllability	7
1.1 Formulation of the control problem	7
1.2 Existence and uniqueness of solutions	8
1.3 Controllability by minimization	9
2 Conjugate Gradient Algorithm	15
2.1 Variational Problem	15
2.2 General conjugate gradient algorithm	16
2.3 Application of conjugate gradient algorithm	17
Conclusion	21

List of figures

List of tables

Introduction

FE

Chapitre 1

The Exact Boundary controllability

1.1 Formulation of the control problem

Let Ω be a bounded domain (that is, non-empty open connected set) in \mathbb{R}^n with boundary $\Gamma = \partial\Omega$ "Sufficiently smooth", Γ_1 be an open nonempty subset of Γ and $\Gamma_2 = \Gamma \setminus \Gamma_1$. With T a given positive number, we consider the following non-homogeneous wave equation :

$$\left\{ \begin{array}{ll} \frac{\partial^2 y}{\partial t^2} - \Delta y = 0 & \text{for } (x, t) \in \Omega \times (0, T), \quad (1.1.1) \\ y(x, t) = 0 & \text{for } (x, t) \in \Gamma_2 \times (0, T), \quad (1.1.2) \\ y(x, t) = v(x, t) & \text{for } (x, t) \in \Gamma_1 \times (0, T), \quad (1.1.2) \\ y(x, 0) = y^0, \quad \frac{\partial y}{\partial t}(x, 0) = y^1 & \text{for } x \in \Omega. \quad (1.1.3) \end{array} \right. \quad (1.1)$$

(1.1.2) : The boundary conditions.

(1.1.3) The initial conditions.

In (1.1), $y^0 \in L^2(\Omega)$, $y^1 \in H^{-1}(\Omega)$, such that $H^{-1}(\Omega)$ is the topological dual space of $H_0^{-1}(\Omega)$ and Δ is the Laplacian operator.

We shall now define the exact boundary controllability for the system (1.1).

Definition 1.1.1. System (1.1) is controllable in time $T > 0$ if for every initial data $(y^0, y^1) \in L^2(\Omega) \times H^{-1}(\Omega)$, we can find a control function $v \in L^2(\Gamma_1 \times (0, T))$ such that the corresponding solution (y, y') of (1.1) verifies

$$y(., T) = y'(., T) = 0. \quad (1.2)$$

Remark 1.1.1. For the existence of solution see section [1.2]

Remark 1.1.2. If the solution of (1.1) verifies (1.2) is also said to be null controllable in time $T > 0$, for more details (see [1], page 100).

The problem that we consider is the following one :
is it possible to find $T > 0$ sufficiently large or optimal and $v \in L^2(\Gamma_1 \times (0, T))$ a boundary

control function such that for any $(y^0, y^1) \in L^2(\Omega) \times H^{-1}(\Omega)$ we have (1.2).

But before that, we have to prove that the system (1.1) admits a solution.

1.2 Existence and uniqueness of solutions

In this section we consider the general case of (1.1) with v not necessarily null and a bounded domain Ω in \mathbb{R}^n .

Definition 1.2.1. For (y^0, y^1) in $L^2(\Omega) \times H^{-1}(\Omega)$ and $v \in L^2(\Gamma_1 \times [0, +\infty))$ a function $y \in C([0, +\infty), L^2(\Omega)) \cap C^1([0, +\infty), H^{-1}(\Omega))$ is called a weak solution of (1.1) if the relation

$$\begin{aligned} \int_{\Omega} y(x, t) \varphi(x) dx - \int_{\Omega} y^0(x) \varphi(x) dx - t \langle y^1, \varphi \rangle_{-1,1} &= \int_0^t \int_0^s \int_{\Omega} y(x, \xi) \Delta \varphi(x) dx d\xi ds \\ &- \int_0^t \int_0^s \int_{\Gamma_1} v(x, \xi) \frac{\partial \varphi}{\partial n}(x) d\sigma d\xi ds, \end{aligned} \quad (1.3)$$

holds for every $t \geq 0$ and every $\varphi \in H^2(\Omega) \cap H_0^1(\Omega)$.

Remark 1.2.1. For x in Γ and φ in $H^2(\Omega)$ we have

$$\frac{\partial \varphi}{\partial n}(x) = \nabla \varphi(x) \cdot \vec{n}(x),$$

is called the normal derivative, with $\vec{n}(x)$ is the normal vector at x

Remark 1.2.2. The normal vector exists because the boundary Γ is sufficiently smooth.

The main result for the existence of solutions of (1.1) is the following :

Theorem 1.2.1. For every v in $L^2(\Gamma_1 \times (0, T))$ and (y^0, y^1) in $L^2(\Omega) \times H^{-1}$ system (1.1) has a unique weak solution

$$(y, y') \in C([0, T], L^2(\Omega) \times H^{-1}),$$

moreover, there exists a constant $C = C(T) > 0$ such that

$$\|(y, y')\|_{L^\infty([0, T], L^2(\Omega) \times H^{-1})} \leq C \left[\|(y^0, y^1)\|_{L^2(\Omega) \times H^{-1}} + \|v\|_{L^2(\Gamma_1 \times (0, T))} \right]$$

Proof :

The theorem is a consequence of the theory of nonhomogeneous evolution equations.

- * Proof with variational method, see [2].
- * Proof with Semigroups operator, see [3].

1.3 Controllability by minimization

In this section we will give a necessary and sufficient condition for the exact controllability of (1.1) and also we will transform the controllability problem to a minimization problem, but before that we need to prove the following lemma :

Lemma 1.3.1. *For every (φ^0, φ^1) in $H_0^1(\Omega) \times L^2(\Omega)$ the following wave equation*

$$\begin{cases} \frac{\partial^2 \varphi}{\partial t^2} - \Delta \varphi = 0 & \text{in } \Omega \times (0, T), \\ \varphi = 0 & \text{in } \Gamma \times (0, T), \\ \varphi(\cdot, 0) = \varphi^0, \quad \frac{\partial \varphi}{\partial t}(\cdot, 0) = \varphi^1 & \text{in } \Omega, \end{cases} \quad (1.4)$$

has a unique solution, moreover (1.4) generates a group of isometries in $H_0^1(\Omega) \times L^2(\Omega)$.

Proof :

this lemma is a consequence of the following classic theorem :

Theorem 1.3.1. (Stone, 1930)

Let H be a Hilbert space and A be a linear operator on H with dense domain, then A generates a C_0 -group of unitary operators if and only if A is skewadjoint ($A' = -A$)

Let $w(t) = (\varphi(t), \varphi'(t))$ and the state space is $H_0^1(\Omega) \times L^2(\Omega)$, with the scalar product

$$\langle (\varphi_1, \varphi_2), (\psi_1, \psi_2) \rangle = \int_{\Omega} \nabla \varphi_1 \nabla \psi_1 + \int_{\Omega} \varphi_2 \psi_2,$$

then the corresponding norm on H given by :

$$\|(\varphi_1, \varphi_2)\|^2 = \int_{\Omega} \|\nabla \varphi_1\|^2 + \int_{\Omega} \|\varphi_2\|^2.$$

By the Poincaré inequality, this norm is equivalent to the usual norm on H .

We define the linear operator A by :

$$D(A) = H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \quad \text{and}$$

$$A = \begin{pmatrix} 0 & I \\ \Delta & 0 \end{pmatrix}.$$

Firstly the domain $D(A)$ is dense in H .

Let $(\varphi_1, \varphi_2), (\psi_1, \psi_2) \in D(A)$, then

$$\begin{aligned}
 \langle A(\varphi_1, \varphi_2), (\psi_1, \psi_2) \rangle &= \langle (\varphi_2, \Delta\varphi_1), (\psi_1, \psi_2) \rangle \\
 &= \int_{\Omega} \nabla\varphi_2 \nabla\psi_1 + \int_{\Omega} \Delta\varphi_1 \psi_2 \\
 &= \int_{\Omega} \nabla\varphi_2 \nabla\psi_1 - \int_{\Omega} \nabla\varphi_1 \nabla\psi_2 \\
 &\quad + \underbrace{\int_{\Gamma} \frac{\partial\varphi_1}{\partial n} \psi_2}_0 \\
 &= - \int_{\Omega} \nabla\varphi_1 \nabla\psi_2 - \int_{\Omega} \varphi_2 \Delta\psi_1 \\
 &= - \langle (\varphi_1, \varphi_2), (\psi_2, \Delta\psi_1) \rangle \\
 &= - \langle (\varphi_1, \varphi_2), A(\psi_1, \psi_2) \rangle.
 \end{aligned}$$

Which implies that the operator A is skewadjoint on H and thus generates a unitary C_0 -group on H by Stone's theorem.

Then if $(\varphi^0, \varphi^1) \in D(A)$ the system (1.4) has a unique strict solution by Semi-group theory, but in the case $(\varphi^0, \varphi^1) \in H \setminus D(A)$, (1.4) has a unique weak solution by follows the proof of lemma.

Proposition 1.3.1. *Assume that $T > 0$ is large enough, then there exists a constant $\hat{C} > 0$ such that :*

$$\int_0^T \int_{\Gamma_1} \left| \frac{\partial\varphi}{\partial n} \right|^2 d\sigma dt \leq \hat{C} \|(\varphi^0, \varphi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)}^2, \quad (1.5)$$

for all $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$ and φ is the solution of (1.4).

Proof :

The inequality (1.5) is a hidden regularity result, that may not be obtained by standard trace results (see, [.]).

The key result in this section is the variational characterization of the controllability of (1.1) given by the following lemma :

Lemma 1.3.2. *The initial data $(y^0, y^1) \in \mathbb{L}^2(\Omega) \times H^{-1}(\Omega)$ is controllable to zero if and only if there exists $v \in L^2(\Gamma_1 \times (0, T))$ such that :*

$$\int_0^T \int_{\Gamma_1} \frac{\partial\varphi}{\partial n} v d\sigma dt = \langle y^1, \varphi^0 \rangle_{-1,1} - \int_{\Omega} y^0 \varphi^1 dx, \quad (1.6)$$

for all $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$ and where φ is the solution of the adjoint problem (1.4)

Proof :

For $(y^0, y^1) \in D(\Omega) \times D(\Omega)$, $(\varphi^0, \varphi^1) \in D(\Omega) \times D(\Omega)$ and $v \in D(\Gamma_1 \times (0, T))$, then y and

φ are regular solutions. Multiplying the equation (1.1.1) in system (1.1) by φ and integrating, we have

$$\begin{aligned}
0 &= \int_0^T \int_{\Omega} (y'' - \Delta y) \varphi dx dt \\
&= \int_{\Omega} \int_0^T y'' \varphi dt dx - \int_0^T \int_{\Omega} \Delta y \varphi dx dt \\
&= \int_{\Omega} \left[[y' \varphi]_0^T - \int_0^T y' \varphi' dt \right] dx + \int_0^T \int_{\Omega} \nabla y \nabla \varphi dx dt \\
&= \int_{\Omega} (y'(T) \varphi(T) - y'(0) \varphi(0)) dx - \int_{\Omega} [y \varphi']_0^T dx + \int_{\Omega} \int_0^T y \varphi'' dt dx \\
&\quad - \int_0^T \int_{\Omega} \Delta \varphi y dx dt + \int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} y d\sigma dt \\
&= \int_{\Omega} (y'(T) \varphi(T) - y'(0) \varphi(0)) dx - \int_{\Omega} (y(T) \varphi'(T) - y(0) \varphi'(0)) dx + \int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} y d\sigma dt
\end{aligned}$$

Hence,

$$\begin{aligned}
\int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} y d\sigma dt &= - \int_{\Omega} (y'(T) \varphi(T) - y'(0) \varphi(0)) dx \\
&\quad + \int_{\Omega} (y(T) \varphi'(T) - y(0) \varphi'(0)) dx,
\end{aligned}$$

finally we have,

$$\begin{aligned}
\int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} y d\sigma dt &= \int_{\Omega} y^1 \varphi^0 dx - \int_{\Omega} y^0 \varphi^1 dx \\
&\quad + \int_{\Omega} y(T) \varphi'(T) dx - \int_{\Omega} y'(T) \varphi(T) dx.
\end{aligned}$$

We Know that $D(\Omega) \times D(\Omega)$ dense in $L^2(\Omega) \times H^{-1}(\Omega)$ as well as in $H_0^1(\Omega) \times L^2(\Omega)$ and $D(\Gamma_1 \times (0, T))$ dense in $L^2(\Gamma_1 \times (0, T))$. by the inequality (1.5), we have $\frac{\partial \varphi}{\partial n}|_{\Gamma_1} \in L^2(\Gamma_1) \times (0, T)$. From a density argument we deduce that for any $(y^0, y^1) \in L^2(\Omega) \times H^{-1}(\Omega)$, $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$ we have,

$$\begin{aligned}
\int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} y d\sigma dt &= \langle y^1 \varphi^0 \rangle_{-1,1} - \int_{\Omega} y^0, \varphi^1 dx \\
&\quad + \langle (\varphi(T), \varphi'(T)), (y(T), y'(T)) \rangle.
\end{aligned}$$

With $\langle \cdot, \cdot \rangle_{-1,1}$ be the duality product between $H^{-1}(\Omega)$ and $H_0^1(\Omega)$. For all $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$, $(y^0, y^1) \in L^2(\Omega) \times H^{-1}(\Omega)$ we introduce the duality product

$$\langle (\varphi^0, \varphi^1), (y^0, y^1) \rangle = \int_{\Omega} y^0 \varphi^1 - \langle y^1, \varphi^0 \rangle_{-1,1}.$$

Such that the wave equation generates a group of isometries in $H_0^1(\Omega) \times L^2(\Omega)$, then (1.6) holds if and only if (y^0, y^1) is controllable to zero in time $T > 0$. This completes the proof. From lemme 1.3.2, the equality (1.6) can be seen as an optimality condition for the minimants of the functional $\mathcal{J} : H_0^1(\Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$, defined by

$$\mathcal{J} = \frac{1}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \varphi}{\partial n} \right|^2 d\sigma dt + \langle (\varphi^0, \varphi^1), (y^0, y^1) \rangle, \quad (1.7)$$

where $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$ and φ is the corresponding solution of (1.4).

Before introducing the main theorem in this section we need to proof that the functional \mathcal{J} has a minimant.

Definition 1.3.1. System (1.4) is said to be observable in time $T > 0$ if there exists a positive a positive constant $C > 0$ such that

$$C \|(\varphi^0, \varphi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)}^2 \leq \int_0^T \int_{\Gamma_1} \left| \frac{\partial \varphi}{\partial n} \right|^2 d\sigma dt, \quad (1.8)$$

for all $(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)$ where φ is the solution of (1.4) with initial data (φ^0, φ^1) .

In the following we assume that there is a positive time T^* such that for any $T > T^*$ the system (1.4) is observable.

On the other hand, the functional \mathcal{J} is continuous, strictly convex and coercive.

It is easy to see that functional \mathcal{J} is continuous, now let $(\varphi^0, \varphi^1), (\psi^0, \psi^1)$ in $H_0^1(\Omega) \times L^2(\Omega)$ and $\lambda \in]0, 1[$, we have,

$$\begin{aligned} \mathcal{J}(\lambda(\varphi^0, \varphi^1) + (1 - \lambda)(\psi^0, \psi^1)) &= \frac{1}{2} \int_0^T \int_{\Gamma_1} \left| \lambda \frac{\partial \varphi}{\partial n} + (1 - \lambda) \frac{\partial \psi}{\partial n} \right|^2 d\sigma dt \\ &+ \langle \lambda(\varphi^0, \varphi^1) + (1 - \lambda)(\psi^0, \psi^1), (y^0, y^1) \rangle \\ &= \lambda \mathcal{J}((\varphi^0, \varphi^1)) + (1 - \lambda) \mathcal{J}((\psi^0, \psi^1)) \\ &- \frac{\lambda(1 - \lambda)}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \varphi}{\partial n} - \frac{\partial \psi}{\partial n} \right|^2 d\sigma dt. \end{aligned}$$

Using the observability inequation, we obtain,

$$\int_0^T \int_{\Gamma_1} \left| \frac{\partial}{\partial n}(\varphi - \psi) \right|^2 d\sigma dt \geq C \|(\varphi^0 - \psi^0, \varphi^1 - \psi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)}^2,$$

then if $(\varphi_0, \varphi_1) \neq (\psi_0, \psi_1)$, then,

$$\mathcal{J}(\lambda(\varphi^0, \varphi^1) + (1 - \lambda)(\psi^0, \psi^1)) < \lambda \mathcal{J}((\varphi^0, \varphi^1)) + (1 - \lambda) \mathcal{J}((\psi^0, \psi^1)).$$

Hence \mathcal{J} is strictly convex.

For the coercivity of the the functional \mathcal{J} , we have,

$$\begin{aligned}
\mathcal{J}((\varphi^0, \varphi^1)) &\geq \frac{1}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \varphi}{\partial n} \right|^2 d\sigma dt - \|(\varphi^0, \varphi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)} \| (y^0, y^1) \|_{L^2(\Omega) \times H^{-1}(\Omega)} \\
&\geq \frac{C}{2} \|(\varphi^0, \varphi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)}^2 - \|(\varphi^0, \varphi^1)\|_{H_0^1(\Omega) \times L^2(\Omega)} \| (y^0, y^1) \|_{L^2(\Omega) \times H^{-1}(\Omega)},
\end{aligned}$$

then $\lim_{\|(\varphi^0, \varphi^1)\| \rightarrow +\infty} \mathcal{J}((\varphi^0, \varphi^1)) = \infty$.

We conclude that the functional \mathcal{J} has a unique minimizer $(\hat{\varphi}^0, \hat{\varphi}^1)$ in $H_0^1(\Omega) \times L^2(\Omega)$, we have,

Theorem 1.3.2. *Let (y^0, y^1) in $L^2(\Omega) \times H^{-1}(\Omega)$ and $(\hat{\varphi}^0, \hat{\varphi}^1)$ in $H_0^1(\Omega) \times L^2(\Omega)$ be the unique minimizer of the functional \mathcal{J} , then the function \hat{v} defined on $\Gamma_1 \times (0, T)$ by :*

$$\hat{v}(x, t) = \frac{\partial \hat{\varphi}}{\partial n}(x, t), \quad (x, t) \in \Gamma_1 \times (0, T),$$

is a control which leads (y^0, y^1) to zero in time $T > 0$.

Proof :

The functional \mathcal{J} achieves its minimum at $(\hat{\varphi}^0, \hat{\varphi}^1)$, then

$$\lim_{h \rightarrow 0} \frac{1}{h} [\mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1) + h(\varphi^0, \varphi^1)) - \mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1))] = 0, \quad (1.9)$$

for all (φ^0, φ^1) in $H_0^1(\Omega) \times L^2(\Omega)$ where φ is the solution of (1.4) with initial data (φ^0, φ^1) . On the other hand, we have,

$$\begin{aligned}
\mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1) + h(\varphi^0, \varphi^1)) - \mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1)) &= \frac{1}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \hat{\varphi}}{\partial n} + h \frac{\partial \varphi}{\partial n} \right|^2 d\sigma dt + \langle (\hat{\varphi}^0, \hat{\varphi}^1) + h(\varphi^0, \varphi^1), (y^0, y^1) \rangle \\
&\quad - \frac{1}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \hat{\varphi}}{\partial n} \right|^2 d\sigma dt - \langle (\hat{\varphi}^0, \hat{\varphi}^1), (y^0, y^1) \rangle,
\end{aligned}$$

hence,

$$\begin{aligned}
\frac{1}{h} [\mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1) + h(\varphi^0, \varphi^1)) - \mathcal{J}((\hat{\varphi}^0, \hat{\varphi}^1))] &= \frac{h}{2} \int_0^T \int_{\Gamma_1} \left| \frac{\partial \varphi}{\partial n} \right|^2 d\sigma dt + \int_0^T \int_{\Gamma_1} \frac{\partial \hat{\varphi}}{\partial n} \frac{\partial \varphi}{\partial n} d\sigma dt \\
&\quad + \langle (\varphi^0, \varphi^1), (y^0, y^1) \rangle,
\end{aligned}$$

and from (1.8) we deduce that

$$\begin{aligned}
\int_0^T \int_{\Gamma_1} \frac{\partial \hat{\varphi}}{\partial n} \frac{\partial \varphi}{\partial n} d\sigma dt &= - \langle (\varphi^0, \varphi^1), (y^0, y^1) \rangle \\
&= - \int_{\Omega} y^0 \varphi^1 dx + \langle y^1, \varphi^0 \rangle_{-1,1},
\end{aligned}$$

for every (φ^0, φ^1) in $H_0^1(\Omega) \times L^2(\Omega)$.

From lemma 1.3.2, it follows that $\hat{v} = \frac{\partial \hat{\varphi}}{\partial n}|_{\Gamma_1}$ is the control for (1.1). This complete the proof. Now we can find the control of the wave equation by minimization of the functional \mathcal{J} , moreover, this control is the control of minimal L^2 -norm :

Proposition 1.3.2. *Let $\hat{v} = \frac{\partial \hat{\varphi}}{\partial n}|_{\Gamma_1}$ be the control given by minimizing the functional \mathcal{J} . If $w \in L^2(\Gamma_1 \times (0, T))$ is any other control for (1.1), then*

$$\|\hat{v}\|_{L^2(\Gamma_1 \times (0, T))} \leq \|w\|_{L^2(\Gamma_1 \times (0, T))}. \quad (1.10)$$

Proof :

Let $(\hat{\varphi}^0, \hat{\varphi}^1) \in H_0^1(\Omega) \times L^2(\Omega)$ the minimizer of the functional \mathcal{J} and w is a control function of (1.1). By taking $(\hat{\varphi}^0, \hat{\varphi}^1)$ as initial data for (1.4), lemma 1.3.2 gives,

$$\int_0^T \int_{\Gamma_1} \frac{\partial \hat{\varphi}}{\partial n} \frac{\partial \hat{\varphi}}{\partial n} d\sigma dt = - \int_{\Omega} y^0 \hat{\varphi}^1 dx + \langle y^1, \hat{\varphi}^0 \rangle_{-1,1},$$

hence,

$$\|\hat{v}\|_{L^2(\Gamma_1 \times (0, T))}^2 = - \int_{\Omega} y^0 \hat{\varphi}^1 dx + \langle y^1, \hat{\varphi}^0 \rangle_{-1,1}.$$

On the other hand,

$$\int_0^T \int_{\Gamma_1} \frac{\partial \hat{\varphi}}{\partial n} w d\sigma dt = - \int_{\Omega} y^0 \hat{\varphi}^1 dx + \langle y^1, \hat{\varphi}^0 \rangle_{-1,1},$$

then,

$$\begin{aligned} \|\hat{v}\|_{L^2(\Gamma_1 \times (0, T))}^2 &= \int_0^T \int_{\Gamma_1} \hat{v} w d\sigma dt \\ &\leq \|\hat{v}\|_{L^2(\Gamma_1 \times (0, T))} \|w\|_{L^2(\Gamma_1 \times (0, T))}. \end{aligned}$$

Consequently, (1.10) is verified and the proof finishes.

Chapitre 2

Conjugate Gradient Algorithm

2.1 Variational Problem

In the previous chapter, we transferred the problem of controllability of (1.1) to a problem of minimization :

$$\min_{(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)} \mathcal{J}((\varphi^0, \varphi^1)), \quad (2.1)$$

where the functional \mathcal{J} is defined by (1.7).

Problem (2.1) can be written as follows :

$$\min_{(\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega)} \left(\frac{1}{2} a((\varphi^0, \varphi^1), (\varphi^0, \varphi^1)) - L((\varphi^0, \varphi^1)) \right), \quad (2.2)$$

where a is defined on $(H_0^1(\Omega) \times L^2(\Omega)) \times (H_0^1(\Omega) \times L^2(\Omega))$ by

$$a((\varphi^0, \varphi^1), (\widetilde{\varphi}^0, \widetilde{\varphi}^1)) = \int_0^T \int_{\Gamma_1} \frac{\partial \varphi}{\partial n} \frac{\partial \widetilde{\varphi}}{\partial n} d\sigma dt, \quad \forall (\varphi^0, \varphi^1), (\widetilde{\varphi}^0, \widetilde{\varphi}^1) \in H_0^1(\Omega) \times L^2(\Omega),$$

such that $\varphi, \widetilde{\varphi}$ respectively the solutions of (1.4) with initial data (φ^0, φ^1) and $(\widetilde{\varphi}^0, \widetilde{\varphi}^1)$, and L is defined on $H_0^1(\Omega) \times L^2(\Omega)$ by

$$\begin{aligned} L((\varphi^0, \varphi^1)) &= - \langle (\varphi^0, \varphi^1), (y^0, y^1) \rangle \\ &= - \int_{\Omega} y^0 \varphi^1 + \langle y^1, \varphi^0 \rangle_{-1,1}, \end{aligned}$$

for all (φ^0, φ^1) in $H_0^1(\Omega) \times L^2(\Omega)$ and (y^0, y^1) in $L^2(\Omega) \times H^{-1}(\Omega)$. we have,

Lemma 2.1.1. ♦ *The operator a is a bilinear form, continuous and $H_0^1(\Omega) \times L^2(\Omega)$ -elliptic. ♦ For all $(y^0, y^1) \in L^2(\Omega) \times H^{-1}(\Omega)$, we have $L \in (H_0^1(\Omega) \times L^2(\Omega))'$.*

Proof :

- it easy to proof that L is linear continuous and a is bilinear
- The continuity of a follows from the inequality (1.5) and the coercivity follows from the inequality of observability (1.8).

Moreover the bilinear form a is symmetric, by follows and with the theorem of Lax-Milgram

the problem (2.2) reads as follows :

$$\begin{cases} \text{Find } (\widehat{\varphi}^0, \widehat{\varphi}^1) \in H_0^1(\Omega) \times L^2(\Omega) \text{ such that} \\ a((\widehat{\varphi}^0, \widehat{\varphi}^1), (\varphi^0, \varphi^1)) = L((\varphi^0, \varphi^1)), \quad \forall (\varphi^0, \varphi^1) \in H_0^1(\Omega) \times L^2(\Omega). \end{cases} \quad (2.3)$$

2.2 General conjugate gradient algorithm

Let H a Hilbert space, a a continuous, symmetric and coercive bilinear form on $H \times H$, and L a continuous linear form on H , the variational problem (2.3) is a particular case of the following general variational problem :

$$\begin{cases} \text{Find } u \in H \text{ such that} \\ a(u, v) = L(v), \quad \forall v \in H. \end{cases} \quad (2.4)$$

With the above hypotheses, problem (2.4) has a unique solution.

For the numerical solution of (2.4) we need the following algorithm, say the conjugate gradient algorithm :

(1) $u^{(0)}$: any arbitrarily vector in H ;

(2) solve

$$\begin{cases} \widetilde{u}^{(0)} \in H \\ \langle \widetilde{u}^{(0)}, v \rangle_H = a(u^{(0)}, v) - L(v), \quad \forall v \in H. \end{cases} \quad (2.5)$$

(3) • If $\widetilde{u}^{(0)}$ is small ($\frac{\|\widetilde{u}^{(0)}\|}{\|u^{(0)}\|} < \epsilon$), take $u = u^{(0)}$;
 • If not, set $\check{u}^{(0)} = \widetilde{u}^{(0)}$;

Assuming that $u^{(n)}$, $\widetilde{u}^{(n)}$, $\check{u}^{(n)}$ are known, compute $u^{(n+1)}$, $\widetilde{u}^{(n+1)}$, $\check{u}^{(n+1)}$:

(4) $\rho_n = \frac{\|\widetilde{u}^{(n)}\|^2}{a(\check{u}^{(n)}, \check{u}^{(n)})}$;

(5) $u^{(n+1)} = u^{(n)} - \rho_n \check{u}^{(n)}$;

(6) solve

$$\begin{cases} \widetilde{u}^{(n+1)} \in H \\ \langle \widetilde{u}^{(n+1)}, v \rangle_H = \langle \widetilde{u}^{(n)}, v \rangle_H - \rho_n a(\check{u}^{(n)}, v), \quad \forall v \in H. \end{cases} \quad (2.6)$$

(7) • If $\widetilde{u}^{(n+1)}$ is small ($\frac{\|\widetilde{u}^{(n+1)}\|}{\|\widetilde{u}^{(0)}\|} < \epsilon$), take $u = u^{(n+1)}$;

• If not,

★ $\gamma_n = \frac{\|\widetilde{u}^{(n+1)}\|^2}{\|\widetilde{u}^{(n)}\|^2}$;

★ $\check{u}^{(n+1)} = \widetilde{u}^{(n+1)} + \gamma_n \check{u}^{(n)}$;

(8) $n = n + 1$ and go to (4) ;

2.3 Application of conjugate gradient algorithm

Let us now apply the general conjugate gradient algorithm to the solution of (2.3).

(1) $(\varphi_0^0, \varphi_0^1) \in H_0^1(\Omega) \times L^2(\Omega) = H$: Initialization ;

(2) solve

$$\begin{cases} (\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1) \in H \\ \langle (\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1), (\varphi^0, \varphi^1) \rangle_H = a((\varphi_0^0, \varphi_0^1), (\varphi^0, \varphi^1)) - L((\varphi^0, \varphi^1)), \quad \forall (\varphi^0, \varphi^1) \in H. \end{cases} \quad (2.7)$$

Consider the following non-homogeneous backward wave equation :

$$\begin{cases} \frac{\partial^2 \psi_0}{\partial t^2} - \Delta \psi_0 = 0 & \text{in } \Omega \times (0, T), \\ \psi_0 = \frac{\partial \varphi_0}{\partial n} & \text{in } \Gamma_1 \times (0, T), \\ \psi_0 = 0 & \text{in } \Gamma_2 \times (0, T), \\ \psi_0(., T) = 0, \quad \frac{\partial \psi_0}{\partial t}(., T) = 0 & \text{in } \Omega. \end{cases}$$

From the lemma 1.3.2, we have,

$$\int_0^T \int_{\Gamma_1} \frac{\partial \varphi_0}{\partial n} \frac{\partial \varphi}{\partial n} d\sigma dt = \langle \psi_0'(0), \varphi^0 \rangle_{-1,1} - \int_{\Omega} \psi_0(0) \varphi^1 dx,$$

for every $(\varphi^0, \varphi^1) \in H$, then we obtain,

$$\begin{aligned} \int_{\Omega} \nabla \widetilde{\varphi}_0^0 \nabla \varphi^0 dx + \int_{\Omega} \widetilde{\varphi}_0^1 \varphi^1 dx &= \langle \psi_0'(0), \varphi^0 \rangle_{-1,1} - \int_{\Omega} \psi_0(0) \varphi^1 dx \\ &+ \int_{\Omega} y^0 \varphi^1 dx - \langle y^1, \varphi^0 \rangle_{-1,1}. \end{aligned}$$

Hence,

$$\langle -\Delta \widetilde{\varphi}_0^0, \varphi^0 \rangle_{-1,1} - \langle \psi_0'(0) - y^1, \varphi^0 \rangle_{-1,1} = \int_{\Omega} (y^0 - \psi_0(0) - \widetilde{\varphi}_0^1) \varphi^1 dx.$$

Finally,

$$\langle (\varphi^0, \varphi^1), (y^0 - \psi_0(0) - \widetilde{\varphi}_0^1, -\Delta \widetilde{\varphi}_0^0 - (\psi_0'(0) - y^1)) \rangle = 0.$$

Its follows : (2)

$$\begin{cases} -\Delta \widetilde{\varphi}_0^0 = \psi_0'(0) - y^1, \\ \widetilde{\varphi}_0^1 = y^0 - \psi_0(0). \end{cases} \quad (2.8)$$

(3) • If $(\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1)$ is small , take $(\widehat{\varphi}^0, \widehat{\varphi}^1) = (\varphi_0^0, \varphi_0^1)$;

• If not, set $(\check{\varphi}_0^0, \check{\varphi}_0^1) = (\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1)$.

Assuming that $(\varphi_n^0, \varphi_n^1)$, $(\widetilde{\varphi}_n^0, \widetilde{\varphi}_n^1)$, $(\check{\varphi}_n^0, \check{\varphi}_n^1)$ and φ_n , ψ_n are known, compute $(\varphi_{n+1}^0, \varphi_{n+1}^1)$, $(\widetilde{\varphi}_{n+1}^0, \widetilde{\varphi}_{n+1}^1)$, $(\check{\varphi}_{n+1}^0, \check{\varphi}_{n+1}^1)$, φ_{n+1} , ψ_{n+1} .

we knew that the form linear $(\varphi^0, \varphi^1) \in H \mapsto a((\check{\varphi}_n^0, \check{\varphi}_n^1), (\varphi^0, \varphi^1))$ is continuous, then by Riesz's theorem there exists unique $(\underline{\varphi}_n^0, \underline{\varphi}_n^1)$ in H , such that

$$a((\check{\varphi}_n^0, \check{\varphi}_n^1), (\varphi^0, \varphi^1)) = \langle (\underline{\varphi}_n^0, \underline{\varphi}_n^1), (\varphi^0, \varphi^1) \rangle, \quad \forall (\varphi^0, \varphi^1) \in H.$$

Like the previous case, we can find $(\underline{\varphi}_n^0, \underline{\varphi}_n^1)$ by :

(4) solve

$$\begin{cases} \frac{\partial^2 \check{\varphi}_n}{\partial t^2} - \Delta \check{\varphi}_n = 0 & \text{in } \Omega \times (0, T), \\ \check{\varphi}_n = 0 & \text{in } \Gamma \times (0, T), \\ \check{\varphi}_n(., 0) = \check{\varphi}_n^0, \quad \frac{\partial \check{\varphi}_n}{\partial t}(., 0) = \check{\varphi}_n^1 & \text{in } \Omega, \end{cases}$$

and then

$$\begin{cases} \frac{\partial^2 \check{\psi}_n}{\partial t^2} - \Delta \check{\psi}_n = 0 & \text{in } \Omega \times (0, T), \\ \check{\psi}_n = \frac{\partial \check{\varphi}_n}{\partial n} & \text{in } \Gamma_1 \times (0, T), \\ \check{\psi}_n = 0 & \text{in } \Gamma_2 \times (0, T), \\ \check{\psi}_n(., T) = 0, \quad \frac{\partial \check{\psi}_n}{\partial t}(., T) = 0 & \text{in } \Omega. \end{cases}$$

Compute now $(\underline{\varphi}_n^0, \underline{\varphi}_n^1) \in H$ by :

$$\begin{cases} -\Delta \underline{\varphi}_n^0 = \check{\psi}_n'(0) & \text{in } \Omega \\ \underline{\varphi}_n^1 = -\check{\psi}_n(0). \end{cases} \quad (2.9)$$

The other steps of the general algorithm are easy to adapt. Now we give the complete algorithm to solve the system (2.3) :

(1) $(\varphi_0^1, \varphi_0^1) \in H_0^1(\Omega) \times L^2(\Omega) = H$ are given ;

(2) solve then

$$\begin{cases} \frac{\partial^2 \varphi_0}{\partial t^2} - \Delta \varphi_0 = 0 & \text{in } \Omega \times (0, T), \\ \varphi_0 = 0 & \text{in } \Gamma \times (0, T), \\ \varphi_0(., T) = 0, \quad \frac{\partial \varphi_0}{\partial t}(., T) = 0 & \text{in } \Omega, \end{cases}$$

and

$$\begin{cases} \frac{\partial^2 \psi_0}{\partial t^2} - \Delta \psi_0 = 0 & \text{in } \Omega \times (0, T), \\ \psi_0 = \frac{\partial \varphi_0}{\partial n} & \text{in } \Gamma_1 \times (0, T), \\ \psi_0 = 0 & \text{in } \Gamma_2 \times (0, T), \\ \psi_0(., T) = 0, \quad \frac{\partial \psi_0}{\partial t}(., T) = 0 & \text{in } \Omega. \end{cases}$$

(3) **Compute** $(\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1) \in H$ **by**

$$\begin{cases} -\Delta \widetilde{\varphi}_0^0 = \psi_0'(0) - y^1, \\ \widetilde{\varphi}_0^0 = 0 \quad \text{in } \Gamma, \end{cases}$$

and

$$\widetilde{\varphi}_0^1 = y^0 - \psi_0(0).$$

(4) • **If** $(\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1)$ **is small** , **take** $(\widehat{\varphi}^0, \widehat{\varphi}^1) = (\varphi_0^0, \varphi_0^1)$;

• **If not**, **set** $(\check{\varphi}_0^0, \check{\varphi}_0^1) = (\widetilde{\varphi}_0^0, \widetilde{\varphi}_0^1)$.

Assuming that $(\varphi_n^0, \varphi_n^1)$, $(\widetilde{\varphi}_n^0, \widetilde{\varphi}_n^1)$, $(\check{\varphi}_n^0, \check{\varphi}_n^1)$ **and** φ_n , ψ_n **are known, compute** $(\varphi_{n+1}^0, \varphi_{n+1}^1)$, $(\widetilde{\varphi}_{n+1}^0, \widetilde{\varphi}_{n+1}^1)$, $(\check{\varphi}_{n+1}^0, \check{\varphi}_{n+1}^1)$, φ_{n+1} , ψ_{n+1} .

Descent :

(5) **Solve**

$$\begin{cases} \frac{\partial^2 \check{\varphi}_n}{\partial t^2} - \Delta \check{\varphi}_n = 0 & \text{in } \Omega \times (0, T), \\ \check{\varphi}_n = 0 & \text{in } \Gamma \times (0, T), \\ \check{\varphi}_n(\cdot, 0) = \check{\varphi}_n^0, \quad \frac{\partial \check{\varphi}_n}{\partial t}(\cdot, 0) = \check{\varphi}_n^1 & \text{in } \Omega, \end{cases}$$

and then

$$\begin{cases} \frac{\partial^2 \check{\psi}_n}{\partial t^2} - \Delta \check{\psi}_n = 0 & \text{in } \Omega \times (0, T), \\ \check{\psi}_n = \frac{\partial \check{\varphi}_n}{\partial n} & \text{in } \Gamma_1 \times (0, T), \\ \check{\psi}_n = 0 & \text{in } \Gamma_2 \times (0, T), \\ \check{\psi}_n(\cdot, T) = 0, \quad \frac{\partial \check{\psi}_n}{\partial t}(\cdot, T) = 0 & \text{in } \Omega. \end{cases}$$

(6) **Compute** $(\underline{\varphi}_n^0, \underline{\varphi}_n^1) \in H$ **by : solve**

$$\begin{cases} -\Delta \underline{\varphi}_n^0 = \check{\psi}_n'(0) & \text{in } \Omega, \\ \underline{\varphi}_n^0 = 0 & \text{in } \Gamma, \end{cases}$$

and

$$\underline{\varphi}_n^1 = -\check{\psi}_n(0).$$

(7) **Compute** ρ_n **by :**

$$\left\{ \begin{aligned} \rho_n &= \frac{\|(\widetilde{\varphi}_n^0, \widetilde{\varphi}_n^1)\|^2}{a((\check{\varphi}_n^0, \check{\varphi}_n^1), (\check{\varphi}_n^0, \check{\varphi}_n^1))}, \\ &= \frac{\|(\widetilde{\varphi}_n^0, \widetilde{\varphi}_n^1)\|^2}{\langle \check{\psi}_n'(0), \check{\varphi}_n^0 \rangle_{-1,1} - \int_{\Omega} \check{\psi}_n(0) \check{\varphi}_n^1 dx}, \\ &= \frac{\int_{\Omega} \|\nabla \widetilde{\varphi}_n^0\|^2 + \int_{\Omega} \|\widetilde{\varphi}_n^1\|^2}{\int_{\Omega} \nabla \underline{\varphi}_n^0 \nabla \check{\varphi}_n^0 dx + \int_{\Omega} \underline{\varphi}_n^1 \check{\varphi}_n^1 dx}. \end{aligned} \right.$$

(8) **Once ρ_n is known, compute :**

$$\oplus (\varphi_{n+1}^0, \varphi_{n+1}^1) = (\varphi_n^0, \varphi_n^1) - \rho_n(\check{\varphi}_n^0, \check{\varphi}_n^1),$$

$$\oplus \varphi_{n+1} = \varphi_n - \rho_n \check{\varphi}_n,$$

$$\oplus \psi_{n+1} = \psi_n - \rho_n \check{\psi}_n,$$

$$\oplus (\widetilde{\varphi}_{n+1}^0, \widetilde{\varphi}_{n+1}^1) = (\widetilde{\varphi}_n^0, \widetilde{\varphi}_n^1) - \rho_n(\underline{\varphi}_n^0, \underline{\varphi}_n^1).$$

Test of the convergence and construction of the new descent direction.

(9) **If $(\widetilde{\varphi}_{n+1}^0, \widetilde{\varphi}_{n+1}^1)$ is small , take $(\widehat{\varphi}^0, \widehat{\varphi}^1) = (\varphi_{n+1}^0, \varphi_{n+1}^1)$.**

If not, compute

$$\gamma_n = \frac{\int_{\Omega} \|\nabla \widetilde{\varphi}_{n+1}^0\|^2 dx + \int_{\Omega} \|\widetilde{\varphi}_{n+1}^1\|^2 dx}{\int_{\Omega} \|\nabla \widetilde{\varphi}_n^0\|^2 dx + \int_{\Omega} \|\widetilde{\varphi}_n^1\|^2 dx},$$

and set

$$(\check{\varphi}_{n+1}^0, \check{\varphi}_{n+1}^1) = (\widetilde{\varphi}_{n+1}^0, \widetilde{\varphi}_{n+1}^1) + \gamma_n(\check{\varphi}_n^0, \check{\varphi}_n^1).$$

(10) $n = n + 1$ **and go to (5).**

Conclusion

OK

Bibliographie

- [1] T. Rossing. Springer Handbook of Acoustics. *Rossing Ed.*, 2007.