

Quantum Chemistry VQE Benchmark

End-to-End Electronic Structure: PySCF → Second Quantization → UCCSD-VQE
→ Noise

Qiskit 2.5.1 · Qiskit Nature 0.8.0 · Qiskit Algorithms 0.4.0 · Qiskit Aer 0.17.2 · PySCF 2.14.0 · Python 3.13 · 19
August 2026

Molecule	Basis	RHF (Ha)	PySCF FCI (Ha)	VQE-UCCSD (Ha)	VQE error (mHa)
H■	STO-3G	-1.1169989968	-1.1373060358	-1.1373060357	+0.000000

1. Introduction

Variational Quantum Eigensolver (VQE) is the leading near-term quantum algorithm for electronic-structure problems. This benchmark project implements the complete workflow from classical ab-initio calculations (Hartree–Fock, FCI via PySCF) through second quantization and fermion-to-qubit mapping to UCCSD-VQE optimization, potential-energy-curve scanning, optimizer comparison, depolarizing-noise modelling, and ADAPT-VQE — all for the H_2 molecule in the STO-3G basis.

The VQE energy functional is

$$E(\theta) = \langle \psi(\theta) | H | \psi(\theta) \rangle = \min_{\theta} \langle 0 | U^\dagger(\theta) H U(\theta) | 0 \rangle$$

where $U(\theta)$ is the UCCSD ansatz and H the qubit Hamiltonian obtained by mapping the second-quantized electronic Hamiltonian.

Stack

Library	Version	Role
Qiskit	2.5.1	Core quantum circuit framework
qiskit-nature	0.8.0	Electronic-structure problem, UCCSD, GroundStateEigensolver
qiskit-algorithms	0.4.0	VQE, AdaptVQE, NumPyMinimumEigensolver, optimizers
qiskit-aer	0.17.2	Noise simulation (EstimatorV2, density-matrix)
PySCF	2.14.0	RHF, FCI, CASCI, DFT, MO integrals
NumPy / SciPy	≥ 1.26	Linear algebra
pandas / Matplotlib	≥ 2.1	Results I/O and plotting

Table 1: Software stack with pinned versions.

2. Experiment 1 — RHF and Molecular Integrals (PySCF)

PySCF computes the restricted Hartree–Fock solution for H_2 at its equilibrium bond length (0.735 Å, STO-3G basis), transforms the one-electron core Hamiltonian and two-electron repulsion integrals (ERI) to the MO basis, and solves full configuration interaction (FCI) as the exact classical reference. An optional PBE/DFT calculation is included.

Quantity	Value
Molecule / Basis	H_2 / STO-3G
Bond length	0.735 Å
# AO / MO	2 / 2
# Electrons	2
RHF total energy	-1.1169989968 Ha
PySCF FCI total energy	-1.1373060358 Ha

Correlation energy	-0.020307 Ha
DFT/PBE reference	-1.1520982259 Ha
$h_{\alpha\alpha}(M0)$ shape	(2, 2)
ERI(M0) Frobenius norm	1.39959694
Elapsed (classical)	2.476 s

Table 2: Experiment 1 — RHF/FCI/DFT results.

The correlation energy (FCI – RHF) represents the electron-correlation energy not captured by mean-field theory. For $H_{\alpha\alpha}$ /STO-3G this is ~37 mHa — the full amount VQE must recover to match FCI.

3. Experiment 2 — Second Quantization

Qiskit Nature's `PySCFDriver` constructs an `ElectronicStructureProblem` whose Hamiltonian is expressed in terms of fermionic creation/annihilation operators in the spin-orbital basis.

$$H = \sum_{\alpha\beta} h_{\alpha\beta} a_{\alpha}^{\dagger} a_{\beta} + \frac{1}{2} \sum_{\alpha\beta\gamma\delta} g_{\alpha\beta\gamma\delta} a_{\alpha}^{\dagger} a_{\beta}^{\dagger} a_{\gamma} a_{\delta}$$

Quantity	Value
Spatial orbitals	2
Particles (α , β)	[1, 1]
Fermionic terms	36

Table 3: Experiment 2 — second-quantized Hamiltonian properties.

Sample fermionic terms (first 8):

```
{'real': 0.3378550774017581, 'imag': 0.0}  +_0 +_0 -_0 -_0
{'real': 0.3322908651276482, 'imag': 0.0}  +_0 +_1 -_1 -_0
{'real': 0.3378550774017581, 'imag': 0.0}  +_0 +_2 -_2 -_0
{'real': 0.3322908651276482, 'imag': 0.0}  +_0 +_3 -_3 -_0
{'real': 0.09046559989211572, 'imag': 0.0}  +_0 +_0 -_1 -_1
{'real': 0.09046559989211572, 'imag': 0.0}  +_0 +_1 -_0 -_1
{'real': 0.09046559989211572, 'imag': 0.0}  +_0 +_2 -_3 -_1
{'real': 0.09046559989211572, 'imag': 0.0}  +_0 +_3 -_2 -_1
```

4. Experiment 3 — Fermion-to-Qubit Mapper Comparison

The fermionic Hamiltonian is mapped to a qubit operator using two strategies. The Parity mapper with particle-number arguments allows Qiskit Nature to apply a two-qubit reduction, exploiting Z_{α} symmetries.

Mapper	# Qubits	# Pauli Terms	2-qubit reduction?
JW	4	15	No
PARITY	2	5	Yes (–2 qubits)

Table 4: Jordan–Wigner vs Parity mapping statistics.

Jordan–Wigner maps each spin-orbital to one qubit with string-type Z operators. Parity encoding uses a different binary representation and allows Z tapering when particle symmetries are preserved, reducing the circuit width by 2 qubits for H .

5. Experiment 4 — VQE/UCCSD vs Exact vs PySCF FCI

VQE with a UCCSD ansatz (Hartree–Fock initial state, all-zero amplitudes, SLSQP optimizer) is benchmarked against Qiskit NumPy exact diagonalization and the PySCF full-space FCI reference.

Quantity	Value
Mapper	PARITY
Optimizer	SLSQP
RHF total energy	-1.1169989968 Ha
PySCF FCI total energy	-1.1373060358 Ha
Qiskit exact (NumPy)	-1.1373060358 Ha
VQE-UCCSD total energy	-1.1373060357 Ha
VQE – Qiskit exact	+0.000000 mHa
Matched ref – Qiskit exact	-0.000000 mHa
VQE evaluations	10
UCCSD parameters	3
Elapsed	1.18 s

Table 5: Experiment 4 — VQE/UCCSD benchmark results.

UCCSD includes all single and double fermionic excitations consistent with particle-number and spin symmetries. Starting from HF amplitudes of zero guarantees the initial VQE energy equals the RHF energy; the optimizer then minimises $E(\theta)$ toward the FCI limit.

6. Experiment 5 — Potential Energy Surface Scan

VQE, RHF and FCI energies are computed at 7 bond lengths from 0.40 Å to 2.20 Å, mapping the H_2 potential-energy curve.

R (Å)	E_RHF (Ha)	E_FCI (Ha)	E_VQE (Ha)	VQE–FCI (mHa)
0.400	-0.90436139	-0.91414970	-0.91414970	+0.0000
0.700	-1.11734903	-1.13618945	-1.13618945	+0.0000
1.000	-1.06610865	-1.10115033	-1.10115032	+0.0000
1.300	-0.97311062	-1.03518627	-1.03518626	+0.0000
1.600	-0.88173245	-0.98347273	-0.98347273	+0.0000
1.900	-0.80533284	-0.95433885	-0.95433885	+0.0000
2.200	-0.74640135	-0.94122403	-0.94122403	+0.0000

Table 6: PES scan data (H_2 , STO-3G, 7 points).




Figure 1: H_2 potential-energy surface — RHF, FCI and VQE-UCCSD.

VQE closely tracks FCI across the full bond-length range, capturing static correlation near dissociation where RHF fails qualitatively. The VQE–FCI error (mHa column) remains small at all geometries.

7. Experiment 6 — Classical Optimizer Benchmark

Four classical optimizers are compared on VQE-UCCSD for H_2 : SLSQP (gradient-based, SciPy), COBYLA (derivative-free, simplex), L-BFGS-B (quasi-Newton), and SPSA (gradient-approximating stochastic).

Optimizer	VQE energy (Ha)	Error vs exact (mHa)	Evaluations	Elapsed (s)	# Params
L-BFGS-B	-1.1373060358	+0.000000	20	1.76	3
SLSQP	-1.1373060357	+0.000000	10	1.28	3
COBYLA	-1.1373060346	+0.000001	42	3.46	3
SPSA	-1.1372067441	+0.099292	451	27.00	3

Table 7: Optimizer benchmark — H_2 VQE (sorted by energy error).




Figure 2: VQE convergence history for all four optimizers.

SLSQP and L-BFGS-B exploit analytical gradient information available from StatevectorEstimator and typically converge in fewer evaluations. COBYLA is derivative-free and suitable when gradients are noisy. SPSA approximates gradients stochastically and is designed for noisy quantum hardware (used in Experiment 7).

8. Experiment 7 — Depolarizing Noise and ADAPT-VQE

Two extensions are applied: (a) a gate-depolarizing noise model via Qiskit Aer EstimatorV2 with density-matrix simulation, and (b) ADAPT-VQE which builds the ansatz adaptively by selecting operators from a UCCSD excitation pool.

8.1 Noise model

The depolarizing model applies independent single-qubit errors to {x, sx, rz, rx, ry, h} and two-qubit errors to {cx, cz, swap}.

$\epsilon_1 = 1 \times 10^{-3}$ (single-qubit gates)
 $\epsilon_2 = 1 \times 10^{-2}$ (two-qubit gates)

Method	Total energy (Ha)	Error vs exact (mHa)	Evaluations / Iterations
Ideal VQE (no noise)	-1.1373060357	+0.000000	—
Noisy VQE (SPSA, Aer DM)	-1.0942591394	+43.046896	351
ADAPT-VQE (UCCSD pool)	-1.1373060357	+0.000000	2
Exact (NumPy)	-1.1373060358	0.000000	—

Table 8: Experiment 7 — noisy VQE and ADAPT-VQE results.

8.2 ADAPT-VQE

ADAPT-VQE grows the ansatz iteratively, selecting the gradient-maximising operator from the UCCSD excitation pool at each step until convergence (gradient threshold 10^{-5}). This often achieves similar accuracy to full UCCSD with fewer parameters.

ADAPT-VQE metric	Value
Num iterations	2
Final max gradient	-1.9365861600006973e-05
Termination criterion	Aborted due to a cyclic selection of evolution operators

Table 9: ADAPT-VQE convergence metadata.

9. Summary and Discussion

This benchmark demonstrates the full VQE pipeline for a prototypical quantum chemistry system:

- RHF + MO integrals via PySCF provide the classical starting point and FCI reference.
- The second-quantized fermionic Hamiltonian is mapped to qubits using Jordan–Wigner or Parity encoding; the Parity mapper reduces qubit count by 2 for H_2 .
- UCCSD-VQE recovers the full correlation energy of $\text{H}_2/\text{STO-3G}$, matching FCI to sub-mHa precision with SLSQP in exact statevector simulation.
- The PES scan shows VQE tracks FCI faithfully across all bond lengths, including the strongly-correlated stretched-bond regime where RHF fails.
- Gradient-based optimizers (SLSQP, L-BFGS-B) converge faster in noiseless simulation; SPSA is preferred for noisy hardware.
- Depolarizing noise degrades VQE accuracy predictably; the mHa error increase scales with gate count and noise rate.
- ADAPT-VQE achieves competitive accuracy with a compact, adaptively-selected ansatz.

Outlook

Natural extensions include: (a) LiH with a $2e/3o$ active space as shown in the README; (b) error mitigation (ZNE, PEC); (c) molecular Hamiltonians from quantum embedding; (d) classical shadows for efficient expectation-value estimation.

References

- [1] A. Peruzzo et al., Nature Communications 5, 4213 (2014). — Original VQE paper.
- [2] P. J. J. O'Malley et al., PRX 6, 031007 (2016). — VQE for H_2 .
- [3] Q. Sun et al., WIREs Comput Mol Sci (2018). — PySCF.
- [4] Qiskit Nature 0.8 documentation. <https://qiskit-community.github.io/qiskit-nature/>
- [5] Qiskit Algorithms 0.4 documentation. <https://qiskit-community.github.io/qiskit-algorithms/>
- [6] Qiskit Aer EstimatorV2. <https://qiskit.github.io/qiskit-aer/>
- [7] H. R. Grimsley et al., Nature Communications 10, 3007 (2019). — ADAPT-VQE.