Table 1 | Restricted mean survival time

Cohort	Endpoint	Group	ICI+CT	CT	Difference (95% CI)	P value
JUPITER-06	PFS, 12 m	B	9.33 (8.56–10.09)	8.61 (7.81–9.41)	0.71 (-0.39–1.82)	.207 a
JUPITER-06	PFS, 12 m	NB	7.30 (6.27–8.33)	8.28 (7.37–9.18)	-0.97 (-2.35–0.40)	.164 a
JUPITER-06	PFS, 12 m	B minus NB	—	—	1.69 (-0.08–3.45)	.061 b
JUPITER-06	OS, 24 m	B	20.01 (18.81–21.22)	18.98 (17.61–20.36)	1.03 (-0.80–2.85)	.270 a
JUPITER-06	OS, 24 m	NB	17.67 (15.99–19.35)	18.28 (16.70–19.86)	-0.61 (-2.91–1.70)	.607 a
JUPITER-06	OS, 24 m	B minus NB	—	—	1.63 (-1.31–4.58)	.276 b
ESCORT-1ST	PFS, 12 m	B	9.15 (8.35–9.95)	8.27 (7.51–9.04)	0.88 (-0.23–1.98)	.121 a
ESCORT-1ST	PFS, 12 m	NB	6.57 (5.63–7.51)	8.75 (7.78–9.72)	-2.18 (-3.53–-0.83)	.002 a
ESCORT-1ST	PFS, 12 m	B minus NB	—	—	3.05 (1.31–4.80)	<.001 b
ESCORT-1ST	OS, 24 m	B	20.40 (19.18–21.62)	18.79 (17.52–20.06)	1.61 (-0.15–3.38)	.073 a
ESCORT-1ST	OS, 24 m	NB	15.85 (14.15–17.55)	18.79 (17.19–20.38)	-2.94 (-5.27–-0.61)	.013 a
ESCORT-1ST	OS, 24 m	B minus NB	—	—	4.55 (1.63–7.48)	.002 b

Units: months. Treatment difference: ICI+CT minus CT.
a: within-group treatment-difference Wald test. b: B minus NB treatment-difference Wald test. All P values nominal.
JUPITER uses OOF groups; ESCORT uses frozen external groups. Greenwood plug-in variances; predictions treated as fixed.

SIMULATED DATA — workflow demonstration only