Table 2 | Fixed-time survival probabilities

Cohort	Endpoint	Group	ICI+CT	CT	Difference (95% CI)	P value
JUPITER-06	PFS, 12 m	B	61.17 (51.75–70.58)	52.48 (42.74–62.21)	8.69 (-4.85–22.23)	.209 a
JUPITER-06	PFS, 12 m	NB	38.96 (28.07–49.85)	44.30 (33.35–55.26)	-5.34 (-20.79–10.10)	.498 a
JUPITER-06	PFS, 12 m	B minus NB	—	—	14.03 (-6.51–34.58)	.181 b
JUPITER-06	OS, 24 m	B	64.08 (54.81–73.34)	52.48 (42.74–62.21)	11.60 (-1.84–25.04)	.091 a
JUPITER-06	OS, 24 m	NB	44.16 (33.06–55.25)	46.84 (35.83–57.84)	-2.68 (-18.30–12.94)	.737 a
JUPITER-06	OS, 24 m	B minus NB	—	—	14.28 (-6.33–34.89)	.174 b
ESCORT-1ST	PFS, 12 m	B	59.18 (49.45–68.91)	45.87 (36.52–55.23)	13.31 (-0.19–26.81)	.053 a
ESCORT-1ST	PFS, 12 m	NB	29.27 (19.42–39.12)	49.30 (37.67–60.92)	-20.03 (-35.27–-4.79)	.010 a
ESCORT-1ST	PFS, 12 m	B minus NB	—	—	33.34 (12.98–53.70)	.001 b
ESCORT-1ST	OS, 24 m	B	65.31 (55.88–74.73)	48.62 (39.24–58.01)	16.68 (3.38–29.98)	.014 a
ESCORT-1ST	OS, 24 m	NB	40.24 (29.63–50.86)	50.70 (39.08–62.33)	-10.46 (-26.20–5.28)	.193 a
ESCORT-1ST	OS, 24 m	B minus NB	—	—	27.14 (6.53–47.75)	.010 b

Units: percentage points (survival rates in %). Treatment difference: ICI+CT minus CT.
a: within-group treatment-difference Wald test. b: B minus NB treatment-difference Wald test. All P values nominal.
JUPITER uses OOF groups; ESCORT uses frozen external groups. Greenwood plug-in variances; predictions treated as fixed.

SIMULATED DATA — workflow demonstration only