Table S1 | Clinical characteristics by treatment arm

Characteristic	JUPITER-06 CT	JUPITER-06 ICI+CT	ESCORT-1ST CT	ESCORT-1ST ICI+CT	Cohort comparison P value
Patients	180	180	180	180	NE
Age, median (IQR)	58.0 (49.0–69.0)	58.5 (47.0–68.0)	58.0 (48.0–69.0)	61.0 (50.0–71.0)	.391
sex: Male	143 (79.4%)	150 (83.3%)	155 (86.1%)	139 (77.2%)	.923
sex: Female	37 (20.6%)	30 (16.7%)	25 (13.9%)	41 (22.8%)	.923
sex: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
ECOG: 0	95 (52.8%)	93 (51.7%)	79 (43.9%)	92 (51.1%)	.205
ECOG: 1	85 (47.2%)	87 (48.3%)	101 (56.1%)	88 (48.9%)	.205
ECOG: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
TPS: <1%	79 (43.9%)	71 (39.4%)	82 (45.6%)	84 (46.7%)	.220
TPS: >=1%	93 (51.7%)	99 (55.0%)	86 (47.8%)	90 (50.0%)	.220
TPS: Missing	8 (4.4%)	10 (5.6%)	12 (6.7%)	6 (3.3%)	—
PFS events	166 (92.2%)	138 (76.7%)	160 (88.9%)	147 (81.7%)	NE
OS events	132 (73.3%)	123 (68.3%)	137 (76.1%)	124 (68.9%)	NE

P values compare the two cohorts overall. Age: Mann–Whitney; categorical variables: Pearson, missing excluded.
Continuous variables: median (IQR). Categorical percentages use all patients in the column. Repeated category P values are one omnibus test.

SIMULATED DATA — workflow demonstration only