Table S2 | JUPITER-06 inclusion comparison

Characteristic	Full trial	Image included	Not included	Included vs excluded P value
Patients	390	360	30	—
Mapped analysis slides	360	360	0	—
Age, median (IQR)	59.0 (49.0–69.0)	58.0 (49.0–69.0)	63.0 (54.5–69.5)	.108
sex: Male	320 (82.1%)	293 (81.4%)	27 (90.0%)	.238
sex: Female	70 (17.9%)	67 (18.6%)	3 (10.0%)	.238
sex: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
ECOG: 0	206 (52.8%)	188 (52.2%)	18 (60.0%)	.412
ECOG: 1	184 (47.2%)	172 (47.8%)	12 (40.0%)	.412
ECOG: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
TPS: <1%	163 (41.8%)	150 (41.7%)	13 (43.3%)	.920
TPS: >=1%	208 (53.3%)	192 (53.3%)	16 (53.3%)	.920
TPS: Missing	19 (4.9%)	18 (5.0%)	1 (3.3%)	—
PFS events	331 (84.9%)	304 (84.4%)	27 (90.0%)	—
OS events	279 (71.5%)	255 (70.8%)	24 (80.0%)	—

Included WSI: 360. Original trial counts require all randomized patients, including those without images.
Age: two-sided Mann–Whitney test. Categorical variables: nominal Pearson test, missing excluded. Repeated category P values refer to one omnibus test.

SIMULATED DATA — workflow demonstration only