Table S3 | ESCORT-1ST inclusion comparison

Characteristic	Full trial	Image included	Not included	Included vs excluded P value
Patients	390	360	30	—
Mapped analysis slides	360	360	0	—
Age, median (IQR)	60.0 (49.0–70.0)	60.0 (49.0–70.0)	61.5 (47.0–72.8)	.889
sex: Male	322 (82.6%)	294 (81.7%)	28 (93.3%)	.106
sex: Female	68 (17.4%)	66 (18.3%)	2 (6.7%)	.106
sex: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
ECOG: 0	187 (47.9%)	171 (47.5%)	16 (53.3%)	.539
ECOG: 1	203 (52.1%)	189 (52.5%)	14 (46.7%)	.539
ECOG: Missing	0 (0.0%)	0 (0.0%)	0 (0.0%)	—
TPS: <1%	179 (45.9%)	166 (46.1%)	13 (43.3%)	.701
TPS: >=1%	192 (49.2%)	176 (48.9%)	16 (53.3%)	.701
TPS: Missing	19 (4.9%)	18 (5.0%)	1 (3.3%)	—
PFS events	333 (85.4%)	307 (85.3%)	26 (86.7%)	—
OS events	284 (72.8%)	261 (72.5%)	23 (76.7%)	—

Included WSI: 360. Original trial counts require all randomized patients, including those without images.
Age: two-sided Mann–Whitney test. Categorical variables: nominal Pearson test, missing excluded. Repeated category P values refer to one omnibus test.

SIMULATED DATA — workflow demonstration only