Table S4 | Fixed-candidate ablations

Cohort	Model	Endpoint	Terms	B / NB	B treatment HR (95% CI)	NB treatment HR (95% CI)	HR B / HR NB (95% CI)	Binary P	Continuous HR (95% CI)	Continuous P
JUPITER-06	Full	PFS	179	204 / 156	0.525 (0.386–0.714)	1.051 (0.753–1.467)	0.499 (0.317–0.786)	.003	1.132 (1.049–1.220)	.001
JUPITER-06	Full	OS	179	204 / 156	0.721 (0.514–1.010)	1.014 (0.716–1.437)	0.711 (0.437–1.154)	.167	1.086 (1.005–1.174)	.037
JUPITER-06	No α	PFS	90	179 / 181	0.452 (0.320–0.640)	1.116 (0.830–1.503)	0.405 (0.257–0.640)	<.001	1.285 (1.156–1.428)	<.001
JUPITER-06	No α	OS	90	179 / 181	0.648 (0.446–0.942)	1.045 (0.759–1.439)	0.620 (0.379–1.015)	.057	1.151 (1.036–1.278)	.009
JUPITER-06	No βT	PFS	178	167 / 193	0.495 (0.347–0.706)	0.980 (0.728–1.318)	0.506 (0.318–0.803)	.004	1.139 (1.054–1.232)	.001
JUPITER-06	No βT	OS	178	167 / 193	0.722 (0.491–1.062)	1.002 (0.734–1.369)	0.721 (0.439–1.184)	.196	1.086 (1.001–1.177)	.047
JUPITER-06	γ only	PFS	89	162 / 198	0.404 (0.276–0.589)	1.089 (0.816–1.453)	0.370 (0.230–0.596)	<.001	1.297 (1.164–1.446)	<.001
JUPITER-06	γ only	OS	89	162 / 198	0.568 (0.377–0.854)	1.097 (0.810–1.484)	0.518 (0.311–0.860)	.011	1.153 (1.036–1.284)	.009
ESCORT-1ST	Full	PFS	179	207 / 153	0.554 (0.405–0.758)	1.408 (1.005–1.973)	0.393 (0.248–0.624)	<.001	1.225 (1.118–1.342)	<.001
ESCORT-1ST	Full	OS	179	207 / 153	0.582 (0.415–0.815)	1.273 (0.887–1.826)	0.457 (0.279–0.749)	.002	1.212 (1.094–1.343)	<.001
ESCORT-1ST	No α	PFS	90	176 / 184	0.510 (0.361–0.722)	1.338 (0.988–1.813)	0.381 (0.240–0.605)	<.001	1.316 (1.172–1.477)	<.001
ESCORT-1ST	No α	OS	90	176 / 184	0.511 (0.349–0.748)	1.258 (0.908–1.742)	0.406 (0.246–0.671)	<.001	1.289 (1.134–1.466)	<.001
ESCORT-1ST	No βT	PFS	178	185 / 175	0.500 (0.357–0.701)	1.380 (1.006–1.892)	0.363 (0.228–0.576)	<.001	1.244 (1.132–1.368)	<.001
ESCORT-1ST	No βT	OS	178	185 / 175	0.511 (0.355–0.735)	1.284 (0.917–1.797)	0.398 (0.243–0.653)	<.001	1.228 (1.104–1.367)	<.001
ESCORT-1ST	γ only	PFS	89	170 / 190	0.522 (0.368–0.742)	1.243 (0.921–1.678)	0.420 (0.265–0.667)	<.001	1.328 (1.179–1.496)	<.001
ESCORT-1ST	γ only	OS	89	170 / 190	0.515 (0.348–0.761)	1.207 (0.875–1.664)	0.427 (0.257–0.708)	<.001	1.302 (1.142–1.484)	<.001

All retained terms refitted to development PFS, ridge .01. Same candidate and fold representations; no candidate reselection.
Binary: Altman–Bland. Continuous: treatment × raw log HR(z), adjusted for main effects. Internal Cox evaluation stratifies by fold.
Terms are coefficient counts, not effective degrees of freedom. Nominal robust inference conditions on predictions; no between-model superiority test.

SIMULATED DATA — workflow demonstration only