

Math 230B Lecture Notes

Lance Remigio

October 9, 2025

Contents

1	Week 5	1
2	Week 6	1
3	Week 7	7
3.1	Kernel and Image	7

1 Week 5

Example ($SL(n, \mathbb{F})$). Suppose $\mathbb{F} = \mathbb{R}$ (or $\mathbb{C}$). We claim that

$$SL(n, \mathbb{F}) = \{A \in GL(n, \mathbb{F}) : \det A = 1\}$$

is a Lie subgroup of $GL(n, \mathbb{F})$.

Indeed, we have that

- $SL(n, \mathbb{F})$ is closed under multiplication, indeed, if $A, B \in SL(n, \mathbb{F})$, then

$$\det(AB) = (\det A)(\det B) = 1 \cdot 1 = 1.$$

So, AB is invertible and $\det(AB) = 1$. This implies that

$$AB \in SL(n, \mathbb{F}).$$

- $SL(n, \mathbb{F})$ contains the inverse of each of its elements. Indeed, suppose $A \in SL(n, \mathbb{F})$, then A is invertible and

$$\det(A^{-1}) = \frac{1}{\det A} = \frac{1}{1} = 1.$$

So, $A^{-1} \in SL(n, \mathbb{F})$.

From the above, we can conclude that $SL(n, \mathbb{F})$ is a subgroup of $GL(n, \mathbb{F})$.

2 Week 6

Example (Special Orthogonal Group). Note that

$$\begin{aligned} SO(n) &= \{A \in M(n, \mathbb{R}) : A^T A = I \text{ and } \det A > 0\} \\ &= \{A \in M(n, \mathbb{R}) : A^T A = I \text{ and } \det A = 1\} \end{aligned}$$

We will prove that $SO(n)$ is a subgroup of $O(n)$.

- Note that $SO(n)$ is closed under multiplication. Indeed, suppose A and B are in $SO(n)$. Our

goal is to show that $AB \in \text{SO}(n)$. Then we have

$$A, B \in \text{SO}(n) \implies A, B \in \text{O}(n) \implies AB \in \text{O}(n).$$

Since $A, B \in \text{SO}(n)$, it follows that $\det A = 1$ and $\det B = 1$. Hence,

$$\det AB = (\det A)(\det B) = 1 \cdot 1 = 1.$$

- Also observe that $\text{SO}(n)$ contains the inverse of each of its elements. Indeed, if $A \in \text{SO}(n)$, then

$$A \in \text{SO}(n) \implies A \in \text{O}(n) \implies A^{-1} \in \text{O}(n). \quad (1)$$

Furthermore,

$$A \in \text{SO}(n) \implies \det A = 1 \implies \det A^{-1} = \frac{1}{1} = 1 \implies A^{-1} \in \text{SO}(n) \quad (2)$$

From (1) and (2), we can see that $\text{SO}(n)$ is a subgroup of $\text{O}(n)$. Now, we need to find out whether $\text{SO}(n)$ is a Lie subgroup of $\text{O}(n)$. Note that $\text{SO}(n) = \text{O}(n) \cap \det^{-1}(0, \infty)$ where $(0, \infty)$ is an open set, $\det$ is a continuous function, and the preimage of an open set will be open in $M(n, \mathbb{R})$. Hence, $\text{SO}(n)$ will be open in $\text{O}(n)$. Thus, $\text{SO}(n)$ is an embedded submanifold of $\text{O}(n)$ with dimension $\frac{n^2-n}{2}$. Thus, $\text{SO}(n)$ is a Lie subgroup of $\text{O}(n)$ with dimension $\frac{n^2-n}{2}$.

Example (Special Unitary Group). Recall that

$$\text{SU}(n) = \{A \in M(n, \mathbb{C}) : A^*A = I \text{ and } \det A = 1\}.$$

We will show that the above is also a subgroup of $\text{U}(n)$. Clearly, we can see that $\text{SU}(n) \subseteq \text{U}(n)$.

- $\text{SU}(n)$ is closed under multiplication. Indeed, suppose A and B are in $\text{SU}(n)$. Then observe that

$$A, B \in \text{SU}(n) \implies A, B \in \text{U}(n) \implies AB \in \text{U}(n) \quad (1)$$

and

$$\begin{aligned} A, B \in \text{SU}(n) &\implies \det A = 1 \text{ and } \det B = 1 \\ &\implies \det AB = 1 \\ &\implies AB \in \text{SU}(n). \end{aligned}$$

- Also, $\text{SU}(n)$ contains the inverse of each of its elements. Suppose $A \in \text{SU}(n)$. Then

$$A \in \text{SU}(n) \implies A \in \text{U}(n) \implies A^{-1} \in \text{U}(n) \quad (3)$$

and

$$A \in \text{SU}(n) \implies \det A = 1 \implies \det A^{-1} = \frac{1}{\det A} = \frac{1}{1} = 1 \quad (4)$$

Now, (3) and (4) imply that $A^{-1} \in \text{SU}(n)$.

Now, we need to find out if $\text{SU}(n)$ is a Lie subgroup of $\text{U}(n)$? Of course! Let $F : \text{U}(n) \rightarrow \mathbb{R}$ be defined as follows:

$$F(A) = \varphi.$$

where we previously proved that if $A \in \text{U}(n)$, then $|\det A| = 1$. So, there exists an angle $\varphi \in (-\pi, \pi]$ such that $\det A = e^{i\varphi}$. It can be shown that

$$\forall A \in \text{U}(n) \quad \text{rank } DF(A) = 1.$$

Thus, $\text{SU}(n) = F^{-1}(0)$ is an $n^2 - 1$ dimensional embedded submanifold of $\text{U}(n)$. Therefore, $\text{SU}(n)$ is a Lie subgroup of $\text{U}(n)$ with dimension $n^2 - 1$.

Definition (Bilinear Form). A function $B : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be a **bilinear form** on $\mathbb{R}^n$ if it satisfies the following condition:

- (1) $\forall x, y, z \in \mathbb{R}^n \quad B(x + y, z) = B(x, z) + B(y, z)$
- (2) $\forall c \in \mathbb{R} \text{ and } \forall x, y \in \mathbb{R}^n \quad B(cx, y) = cB(x, y)$
- (3) $\forall x, y, z \in \mathbb{R}^n \quad B(x, y + z) = B(x, y) + B(x, z)$
- (4) $\forall c \in \mathbb{R} \quad \forall x, y \in \mathbb{R}^n \quad B(x, cy) = cb(x, y)$

Moreover,

- The bilinear form B is said to be **symmetric** if

$$\forall x, y \in \mathbb{R}^n \quad B(x, y) = B(y, x).$$

- The bilinear form B is said to be **skew-symmetric** if

$$\forall x, y \in \mathbb{R}^n \quad B(x, y) = -B(y, x).$$

- The bilinear form B is said to be **non-degenerate** if the following statement is true

$$B(x, y) = 0 \quad \forall y \in \mathbb{R}^n \implies x = 0.$$

- A symmetric bilinear form B is said to be **positive definite** if

$$B(x, x) > 0 \quad \forall x \neq 0.$$

Example (Standard Inner Product on $\mathbb{R}^n$). The standard inner product in $\mathbb{R}^n$

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

is a symmetric positive definite bilinear form on $\mathbb{R}^n$.

For example, let's verify condition (1) for bilinearity:

$$\begin{aligned} \langle x + y, z \rangle &= \sum_{i=1}^n [(x_i + y_i)z_i] \\ &= \sum_{i=1}^n [x_i z_i + y_i z_i] \\ &= \sum_{i=1}^n x_i z_i + \sum_{i=1}^n y_i z_i \\ &= \langle x, z \rangle + \langle y, z \rangle. \end{aligned}$$

I leave you to check the other conditions yourself.

In general, a symmetric positive definite bilinear form on a real vector space V , by definition, is called an **inner product** on V .

Example (Standard Symplectic Form on $\mathbb{R}^n$). Define $w : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$ as follows:

$$\forall x, y \in \mathbb{R}^{2n} \quad w(x, y) = \sum_{j=1}^n x_j y_{n+j} - x_{n+j} y_j.$$

For example,

$$\begin{aligned} n = 1 &\implies w(x, y) = x_1y_2 + x_2y_1 \\ n = 2 &\implies w(x, y) = (x_1y_3 - x_3y_1) + (x_2y_4 - x_4y_2) \\ &\vdots \end{aligned}$$

If we let $\Omega = \begin{pmatrix} O_{n \times n} & I_{n \times n} \\ -I_{n \times n} & O_{n \times n} \end{pmatrix}$, then we can write the standard symplectic form in the following way:

$$w(x, y) = \langle x, \Omega y \rangle$$

where $\langle, \rangle$ is the standard inner product in $\mathbb{R}^{2n}$.

Proposition (Preserving the standard symplectic form on $\mathbb{R}^{2n}$). Suppose $A \in M(2n, \mathbb{R})$. We say that A preserves the standard symplectic form on $\mathbb{R}^{2n}$ if and only if $A^T \Omega A = \Omega$.

Proof. We say that $A \in M(2n, \mathbb{R})$ preserves the standard symplectic form on $\mathbb{R}^{2n}$ if and only if

$$\begin{aligned} \forall x, y \in \mathbb{R}^{2n} \quad w(Ax, Ay) = w(x, y) &\iff \langle Ax, \Omega(Ay) \rangle = \langle x, \Omega y \rangle \\ &\iff \langle x, A^T \Omega Ay \rangle = \langle x, \Omega y \rangle \\ &\iff \forall y \in \mathbb{R}^{2n} \quad A^T \Omega Ay = \Omega y \\ &\iff A^T \Omega A = \Omega. \end{aligned}$$

■

Proposition. If $A^T \Omega A = \Omega$, then A is invertible.

Proof. Suppose $A^T \Omega A = \Omega$, then we have

$$\begin{aligned} \det(A^T \Omega A) &= \det(\Omega) \\ \implies \det(A^T) \det(\Omega) \det(A) &= \det(\Omega) \\ \implies (\det(A))^2 \det(\Omega) &= \det(\Omega) \\ \implies (\det(A))^2 &= 1 \\ \implies \det(A) &= \pm 1 \\ \implies A &\text{ is invertible.} \end{aligned}$$

■

Proposition. $A^T \Omega A = \Omega$ if and only if $-\Omega A^T \Omega = A^{-1}$ where A is invertible.

Proof. We have

$$\begin{aligned} -\Omega A^T \Omega = A^{-1} &\iff -A^T \Omega = \Omega^{-1} A^{-1} \\ &\text{left multiply by } \Omega^{-1} \\ &\iff -A^T \Omega A = \Omega^{-1} \\ &\text{right multiply by } A \\ &\iff -A^T \Omega A = \Omega \\ &\text{ } \Omega^{-1} = -\Omega \\ &\iff A^T \Omega A = \Omega. \end{aligned}$$

■

Example (Real Symplectic Group). $\text{Sp}(2n, \mathbb{R}) = \{A \in M(2n, \mathbb{R}) : A^T \Omega A = \Omega\}$ is a subgroup of $\text{GL}(2n, \mathbb{R})$.

Proof. We just proved that every matrix in $\text{SP}(2n, \mathbb{R})$ is invertible. So,

$$\text{SP}(2n, \mathbb{R}) \subseteq \text{GL}(2n, \mathbb{R}).$$

- We prove that $\text{SP}(2n, \mathbb{R})$ is closed under multiplication. Suppose $A, B \in \text{SP}(2n, \mathbb{R})$. We have

$$\begin{aligned} (AB)^T \Omega (AB) &= B^T (A^T \Omega A) B \\ &= B^T \Omega B \\ &= \Omega. \end{aligned}$$

Hence, we have $AB \in \text{SP}(2n, \mathbb{R})$.

- $\text{SP}(2n, \mathbb{R})$ contains the inverse of each of its elements. Suppose $A \in \text{SP}(2n, \mathbb{R})$. We have

$$\begin{aligned} A^T \Omega A &= A \implies (A^T)^{-1} \Omega A = (A^T)^{-1} \Omega \\ &\implies \Omega = (A^T)^{-1} \Omega A^{-1} \\ &\implies \Omega = (A^{-1})^T \Omega A^{-1} \\ &\implies A^{-1} \in \text{SP}(2n, \mathbb{R}) \end{aligned}$$

From the above, we can see that $\text{SP}(2n, \mathbb{R})$ is a subgroup of $\text{GL}(2n, \mathbb{R})$. Now, we will prove that $\text{SP}(2n, \mathbb{R})$ is a Lie subgroup of $\text{GL}(2n, \mathbb{R})$. Define the map

$$G : \text{GL}(2n, \mathbb{R}) \subseteq \mathbb{R}^{4n^2} \rightarrow M(2n, \mathbb{R}) \cong \mathbb{R}^{4n^2} \text{ by } A \mapsto A^T \Omega A - \Omega.$$

Clearly, $\text{SP}(2n, \mathbb{R}) = F^{-1}(\{0\})$. It can be shown that

$$\forall A \in \text{GL}(2n, \mathbb{R}) \quad DF(A) \text{ has rank } \frac{4n^2 - 2n}{2} = 2n^2 - n.$$

Hence, $\text{SP}(2n, \mathbb{R})$ is an embedded submanifold of $\text{GL}(2n, \mathbb{R})$ with dimension $4n^2 - (2n^2 - n) = 2n^2 + n$. Thus, $\text{SP}(2n, \mathbb{R})$ is a Lie subgroup of $\text{GL}(2n, \mathbb{R})$. ■

Definition (Homomorphisms, Isomorphisms, Automorphisms). Let G and H be two groups.

- A function $\varphi : G \rightarrow H$ is said to be a **homomorphism** if it preserves group multiplication

$$\forall g_1, g_2 \in G \quad \varphi(g_1 g_2) = \varphi(g_1) \varphi(g_2).$$

- A bijective homomorphism is called an **isomorphism**. If there is an Isomorphism between G and H , we say G and H are isomorphic, and we write $G \cong H$.
- An **endomorphism** of G is a homomorphism from G to itself.
- An **automorphism** of G is an isomorphism from G to itself.

Definition (Lie Group Homomorphism). Let G and H be two Lie groups.

- A function $f : G \rightarrow H$ is said to be a **Lie group homomorphism** if

- (i) f is a group homomorphism
- (ii) f is smooth

- A function $f : G \rightarrow H$ is said to be a **Lie group isomorphism** if

- (i) f is a group isomorphism
- (ii) f is smooth

(iii) f^{-1} is smooth

If f is smooth and its inverse is smooth, then f is a diffeomorphism.

Example (The Determinant is a Lie Group Homomorphism). The determinant $\det : GL(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ is a Lie Group Homomorphism:

- (i) $\forall A, B \in GL(n, \mathbb{R})$, we have $\det(AB) = \det(A)\det(B)$. Hence, $\det$ is a group homomorphism.
- (ii) $\det A$ is a polynomial in terms of the entries of A and so differentiable. Hence, the $\det$ is smooth.

Similarly, the $\det GL(n, \mathbb{C}) \rightarrow \mathbb{C}^*$ is a Lie Group Homomorphism.

Example (Left Translation/right translation). Let G be a Lie Group and let $g \in G$. The **left translation** by g is the map

$$L_g : G \rightarrow G \quad h \mapsto gh.$$

The **right translation** by g is the map

$$R_g : G \rightarrow G \quad h \mapsto hg.$$

Notice that L_g is a composition of smooth maps and so it is smooth:

$$G \rightarrow G \times G \rightarrow G \quad h \mapsto (g, h) \mapsto gh.$$

Starting from left to right, we see that the first mapping is smooth because

- Fact 1: $F = (F_1, F_2)$ is smooth if and only if F_1 and F_2 are smooth.
- Fact 2: $h \mapsto g$ is a constant function implies that it is smooth.
- Fact 3: $h \mapsto h$ is the identity which is also smooth.

The second mapping $G \times G \rightarrow G$ is smooth because multiplication in G is smooth since G is a Lie group.

In fact, $L_g : G \rightarrow G$ is a diffeomorphism because the inverse L_g is $L_{g^{-1}}$:

$$\forall h \quad L_{g^{-1}} \circ L_g(h) = g^{-1}(gh) = h.$$

Unfortunately, L_g in general is NOT a group homomorphism; that is, in general

$$L_g(h_1 h_2) \neq L_g(h_1) L_g(h_2).$$

Similarly, the right translation map also satisfies the properties above but is NOT a group homomorphism.

Example (Conjugation by $g \in G$). Let G be a Lie group and let $g \in G$. The conjugation by g is the map:

$$C_g : G \rightarrow G \quad h \mapsto ghg^{-1}.$$

Note that C_g is a diffeomorphism because it is a composition of diffeomorphisms:

$$C_g = L_g \circ R_{g^{-1}}.$$

In fact, $C_g^{-1} = C_{g^{-1}}$. Also, note that

$$\forall h_1, h_2 \in G \quad C_g(h_1 h_2) = g(h_1 h_2)g^{-1} = (gh_1 g^{-1})(gh_2 g^{-1}) = C_g(h_1)C_g(h_2).$$

Hence, $C_g : G \rightarrow G$ is a Lie group isomorphism (i.e a Lie group automorphism).

3 Week 7

3.1 Kernel and Image

Definition (Kernel and Image). Let G and H be two groups. Let $f : G \rightarrow H$ be a homomorphism.

- The **kernel** of f is the set

$$\ker f = \{g \in G : f(g) = 1_H\} = f^{-1}(\{1_H\}) \subseteq G.$$

- The **image** of f is

$$\operatorname{Im} f = \{f(g) : g \in G\} = f(G) \subseteq H.$$

Both of these sets are subgroups of G and H , respectively.

Example. Consider the determinant $\det : \operatorname{GL}(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ is a homomorphism. Thus,

$$\ker(\det) = \{A \in \operatorname{GL}(n, \mathbb{R}) : \det A = 1\} = \operatorname{SL}(n, \mathbb{R}).$$

Remark (Topology). If X and Y are two topological spaces, we may consider them to be the same if there exists a function $f : X \rightarrow Y$ such that

- (1) f is bijective
- (2) f is continuous
- (3) f^{-1} is continuous

In the context of manifold theory, if M and N are two manifolds, we may consider them to be the same if there exists $f : M \rightarrow N$ such that

- (1) f is bijective
- (2) f is smooth
- (3) f^{-1} is smooth

In the context of group theory, if G and H are two groups, we may consider them to be the same if

- (1) f is bijective
- (2) f is a homomorphism

The last remark gives us an explanation of why (ii) is important.

Remark (Fact from Linear Algebra). Suppose A and B are two $n \times n$ matrices and A is invertible. Then

$$\operatorname{rank}(AB) = \operatorname{rank}(B) \quad \operatorname{rank}(BA) = \operatorname{rank}(B).$$

Theorem (Kernel of a Lie Group Homomorphism). Let G and H be Lie groups. Let $f : G \rightarrow H$ be a Lie group homomorphism. Then

- (i) The rank of the derivative of f is constant for all points in G ; that is, there exists $k \in \mathbb{N}$ such that for all $x \in G$, $Df(x) = k$.

- (ii) $\ker(f)$ is an embedded submanifold of G of dimension $\dim(G) - k$ (and so it is a Lie subgroup of G)
- (iii) $\text{Im}(f)$ can be equipped with a smooth structure that makes it a Lie subgroup of H .

Proof. Note that (ii) is a direct consequence of (i) and Fact 2 from many weeks ago. Here we will prove (i). It suffices to show that for all $x \in G$, $\text{rank}(Df(x)) = \text{rank}(Df(e))$ where $e = 1_G$. For all $a, x \in G$, we have

$$\begin{aligned} f(x) &= f(aa^{-1}x) = f(a) \cdot f(a^{-1}x) \\ &= (L_{f(a)} \circ f \circ L_{a^{-1}})(x). \end{aligned}$$

By the chain rule,

$$Df(x) = DL_{f(a)}(f(a^{-1}x)) \cdot Df(a^{-1}x) \cdot DL_{a^{-1}}(x).$$

Since $L_{f(a)}$ and $L_{a^{-1}}$ are diffeomorphisms, $DL_{f(a)}$ and $DL_{a^{-1}}$ are invertible at any point. From a fact in linear algebra, for all $x, a \in G$

$$\text{rank}(Df(x)) = \text{rank}(Df(a^{-1}x)).$$

In particular, if we set $a = x$, we get

$$\text{rank}(Df(x)) = \text{rank}(Df(e))$$

which is our desired result. ■

Remark. Suppose G is a diffeomorphism. We have

$$(G^{-1} \circ G)(x) = x \implies DG^{-1}(G(x))DG(x) = I$$

Hence, $DG(x)$ is invertible.

Definition (Subgroups and cosets (very important!)). Let G be a group. Given a subgroup H of G and $g \in G$, the **left coset** of H determined by g is the set

$$gH = \{gh : h \in H\}.$$

The **right coset** of H determined by g is the set

$$Hg = \{hg : h \in H\}.$$

Note that H itself is a coset for $g = 1_G$.

You need a subgroup and an element to talk about a coset.

Example ($G = (\mathbb{R}^2, +)$, $H = \text{y-axis}$ Very important!). Observe that $H = \{(0, y) : y \in \mathbb{R}\}$ is a subgroup of $G = (\mathbb{R}^2, +)$. Let's determine the left cosets of H . For all $(a, b) \in \mathbb{R}^2$, we have

$$\begin{aligned} (a, b) + H &= \{(a, b) + (0, y) : y \in \mathbb{R}\} = \{(a, 0 + y) : y \in \mathbb{R}\} \\ &= \{(a, y) : y \in \mathbb{R}\}. \end{aligned}$$

This represents the vertical line at $x = a$. So, in this example, G decomposes into disjoint cosets. Roughly speaking, the plane is the union of parallel lines. This organizes the mess that is $\mathbb{R}^2$ into more manageable chunks to understand. Is this by accident? NO!

Note that two distinct $g \in G$ can give us the same coset (or vertical line in our example). That

is, consider the equivalent cosets

$$(1, 0) + H = (1, 1) + H = \text{vertical line at } x = a.$$

Theorem. Let G be a group and H be a subgroup of G .

- (i) $G = \bigcup_{g \in G} gH$
- (ii) Suppose $g_1, g_2 \in G$. Then either $g_1H = g_2H$ or $g_1H \cap g_2H = \emptyset$.
- (iii) $g_1H = g_2H$ if and only if $g_1^{-1}g_2 \in H$.

Proof. (i) For all $g \in G$, clearly $gH \subseteq G$ since gH is a subgroup of G . Therefore,

$$\bigcup_{g \in G} gH \subseteq G.$$

Suppose $g' \in G$. Since $1 \in H$, we have $g'1 \in g'H$. Thus, $g' \in g'H \subseteq \bigcup_{g \in G} gH$. So, $G \subseteq \bigcup_{g \in G} gH$. Hence,

$$G = \bigcup_{g \in G} gH.$$

- (ii) It suffices to show that if $g_1H \cap g_2H \neq \emptyset$, then $g_1H = g_2H$. Suppose $g \in g_1H \cap g_2H$. Thus,

$$g = g_1h_1 = g_2h_2 \text{ for some } g_1 \text{ and } g_2 \text{ in } H.$$

We have

$$\begin{aligned} g_1 = g_2h_2h_1^{-1} &\implies g_1H = g_2h_2h_1^{-1}H \\ &\implies g_1H = g_2(h_2h_1^{-1}H) \\ &\implies g_1H = g_2H. \end{aligned}$$

- (iii) ($\Leftarrow$) Suppose $g_1^{-1}g_2 \in H$. Our goal is to show that $g_1H = g_2H$. By (ii), it suffices to show that $g_1H \cap g_2H \neq \emptyset$. We have

$$\begin{aligned} g_1^{-1}g_2 \in H &\implies g_1(g_1^{-1}g_2) \in g_1H \\ &\implies g_2 \in g_1H. \end{aligned}$$

Similarly, we have $g_2 \in g_2H$. Thus, $g_2 \in g_1H \cap g_2H$. So, we have $g_1H \cap g_2H \neq \emptyset$.

($\Rightarrow$) Assume $g_1H = g_2H$. We claim that $g_1^{-1}g_2 \in H$.

$$\begin{aligned} g_1H = g_2H &\implies g_2 \in g_1H \\ &\implies g_2 = g_1h \text{ for some } h \in H \\ &\implies g_1^{-1}g_2 = h \text{ for some } h \in H \\ &\implies g_1^{-1}g_2 \in H. \end{aligned}$$

■

Theorem (Congruence modulo H). Let G be a group and H be a subgroup. Define the relation **congruence modulo H** on G as follows:

$$g_1 \sim g_2 \iff g_1^{-1}g_2 \in H.$$

Then the above relation is an equivalence relation, and its equivalence classes are precisely the left cosets of H .

Remark. • When we look at a group G , one very natural perspective is to view it as a disjoint union of cosets of a subgroup H .

- This viewpoint organizes the elements of G and often makes it easier to extract information about the structure of the group.
- In a way, the collection of cosets is simpler and more manageable than the group itself.

Remark. • It is tempting to try to make this collection of cosets into a group, using the natural operation

$$(g_1H)(g_2H) := (g_1g_2)H.$$

However, this does not always work; in general, the operation may be ill-defined.

- The question is as follows: *Under what conditions is the above operation well-defined?*

Lemma. Let G be a group and H be a subgroup of G . The following statements are equivalent:

- (1) $\forall g \in G, \forall h \in H, ghg^{-1} \in H$
- (2) $\forall g \in G, \forall h \in H, gHg^{-1} = H$
- (3) $\forall g \in G, gH = Hg$.

Definition (Normal Subgroups). In the case that any of the above equivalent conditions holds, we say H is a normal subgroup of G .

Notationally, normal subgroups can be represented by $H \trianglelefteq G$. If G is abelian, any subgroup of G is normal.

Theorem. Suppose G is a group and H is a subgroup of G . The operation

$$(g_1H) \cdot (g_2H) = (g_1g_2) \cdot H$$

is well-defined on the collection of left cosets of H if and only if H is a normal subgroup of G .

Proof. ($\implies$) Assume that the operation above is well-defined. We want to show that H is a normal subgroup of G . That is, it suffices to show that for all $g \in G$ and $h \in H$, we have

$$gHg^{-1} \in H.$$

That is, we want to show that (using the previous result $aH = bH \iff b^{-1}a \in H$)

$$g^{-1}H = (gh)^{-1}H.$$

That is, we want to show that

$$g^{-1}H = h^{-1}g^{-1}H \tag{*}$$

Since the coset operation is well defined, we have

$$\begin{aligned} g^{-1}H &= (1g^{-1})H = (1H)(g^{-1}H) \\ &\stackrel{1H=H=h^{-1}H}{=} (h^{-1}H)(g^{-1}H) \\ &= (h^{-1}g^{-1})H. \end{aligned}$$

($\Leftarrow$) Assume that $H \trianglelefteq G$. We want to show that the above operation is well-defined. Suppose $g_1H = g'_1H$ and $g_2H = g'_2H$. Our goal is to show that

$$(g_1g_2) = (g'_1g'_2)H. \quad (*)$$

Since H is normal, (that is, $g'_1H = Hg'_2$)

$$\begin{aligned} g'_1g'_2H &= g'_1Hg'_2 = g_1Hg'_2 & (g'_1H &= g_1H) \\ &= g_1g'_2H & (H \trianglelefteq G \implies Hg'_1 &= g'_2H) \\ &= g_1g_2H. & (g'_2H &= g_2H) \end{aligned}$$

and we are done. ■

Theorem (Quotient Theorem for Groups). If H is a normal subgroup of G , then the operation

$$(g_1H) \cdot (g_2H) = (g_1g_2)H$$

is well-defined on cosets

The group of cosets is called the **quotient group of G by H** , and is written G/H .

Remark. The map $\pi : G \rightarrow G/H$ where $g \mapsto gH$ is a surjective homomorphism and is called the **quotient map**.

By studying quotient groups of a group, which in a sense organize the group's elements into simpler chunks, we can gain valuable information about the group itself. For example, we can show that if $G/Z(G)$ is cyclic, then G is abelian (where $Z(G)$ is the center of G).

Every time we have a homomorphism and we look at the kernel of said homomorphism, they will always be normal subgroups.

Remark (Observation). Suppose G is a Lie group and N is a normal algebraic subgroup of G . Thus, G/N will be a new algebraic group. The question is:

Can we make G/N a Lie group?

It seems reasonable to look for a smooth structure on G/N that at least ensures that the quotient map $\pi : G \rightarrow G/N$ is smooth.

What is the kernel of π ? It is

$$\begin{aligned} \ker(\pi) &= \pi^{-1}\{1_{G/N}\} = \{g \in G : \pi(g) = 1_{G/N}\} \\ &= \{g \in G : \pi(g) = N\} \\ &= \{g \in G : gN = N\} \\ &= N. \end{aligned}$$

(very important!)

We can conclude from this result? If we manage to find a smooth structure on G/N that ensures $\pi : G \rightarrow G/N$ is smooth, then N is necessarily a topologically closed subset of G . If it is not closed, then there is no way to define a smooth structure on the homomorphism π .

Theorem (Quotient Theorem for Lie Groups). Let G be a Lie group, $N \trianglelefteq G$, and N is topologically closed subset of G . Then G/N can be equipped with a smooth structure that turns it into a Lie group such that the quotient map $\pi : G \rightarrow G/N$ is a surjective smooth map.

Example $(\mathbb{R}/\mathbb{Z})$. Note that $G = (\mathbb{R}, +)$ is a Lie group and $N = (\mathbb{Z}, +)$ is a subgroup of G . Since G is abelian, $N \trianglelefteq G$. Now, is N a closed subset? Yes, because the complement of $\mathbb{Z}$ (that is $\mathbb{R} \setminus \mathbb{Z}$) is open, so $\mathbb{Z}$ is a closed subset of $\mathbb{R}$. Using the Quotient theorem for Lie groups, we have that $\mathbb{R}/\mathbb{Z}$ is a Lie group.

As a matter of fact, the mapping

$$F : \mathbb{R}/\mathbb{Z} \rightarrow S^1 = U(1) \quad x + \mathbb{Z} \mapsto e^{2\pi i x}$$

is a well-defined isomorphism of Lie group. And so,

$$\mathbb{R}/\mathbb{Z} \cong S^1.$$

Why is this well-defined? Suppose $x + \mathbb{Z} = y + \mathbb{Z}$. We need to show that $e^{2\pi i x} = e^{2\pi i y}$. We have

$$\begin{aligned} x + \mathbb{Z} = y + \mathbb{Z} &\implies x - y \in \mathbb{Z} \\ &\implies e^{2\pi i(x-y)} = \cos(2\pi(x-y)) + i \sin(2\pi(x-y)) \\ &\implies e^{2\pi i(x-y)} = 1 + i0 = 1 \\ &\implies e^{2\pi i x} \cdot e^{-2\pi i y} = 1 \\ &\implies e^{2\pi i x} = e^{2\pi i y}. \end{aligned}$$

Similarly, we can show that

$$\mathbb{R}^n/\mathbb{Z}^n \cong (S^1) \times \cdots (S^1) \quad (n\text{-dimensional torus})$$

mapped by $(x_1, \dots, x_n) + \mathbb{Z}^n \mapsto (e^{2\pi i x_1}, e^{2\pi i x_2}, \dots, e^{2\pi i x_n})$.

Example. Let $G = \text{SU}(2)$ be a Lie group, $N = \{-I, +I\}$ is a normal subgroup of $\text{SU}(2)$, then

$$\forall g \in \text{SU}(2) \quad g(\pm I)g^{-1} = \pm g g^{-1} = \pm I \in N$$

where N is closed G (N is finite subset of G). Hence, $\text{SU}(2)/\{-I, I\}$ is a Lie group. In fact,

$$\text{SU}(2)/\{-I, +I\} \cong \text{SO}(3).$$