

Math 230B Lecture Notes

Lance Remigio

October 30, 2025

Contents

1	Week 5	1
2	Week 6	2
3	Week 7	7
3.1	Kernel and Image	7
3.2	A Common Theme in Mathematics	8
3.3	Kernel of a Lie Group Homomorphism	8
3.4	Subgroups and Cosets	9
3.5	Normal Subgroups	11
3.6	Making G/N a Lie Group	12
4	Week 8	13
4.1	Conjugate subgroups, Conjugacy Class	13
4.2	Review of Linear Algebra	14
4.3	Remark about First Isomorphism Theorem	15
4.4	Consequence of Yoneda Lemma	15
4.5	Second Isomorphism Theorem	15
4.6	Third Isomorphism Theorem	16
4.7	The Correspondence Theorem	16
4.8	Simple Groups	16
4.9	Direct Product of Groups	17
5	Week 9	18
6	Week 10	20
6.1	Tangent Spaces	20

1 Week 5

Example ($SL(n, \mathbb{F})$). Suppose $\mathbb{F} = \mathbb{R}$ (or $\mathbb{C}$). We claim that

$$SL(n, \mathbb{F}) = \{A \in GL(n, \mathbb{F}) : \det A = 1\}$$

is a Lie subgroup of $GL(n, \mathbb{F})$.

Indeed, we have that

- $SL(n, \mathbb{F})$ is closed under multiplication, indeed, if $A, B \in SL(n, \mathbb{F})$, then

$$\det(AB) = (\det A)(\det B) = 1 \cdot 1 = 1.$$

So, AB is invertible and $\det(AB) = 1$. This implies that

$$AB \in SL(n, \mathbb{F}).$$

- $SL(n, \mathbb{F})$ contains the inverse of each of its elements. Indeed, suppose $A \in SL(n, \mathbb{F})$, then A is invertible and

$$\det(A^{-1}) = \frac{1}{\det A} = \frac{1}{1} = 1.$$

So, $A^{-1} \in SL(n, \mathbb{F})$.

From the above, we can conclude that $SL(n, \mathbb{F})$ is a subgroup of $GL(n, \mathbb{F})$.

2 Week 6

Example (Special Orthogonal Group). Note that

$$\begin{aligned} SO(n) &= \{A \in M(n, \mathbb{R}) : A^T A = I \text{ and } \det A > 0\} \\ &= \{A \in M(n, \mathbb{R}) : A^T A = I \text{ and } \det A = 1\} \end{aligned}$$

We will prove that $SO(n)$ is a subgroup of $O(n)$.

- Note that $SO(n)$ is closed under multiplication. Indeed, suppose A and B are in $SO(n)$. Our goal is to show that $AB \in SO(n)$. Then we have

$$A, B \in SO(n) \implies A, B \in O(n) \implies AB \in O(n).$$

Since $A, B \in SO(n)$, it follows that $\det A = 1$ and $\det B = 1$. Hence,

$$\det AB = (\det A)(\det B) = 1 \cdot 1 = 1.$$

- Also observe that $SO(n)$ contains the inverse of each of its elements. Indeed, if $A \in SO(n)$, then

$$A \in SO(n) \implies A \in O(n) \implies A^{-1} \in O(n). \quad (1)$$

Furthermore,

$$A \in SO(n) \implies \det A = 1 \implies \det A^{-1} = \frac{1}{1} = 1 \implies A^{-1} \in SO(n) \quad (2)$$

From (1) and (2), we can see that $SO(n)$ is a subgroup of $O(n)$. Now, we need to find out whether $SO(n)$ is a Lie subgroup of $O(n)$. Note that $SO(n) = O(n) \cap \det^{-1}(0, \infty)$ where $(0, \infty)$ is an open set, $\det$ is a continuous function, and the preimage of an open set will be open in $M(n, \mathbb{R})$. Hence, $SO(n)$ will be open in $O(n)$. Thus, $SO(n)$ is an embedded submanifold of $O(n)$ with dimension $\frac{n^2-n}{2}$. Thus, $SO(n)$ is a Lie subgroup of $O(n)$ with dimension $\frac{n^2-n}{2}$.

Example (Special Unitary Group). Recall that

$$SU(n) = \{A \in M(n, \mathbb{C}) : A^* A = I \text{ and } \det A = 1\}.$$

We will show that the above is also a subgroup of $U(n)$. Clearly, we can see that $SU(n) \subseteq U(n)$.

- $SU(n)$ is closed under multiplication. Indeed, suppose A and B are in $SU(n)$. Then observe that

$$A, B \in SU(n) \implies A, B \in U(n) \implies AB \in U(n) \quad (1)$$

and

$$\begin{aligned} A, B \in SU(n) &\implies \det A = 1 \text{ and } \det B = 1 \\ &\implies \det AB = 1 \\ &\implies AB \in SU(n). \end{aligned}$$

- Also, $SU(n)$ contains the inverse of each of its elements. Suppose $A \in SU(n)$. Then

$$A \in SU(n) \implies A \in U(n) \implies A^{-1} \in U(n) \quad (3)$$

and

$$A \in SU(n) \implies \det A = 1 \implies \det A^{-1} = \frac{1}{\det A} = \frac{1}{1} = 1 \quad (4)$$

Now, (3) and (4) imply that $A^{-1} \in SU(n)$.

Now, we need to find out if $SU(n)$ a Lie subgroup of $U(n)$? Of course! Let $F : U(n) \rightarrow \mathbb{R}$ be defined as follows:

$$F(A) = \varphi.$$

where we previously proved that if $A \in U(n)$, then $|\det A| = 1$. So, there exists an angle $\varphi \in (-\pi, \pi]$ such that $\det A = e^{i\varphi}$. It can be shown that

$$\forall A \in U(n) \quad \text{rank} DF(A) = 1.$$

Thus, $SU(n) = F^{-1}(0)$ is an $n^2 - 1$ dimensional embedded submanifold of $U(n)$. Therefore, $SU(n)$ is a Lie subgroup of $U(n)$ with dimension $n^2 - 1$.

Definition (Bilinear Form). A function $B : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be a **bilinear form** on $\mathbb{R}^n$ if it satisfies the following condition:

- (1) $\forall x, y, z \in \mathbb{R}^n \quad B(x + y, z) = B(x, z) + B(y, z)$
- (2) $\forall c \in \mathbb{R} \text{ and } \forall x, y \in \mathbb{R}^n \quad B(cx, y) = cB(x, y)$
- (3) $\forall x, y, z \in \mathbb{R}^n \quad B(x, y + z) = B(x, y) + B(x, z)$
- (4) $\forall c \in \mathbb{R} \quad \forall x, y \in \mathbb{R}^n \quad B(x, cy) = cB(x, y)$

Moreover,

- The bilinear form B is said to be **symmetric** if

$$\forall x, y \in \mathbb{R}^n \quad B(x, y) = B(y, x).$$

- The bilinear form B is said to be **skew-symmetric** if

$$\forall x, y \in \mathbb{R}^n \quad B(x, y) = -B(y, x).$$

- The bilinear form B is said to be **non-degenerate** if the following statement is true

$$B(x, y) = 0 \quad \forall y \in \mathbb{R}^n \implies x = 0.$$

- A symmetric bilinear form B is said to be **positive definite** if

$$B(x, x) > 0 \quad \forall x \neq 0.$$

Example (Standard Inner Product on $\mathbb{R}^n$). The standard inner product in $\mathbb{R}^n$

$$\langle x, y \rangle = \sum_{i=1}^n x_i y_i$$

is a symmetric positive definite bilinear form on $\mathbb{R}^n$.

For example, let's verify condition (1) for bilinearity:

$$\begin{aligned}
 \langle x + y, z \rangle &= \sum_{i=1}^n [(x_i + y_i)z_i] \\
 &= \sum_{i=1}^n [x_i z_i + y_i z_i] \\
 &= \sum_{i=1}^n x_i z_i + \sum_{i=1}^n y_i z_i \\
 &= \langle x, z \rangle + \langle y, z \rangle.
 \end{aligned}$$

I leave you to check the other conditions yourself.

In general, a symmetric positive definite bilinear form on a real vector space V , by definition, is called an **inner product** on V .

Example (Standard Symplectic Form on $\mathbb{R}^n$). Define $w : \mathbb{R}^{2n} \times \mathbb{R}^{2n} \rightarrow \mathbb{R}$ as follows:

$$\forall x, y \in \mathbb{R}^{2n} \quad w(x, y) = \sum_{j=1}^n x_j y_{n+j} - x_{n+j} y_j.$$

For example,

$$\begin{aligned}
 n = 1 &\implies w(x, y) = x_1 y_2 + x_2 y_1 \\
 n = 2 &\implies w(x, y) = (x_1 y_3 - x_3 y_1) + (x_2 y_4 - x_4 y_2) \\
 &\vdots
 \end{aligned}$$

If we let $\Omega = \begin{pmatrix} O_{n \times n} & I_{n \times n} \\ -I_{n \times n} & O_{n \times n} \end{pmatrix}$, then we can write the standard symplectic form in the following way:

$$w(x, y) = \langle x, \Omega y \rangle$$

where $\langle, \rangle$ is the standard inner product in $\mathbb{R}^{2n}$.

Proposition (Preserving the standard symplectic form on $\mathbb{R}^{2n}$). Suppose $A \in M(2n, \mathbb{R})$. We say that A preserves the standard symplectic form on $\mathbb{R}^{2n}$ if and only if $A^T \Omega A = \Omega$.

Proof. We say that $A \in M(2n, \mathbb{R})$ preserves the standard symplectic form on $\mathbb{R}^{2n}$ if and only if

$$\begin{aligned}
 \forall x, y \in \mathbb{R}^{2n} \quad w(Ax, Ay) = w(x, y) &\iff \langle Ax, \Omega(Ay) \rangle = \langle x, \Omega y \rangle \\
 &\iff \langle x, A^T \Omega A y \rangle = \langle x, \Omega y \rangle \\
 &\iff \forall y \in \mathbb{R}^{2n} \quad A^T \Omega A y = \Omega y \\
 &\iff A^T \Omega A = \Omega.
 \end{aligned}$$

■

Proposition. If $A^T \Omega A = \Omega$, then A is invertible.

Proof. Suppose $A^T \Omega A = \Omega$, then we have

$$\begin{aligned}
 \det(A^T \Omega A) &= \det(\Omega) \\
 \implies \det(A^T) \det(\Omega) \det(A) &= \det(\Omega) \\
 \implies (\det(A))^2 \det(\Omega) &= \det(\Omega) \\
 \implies (\det(A))^2 &= 1 \\
 \implies \det(A) &= \pm 1 \\
 \implies A &\text{ is invertible.}
 \end{aligned}$$

■

Proposition. $A^T \Omega A = \Omega$ if and only if $-\Omega A^T \Omega = A^{-1}$ where A is invertible.

Proof. We have

$$\begin{aligned}
 -\Omega A^T \Omega = A^{-1} &\quad \Longleftrightarrow \quad -A^T \Omega = \Omega^{-1} A^{-1} \\
 &\quad \text{left multiply by } \Omega^{-1} \\
 &\quad \Longleftrightarrow \quad -A^T \Omega A = \Omega^{-1} \\
 &\quad \text{right multiply by } A \\
 &\quad \Longleftrightarrow \quad -A^T \Omega A = \Omega \\
 &\quad \Omega^{-1} = -\Omega \\
 &\quad \Longleftrightarrow \quad A^T \Omega A = \Omega.
 \end{aligned}$$

■

Example (Real Symplectic Group). $\text{Sp}(2n, \mathbb{R}) = \{A \in M(2n, \mathbb{R}) : A^T \Omega A = \Omega\}$ is a subgroup of $\text{GL}(2n, \mathbb{R})$.

Proof. We just proved that every matrix in $\text{SP}(2n, \mathbb{R})$ is invertible. So,

$$\text{SP}(2n, \mathbb{R}) \subseteq \text{GL}(2n, \mathbb{R}).$$

- We prove that $\text{SP}(2n, \mathbb{R})$ is closed under multiplication. Suppose $A, B \in \text{SP}(2n, \mathbb{R})$. We have

$$\begin{aligned}
 (AB)^T \Omega (AB) &= B^T (A^T \Omega A) B \\
 &= B^T \Omega B \\
 &= \Omega.
 \end{aligned}$$

Hence, we have $AB \in \text{SP}(2n, \mathbb{R})$.

- $\text{SP}(2n, \mathbb{R})$ contains the inverse of each of its elements. Suppose $A \in \text{SP}(2n, \mathbb{R})$. We have

$$\begin{aligned}
 A^T \Omega A = \Omega &\implies (A^T)^{-1} \Omega A = (A^T)^{-1} \Omega \\
 &\implies \Omega = (A^T)^{-1} \Omega A^{-1} \\
 &\implies \Omega = (A^{-1})^T \Omega A^{-1} \\
 &\implies A^{-1} \in \text{SP}(2n, \mathbb{R})
 \end{aligned}$$

From the above, we can see that $\text{SP}(2n, \mathbb{R})$ is a subgroup of $\text{GL}(2n, \mathbb{R})$. Now, we will prove that

$\mathrm{SP}(2n, \mathbb{R})$ is a Lie subgroup of $\mathrm{GL}(2n, \mathbb{R})$. Define the map

$$G : \mathrm{GL}(2n, \mathbb{R}) \subseteq \mathbb{R}^{4n^2} \rightarrow M(2n, \mathbb{R}) \cong \mathbb{R}^{4n^2} \text{ by } A \mapsto A^T \Omega A - \Omega.$$

Clearly, $\mathrm{SP}(2n, \mathbb{R}) = F^{-1}(\{0\})$. It can be shown that

$$\forall A \in \mathrm{GL}(2n, \mathbb{R}) \quad DF(A) \text{ has rank } \frac{4n^2 - 2n}{2} = 2n^2 - n.$$

Hence, $\mathrm{SP}(2n, \mathbb{R})$ is an embedded submanifold of $\mathrm{GL}(2n, \mathbb{R})$ with dimension $4n^2 - (2n^2 - n) = 2n^2 + n$. Thus, $\mathrm{SP}(2n, \mathbb{R})$ is a Lie subgroup of $\mathrm{GL}(2n, \mathbb{R})$. ■

Definition (Homomorphisms, Isomorphisms, Automorphisms). Let G and H be two groups.

- A function $\varphi : G \rightarrow H$ is said to be a **homomorphism** if it preserves group multiplication

$$\forall g_1, g_2 \in G \quad \varphi(g_1 g_2) = \varphi(g_1) \varphi(g_2).$$

- A bijective homomorphism is called an **isomorphism**. If there is an Isomorphism between G and H , we say G and H are isomorphic, and we write $G \cong H$.
- An **endomorphism** of G is a homomorphism from G to itself.
- An **automorphism** of G is an isomorphism from G to itself.

Definition (Lie Group Homomorphism). Let G and H be two Lie groups.

- A function $f : G \rightarrow H$ is said to be a **Lie group homomorphism** if
 - (i) f is a group homomorphism
 - (ii) f is smooth
- A function $f : G \rightarrow H$ is said to be a **Lie group isomorphism** if
 - (i) f is a group isomorphism
 - (ii) f is smooth
 - (iii) f^{-1} is smooth

If f is smooth and its inverse is smooth, then f is a diffeomorphism.

Example (The Determinant is a Lie Group Homomorphism). The determinant $\det : \mathrm{GL}(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ is a Lie Group Homomorphism:

- (i) $\forall A, B \in \mathrm{GL}(n, \mathbb{R})$, we have $\det(AB) = \det(A)\det(B)$. Hence, $\det$ is a group homomorphism.
- (ii) $\det A$ is a polynomial in terms of the entries of A and so differentiable. Hence, the $\det$ is smooth.

Similarly, the $\det : \mathrm{GL}(n, \mathbb{C}) \rightarrow \mathbb{C}^*$ is a Lie Group Homomorphism.

Example (Left Translation/right translation). Let G be a Lie Group and let $g \in G$. The **left translation** by g is the map

$$L_g : G \rightarrow G \quad h \mapsto gh.$$

The **right translation by g** is the map

$$R_g : G \rightarrow G \quad h \mapsto hg.$$

Notice that L_g is a composition of smooth maps and so it is smooth:

$$G \rightarrow G \times G \rightarrow G \quad h \mapsto (g, h) \mapsto gh.$$

Starting from left to right, we see that the first mapping is smooth because

- Fact 1: $F = (F_1, F_2)$ is smooth if and only if F_1 and F_2 are smooth.
- Fact 2: $h \mapsto g$ is a constant function implies that it is smooth.
- Fact 3: $h \mapsto h$ is the identity which is also smooth.

The second mapping $G \times G \rightarrow G$ is smooth because multiplication in G is smooth since G is a Lie group.

In fact, $L_g : G \rightarrow G$ is a diffeomorphism because the inverse L_g is $L_{g^{-1}}$:

$$\forall h \quad L_{g^{-1}} \circ L_g(h) = g^{-1}(gh) = h.$$

Unfortunately, L_g in general is NOT a group homomorphism; that is, in general

$$L_g(h_1 h_2) \neq L_g(h_1) L_g(h_2).$$

Similarly, the right translation map also satisfies the properties above but is NOT a group homomorphism.

Example (Conjugation by $g \in G$). Let G be a Lie group and let $g \in G$. The conjugation by g is the map:

$$C_g : G \rightarrow G \quad h \mapsto ghg^{-1}.$$

Note that C_g is a diffeomorphism because it is a composition of diffeomorphisms:

$$C_g = L_g \circ R_{g^{-1}}.$$

In fact, $C_g^{-1} = C_{g^{-1}}$. Also, note that

$$\forall h_1, h_2 \in G \quad C_g(h_1 h_2) = g(h_1 h_2)g^{-1} = (gh_1 g^{-1})(gh_2 g^{-1}) = C_g(h_1) C_g(h_2).$$

Hence, $C_g : G \rightarrow G$ is a Lie group isomorphism (i.e a Lie group automorphism).

3 Week 7

3.1 Kernel and Image

Definition (Kernel and Image). Let G and H be two groups. Let $f : G \rightarrow H$ be a homomorphism.

- The **kernel** of f is the set

$$\ker f = \{g \in G : f(g) = 1_H\} = f^{-1}(\{1_H\}) \subseteq G.$$

- The **image** of f is

$$\operatorname{Im} f = \{f(g) : g \in G\} = f(G) \subseteq H.$$

Both of these sets are subgroups of G and H , respectively.

Example. Consider the determinant $\det : \operatorname{GL}(n, \mathbb{R}) \rightarrow \mathbb{R}^*$ is a homomorphism. Thus,

$$\ker(\det) = \{A \in \operatorname{GL}(n, \mathbb{R}) : \det A = 1\} = \operatorname{SL}(n, \mathbb{R}).$$

3.2 A Common Theme in Mathematics

Remark (Topology). If X and Y are two topological spaces, we may consider them to be the same if there exists a function $f : X \rightarrow Y$ such that

- (1) f is bijective
- (2) f is continuous
- (3) f^{-1} is continuous

In the context of manifold theory, if M and N are two manifolds, we may consider them to be the same if there exists $f : M \rightarrow N$ such that

- (1) f is bijective
- (2) f is smooth
- (3) f^{-1} is smooth

In the context of group theory, if G and H are two groups, we may consider them to be the same if

- (1) f is bijective
- (2) f is a homomorphism

The last remark gives us an explanation of why (ii) is important.

3.3 Kernel of a Lie Group Homomorphism

Remark (Fact from Linear Algebra). Suppose A and B are two $n \times n$ matrices and A is invertible. Then

$$\text{rank}(AB) = \text{rank}(B) \quad \text{rank}(BA) = \text{rank}(B).$$

Theorem (Kernel of a Lie Group Homomorphism). Let G and H be Lie groups. Let $f : G \rightarrow H$ be a Lie group homomorphism. Then

- (i) The rank of the derivative of f is constant for all points in G ; that is, there exists $k \in \mathbb{N}$ such that for all $x \in G$, $\text{rank}(Df(x)) = k$.
- (ii) $\ker(f)$ is an embedded submanifold of G of dimension $\dim(G) - k$ (and so it is a Lie subgroup of G)
- (iii) $\text{Im}(f)$ can be equipped with a smooth structure that makes it a Lie subgroup of H .

Proof. Note that (ii) is a direct consequence of (i) and Fact 2 from many weeks ago. Here we will prove (i). It suffices to show that for all $x \in G$, $\text{rank}(Df(x)) = \text{rank}(Df(e))$ where $e = 1_G$. For all $a, x \in G$, we have

$$\begin{aligned} f(x) &= f(aa^{-1}x) = f(a) \cdot f(a^{-1}x) \\ &= (L_{f(a)} \circ f \circ L_{a^{-1}})(x). \end{aligned}$$

By the chain rule,

$$Df(x) = DL_{f(a)}(f(a^{-1}x)) \cdot Df(a^{-1}x) \cdot DL_{a^{-1}}(x).$$

Since $L_{f(a)}$ and $L_{a^{-1}}$ are diffeomorphisms, $DL_{f(a)}$ and $DL_{a^{-1}}$ are invertible at any point. From a

fact in linear algebra, for all $x, a \in G$

$$\text{rank}(Df(x)) = \text{rank}(Df(a^{-1}x)).$$

In particular, if we set $a = x$, we get

$$\text{rank}(Df(x)) = \text{rank}(Df(e))$$

which is our desired result. ■

Remark. Suppose G is a diffeomorphism. We have

$$(G^{-1} \circ G)(x) = x \implies DG^{-1}(G(x))DG(x) = I$$

Hence, $DG(x)$ is invertible.

3.4 Subgroups and Cosets

Definition (Subgroups and cosets (very important!)). Let G be a group. Given a subgroup H of G and $g \in G$, the **left coset** of H determined by g is the set

$$gH = \{gh : h \in H\}.$$

The **right coset** of H determined by g is the set

$$Hg = \{hg : h \in H\}.$$

Note that H itself is a coset for $g = 1_G$.

You need a subgroup and an element to talk about a coset.

Example ($G = (\mathbb{R}^2, +)$, $H = \text{y-axis}$ (Very important!)). Observe that $H = \{(0, y) : y \in \mathbb{R}\}$ is a subgroup of $G = (\mathbb{R}^2, +)$. Let's determine the left cosets of H . For all $(a, b) \in \mathbb{R}^2$, we have

$$\begin{aligned} (a, b) + H &= \{(a, b) + (0, y) : y \in \mathbb{R}\} = \{(a, 0 + y) : y \in \mathbb{R}\} \\ &= \{(a, y) : y \in \mathbb{R}\}. \end{aligned}$$

This represents the vertical line at $x = a$. So, in this example, G decomposes into disjoint cosets. Roughly speaking, the plane is the union of parallel lines. This organizes the mess that is $\mathbb{R}^2$ into more manageable chunks to understand. Is this by accident? NO!

Note that two distinct $g \in G$ can give us the same coset (or vertical line in our example). That is, consider the equivalent cosets

$$(1, 0) + H = (1, 1) + H = \text{vertical line at } x = 1.$$

Theorem. Let G be a group and H be a subgroup of G .

- (i) $G = \bigcup_{g \in G} gH$
- (ii) Suppose $g_1, g_2 \in G$. Then either $g_1H = g_2H$ or $g_1H \cap g_2H = \emptyset$.
- (iii) $g_1H = g_2H$ if and only if $g_1^{-1}g_2 \in H$.

Proof. (i) For all $g \in G$, clearly $gH \subseteq G$ since gH is a subgroup of G . Therefore,

$$\bigcup_{g \in G} gH \subseteq G.$$

Suppose $g' \in G$. Since $1 \in H$, we have $g'1 \in g'H$. Thus, $g' \in g'H \subseteq \bigcup_{g \in G} gH$. So, $G \subseteq \bigcup_{g \in G} gH$. Hence,

$$G = \bigcup_{g \in G} gH.$$

(ii) It suffices to show that if $g_1H \cap g_2H \neq \emptyset$, then $g_1H = g_2H$. Suppose $g \in g_1H \cap g_2H$. Thus,

$$g = g_1h_1 = g_2h_2 \text{ for some } g_1 \text{ and } g_2 \text{ in } H.$$

We have

$$\begin{aligned} g_1 = g_2h_2h_1^{-1} &\implies g_1H = g_2h_2h_1^{-1}H \\ &\implies g_1H = g_2(h_2h_1^{-1}H) \\ &\implies g_1H = g_2H. \end{aligned}$$

(iii) ($\Leftarrow$) Suppose $g_1^{-1}g_2 \in H$. Our goal is to show that $g_1H = g_2H$. By (ii), it suffices to show that $g_1H \cap g_2H \neq \emptyset$. We have

$$\begin{aligned} g_1^{-1}g_2 \in H &\implies g_1(g_1^{-1}g_2) \in g_1H \\ &\implies g_2 \in g_1H. \end{aligned}$$

Similarly, we have $g_2 \in g_2H$. Thus, $g_2 \in g_1H \cap g_2H$. So, we have $g_1H \cap g_2H \neq \emptyset$.

($\Rightarrow$) Assume $g_1H = g_2H$. We claim that $g_1^{-1}g_2 \in H$.

$$\begin{aligned} g_1H = g_2H &\implies g_2 \in g_1H \\ &\implies g_2 = g_1h \text{ for some } h \in H \\ &\implies g_1^{-1}g_2 = h \text{ for some } h \in H \\ &\implies g_1^{-1}g_2 \in H. \end{aligned}$$

■

Theorem (Congruence modulo H). Let G be a group and H be a subgroup. Define the relation **congruence modulo H** on G as follows:

$$g_1 \sim g_2 \iff g_1^{-1}g_2 \in H.$$

Then the above relation is an equivalence relation, and its equivalence classes are precisely the left cosets of H .

Remark. • When we look at a group G , one very natural perspective is to view it as a disjoint union of cosets of a subgroup H .

- This viewpoint organizes the elements of G and often makes it easier to extract information about the structure of the group.
- In a way, the collection of cosets is simpler and more manageable than the group itself.

Remark. • It is tempting to try to make this collection of cosets into a group, using the natural operation

$$(g_1H)(g_2H) := (g_1g_2)H.$$

However, this does not always work; in general, the operation may be ill-defined.

- The question is as follows: *Under what conditions is the above operation well-defined?*

Lemma. Let G be a group and H be a subgroup of G . The following statements are equivalent:

- (1) $\forall g \in G, \forall h \in H, ghg^{-1} \in H$
- (2) $\forall g \in G, \forall h \in H, gHg^{-1} = H$
- (3) $\forall g \in G, gH = Hg$.

3.5 Normal Subgroups

Definition (Normal Subgroups). In the case that any of the above equivalent conditions holds, we say H is a normal subgroup of G .

Notationally, normal subgroups can be represented by $H \trianglelefteq G$. If G is abelian, any subgroup of G is normal.

Theorem. Suppose G is a group and H is a subgroup of G . The operation

$$(g_1H) \cdot (g_2H) = (g_1g_2) \cdot H$$

is well-defined on the collection of left cosets of H if and only if H is a normal subgroup of G .

Proof. ($\implies$) Assume that the operation above is well-defined. We want to show that H is a normal subgroup of G . That is, it suffices to show that for all $g \in G$ and $h \in H$, we have

$$ghg^{-1} \in H.$$

That is, we want to show that (using the previous result $aH = bH \iff b^{-1}a \in H$)

$$g^{-1}H = (gh)^{-1}H.$$

That is, we want to show that

$$g^{-1}H = h^{-1}g^{-1}H \tag{*}$$

Since the coset operation is well defined, we have

$$\begin{aligned} g^{-1}H &= (1g^{-1})H = (1H)(g^{-1}H) \\ &\quad \underbrace{=}_{1H=H=h^{-1}H} (h^{-1}H)(g^{-1}H) \\ &= (h^{-1}g^{-1})H. \end{aligned}$$

($\impliedby$) Assume that $H \trianglelefteq G$. We want to show that the above operation is well-defined. Suppose $g_1H = g'_1H$ and $g_2H = g'_2H$. Our goal is to show that

$$(g_1g_2)H = (g'_1g'_2)H. \tag{*}$$

Since H is normal, (that is, $g'_2H = Hg'_2$ and $g'_1H = Hg'_1$)

$$\begin{aligned} g'_1g'_2H &= g'_1Hg'_2 = g_1Hg'_2 & (g'_1H &= g_1H) \\ &= g_1g'_2H & (H \trianglelefteq G \implies Hg'_2 &= g'_2H) \\ &= g_1g_2H. & (g'_2H &= g_2H) \end{aligned}$$

and we are done. ■

Theorem (Quotient Theorem for Groups). If H is a normal subgroup of G , then the operation

$$(g_1H) \cdot (g_2H) = (g_1g_2)H$$

is well-defined on cosets

The group of cosets is called the **quotient group of G by H** , and is written G/H .

Remark. The map $\pi : G \rightarrow G/H$ where $g \mapsto gH$ is a surjective homomorphism and is called the **quotient map**.

By studying quotient groups of a group which, in a sense, organizes the group's elements into simpler chunks, we can gain valuable information about the group itself. For example, we can show that if $G/Z(G)$ is cyclic, then G is abelian (where $Z(G)$ is the center of G).

The kernel of the quotient map will be a normal subgroup of G .

3.6 Making G/N a Lie Group

Remark (Observation). Suppose G is a Lie group and N is a normal algebraic subgroup of G . From this we can construct a new algebraic group G/N . In the context of Lie Theory, an important question arises:

Can we make G/N a Lie group?

It seems reasonable to look for a smooth structure on G/N that at least ensures that the quotient map $\pi : G \rightarrow G/N$ is smooth.

What is the kernel of π ? It is

$$\begin{aligned} \ker(\pi) &= \pi^{-1}\{1_GN\} = \{g \in G : \pi(g) = 1_GN\} \\ &= \{g \in G : \pi(g) = N\} \\ &= \{g \in G : gN = N\} \\ &= N. \end{aligned}$$

(very important!)

What can we conclude from this result? If we manage to find a smooth structure on G/N that ensures $\pi : G \rightarrow G/N$ is smooth, then N must be necessarily a topologically closed subset of G (recall that if π is smooth, then it is continuous and preimages of continuous functions of closed sets are closed). If it is not closed, then there is no way to define a smooth structure on the homomorphism π .

Theorem (Quotient Theorem for Lie Groups). Let G be a Lie group, $N \trianglelefteq G$, and N is topologically closed subset of G . Then G/N can be equipped with a smooth structure that turns it into a Lie group such that the quotient map $\pi : G \rightarrow G/N$ is a surjective smooth map.

Example ($\mathbb{R}/\mathbb{Z}$). Note that $G = (\mathbb{R}, +)$ is a Lie group and $N = (\mathbb{Z}, +)$ is a subgroup of G . Since G is abelian, $N \trianglelefteq G$. Now, is N a closed subset? Yes, because the complement of $\mathbb{Z}$ (that is $\mathbb{R} \setminus \mathbb{Z}$) is open, so $\mathbb{Z}$ is a closed subset of $\mathbb{R}$. Using the Quotient theorem for Lie groups, we have that $\mathbb{R}/\mathbb{Z}$ is a Lie group.

As a matter of fact, the mapping

$$\begin{aligned} F : \mathbb{R}/\mathbb{Z} &\longrightarrow S^1 = \mathrm{U}(1) \\ x + \mathbb{Z} &\longmapsto e^{2\pi i x} \end{aligned}$$

is a well-defined isomorphism of Lie group. And so,

$$\mathbb{R}/\mathbb{Z} \cong S^1.$$

Why is this well-defined? Suppose $x + \mathbb{Z} = y + \mathbb{Z}$. We need to show that $e^{2\pi i x} = e^{2\pi i y}$. We have

$$\begin{aligned} x + \mathbb{Z} = y + \mathbb{Z} &\implies x - y \in \mathbb{Z} \\ &\implies e^{2\pi i(x-y)} = \cos(2\pi(x-y)) + i\sin(2\pi(x-y)) \\ &\implies e^{2\pi i(x-y)} = 1 + i0 = 1 \\ &\implies e^{2\pi i x} \cdot e^{-2\pi i y} = 1 \\ &\implies e^{2\pi i x} = e^{2\pi i y}. \end{aligned}$$

Similarly, we can show that

$$\mathbb{R}^n/\mathbb{Z}^n \cong (S^1) \times \cdots \times (S^1) \quad (n\text{-dimensional torus})$$

mapped by $(x_1, \dots, x_n) + \mathbb{Z}^n \mapsto (e^{2\pi i x_1}, e^{2\pi i x_2}, \dots, e^{2\pi i x_n})$.

Example. Let $G = \mathrm{SU}(2)$ be a Lie group, $N = \{-I, +I\}$ is a normal subgroup of $\mathrm{SU}(2)$, then

$$\forall g \in \mathrm{SU}(2) \quad g(\pm I)g^{-1} = \pm gg^{-1} = \pm I \in N$$

where N is closed G (N is finite subset of G). Hence, $\mathrm{SU}(2)/\{-I, I\}$ is a Lie group. In fact,

$$\mathrm{SU}(2)/\{-I, +I\} \cong \mathrm{SO}(3).$$

4 Week 8

4.1 Conjugate subgroups, Conjugacy Class

Let G be a group. Recall that for each $g \in G$, the map conjugation by g , $C_g : G \rightarrow Gh \mapsto ghg^{-1}$ is a homomorphism. Since $C_g : G \rightarrow G$ is a homomorphism, for any subgroup, $H \leq G$, $C_g(H)$ will be a subgroup of G . Note that if H is a normal subgroup of G , then

$$C_g(H) = gHg^{-1} = H. \quad \forall g \in G$$

So,

$$\{C_g(H) : g \in G\} = H$$

which is just H . We can use this set as a measure of how our subgroup deviates from being normal. This motivates the following definition.

Definition (Conjugacy Class). For any subgroup $H \leq G$, we define the **conjugacy class** of H as follows:

$$\{C_g(H) : g \in G\} = \{gHg^{-1} : g \in G\}.$$

If $\tilde{H} \in \{C_g(H) : g \in G\}$, we say that subgroups H and $\tilde{H}$ are conjugate.

Example. Let $\theta \in \mathbb{R}$ be a fixed number. Consider the cyclic subgroup of $\text{SO}(3)$ generated by R_θ^z :

$$H = \langle R_\theta^z \rangle = \{R_{k\theta}^z : k \in \mathbb{Z}\}.$$

Note that for all $g \in \text{SO}(3)$, we have

$$\begin{aligned} gHg^{-1} &= \{gR_{k\theta}^zg^{-1} : k \in \mathbb{Z}\} \\ &= \text{the cyclic subgroup of } \text{SO}(3) \\ &\quad \text{generated by rotation } \theta \text{ radians about the axis} \\ &\quad \text{determined by the 3rd column of } g. \end{aligned}$$

Hence, we have

$$\begin{aligned} \text{Conjugacy class of } H &= \{gHg^{-1} : g \in \text{SO}(3)\} \\ &= \text{the collection of all cyclic subgroups} \\ &\quad \text{of } \text{SO}(3) \text{ generated by rotation } \theta \text{ radians about} \\ &\quad \text{an arbitrary fixed axis.} \end{aligned}$$

Theorem (First Isomorphism Theorem). Let $f : G \rightarrow H$ be a group homomorphism. Then $\varphi : G/\ker(f) \rightarrow \text{Im}(f)$ where $g\ker(f) \mapsto f(g)$.

Example (Observation of rotational matrices). • Rotation by an angle θ about the origin in the xy -plane can be described by the matrix

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}.$$

• Rotation by an angle θ about the z -axis in the xyz -space can be described by the matrix

$$R_\theta^z = \begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Example. Let ℓ be a directed line that passes through the origin in the xyz -space. In what follows, we will find a formula for rotation by an angle θ about ℓ (an arbitrary axis). Let $e_1 = (1, 0, 0)$, $e_2 = (0, 1, 0)$, $e_3 = (0, 0, 1)$ be the standard basis. Now, choose a new orthonormal basis $\{\hat{e}_1, \hat{e}_2, \hat{e}_3\}$ such that

- (1) It is right-handed ($\hat{e}_3 = \hat{e}_1 \times \hat{e}_2$)
- (2) $\hat{e}_3$ lies on ℓ (and points in the same direction as ℓ). Let $A = P_{\beta \leftarrow \hat{\beta}}$ be the change of basis matrix from $\hat{\beta}$ coordinate to β coordinates. Suppose w is a vector in $\mathbb{R}^3$. We can write $w = w_1e_1 + w_2e_2 + w_3e_3$ and $w = \hat{w}_1\hat{e}_1 + \hat{w}_2\hat{e}_2 + \hat{w}_3\hat{e}_3$. Recall that

$$[w]_\beta = \begin{pmatrix} w_1 \\ w_2 \\ w_3 \end{pmatrix}, \quad [w]_{\hat{\beta}} = \begin{pmatrix} \hat{w}_1 \\ \hat{w}_2 \\ \hat{w}_3 \end{pmatrix}.$$

4.2 Review of Linear Algebra

Suppose V and W are vector spaces with dimensions n and m , respectively. Suppose T is a linear transformation from V to W .

Remark (Huge Fact). If we choose a basis $\{b_1, \dots, b_n\}$ for V a basis $\{f_1, \dots, f_m\}$ for W , then T can be represented by an $m \times n$ matrix, denoted by $M(T)$ in the following sense: Given any vector $v = v_1b_1 + \dots + v_nb_n$ is equal to $w_1f_1 + w_2f_2 + \dots + w_mf_m$ where

$$\begin{pmatrix} w_1 \\ \vdots \\ w_m \end{pmatrix} = M(T) \begin{pmatrix} v_1 \\ \vdots \\ v_n \end{pmatrix}.$$

It can be shown that

$$M(T) = (T(b_1) \quad T(b_2) \quad \dots \quad T(b_n))$$

where for each $1 \leq i \leq n$, $T(b_i)$ will be expressed in terms of the basis in $\{b_1, \dots, b_n\}$.

Now, suppose V is a vector space and $\{b_1, \dots, b_n\}$ is one basis for V and $\{\hat{b}_1, \dots, \hat{b}_n\}$ is a second basis for V . We may consider the identity map

$$I : V \rightarrow V \quad I(x) = x$$

and equip the domain with $\{\hat{b}_1, \dots, \hat{b}_n\}$ basis, and equip the codomain with $b_1, \dots, b_n$. Then the $M(I)$ is exactly the change of coordinate matrix $P_{\beta \leftarrow \hat{\beta}}$ from $\hat{\beta}$ to β . Note that $A \in \text{SO}(3)$ because the columns of A are orthonormal and A preserves orientation. Let $v = v_1e_1 + v_2e_2 + v_3e_3$ be a vector in $\mathbb{R}^3$; that is, $[v]_\beta = (v_1, v_2, v_3)$. Let w be the vector that is obtained by rotating v by θ radians about the ℓ axis. We claim that

$$[w]_\beta = AR_\theta^z A^{-1}[v]_\beta.$$

Indeed, $A^{-1}[v]_\beta = [v]_{\hat{\beta}}$. Now, in the $\hat{\beta}$ basis, the ℓ axis is the same as the z -axis. Thus, the vector w that is obtained by rotation, θ radians about ℓ will have coordinates given by

$$[w]_{\hat{\beta}} = R_\theta^z[v]_{\hat{\beta}} = R_\theta^z A^{-1}[v]_\beta.$$

Therefore,

$$[w]_\beta = A[w]_{\hat{\beta}} = AR_\theta^z A^{-1}[v]_\beta.$$

That is,

$$[w]_\beta = (AR_\theta^z A^{-1})[v]_\beta.$$

4.3 Remark about First Isomorphism Theorem

Suppose G is a group and K is a normal subgroup of G . Then, as we know, the quotient map $\pi : G \rightarrow G/K$ where $g \mapsto gk$ is a surjective homomorphism. The above theorem for groups tells us that any surjective homomorphism with domain G is of the above form. That is, if $f : G \rightarrow H$ is a surjective homomorphism, we have

$$H \cong G/\text{Ker}(f).$$

Hence, your surjective homomorphism is, indeed, $G \rightarrow G/\text{Ker}(f)$. This theorem is giving us a way to construct all homomorphisms with domain G . In fact, we can make a connection to the following remarkable result from Category Theory.

4.4 Consequence of Yoneda Lemma

If you want to understand a mathematical object X , it suffices to understand all the structure preserving maps with domain X .

4.5 Second Isomorphism Theorem

Let G be a group. Let $K \trianglelefteq G$. As we know, the quotient map $\pi : G \rightarrow G/K$ is a surjective homomorphism. Suppose H is a subgroup of G . What is exactly $\pi(H)$ (we know that it must be a subgroup of G/K)? By the correspondence theorem, we know that

$$\pi(H) = (\text{a subgroup of } G \text{ that contains } K)/K.$$

The answer is:

$$\pi(H) = HK/K.$$

Theorem (Second Isomorphism Theorem). Let G be a group, H is a subgroup of G , K is a normal subgroup of G , then

$$\varphi : HK/K \rightarrow H/(H \cap K), \quad hk \mapsto h(H \cap K)$$

is an isomorphism

4.6 Third Isomorphism Theorem

Let G be a group, according to the first isomorphism theorem, in order to study homomorphisms with domain G , it suffices to study quotients G/N where N is a normal subgroup of G . Will it be helpful now to consider the quotients of G/N ? No! There will be no new information if you construct the quotients of G/N .

Theorem (Third Isomorphism Theorem). Let G be a group, K be a normal subgroup of G , N is a subgroup of K , and N is a normal subgroup of G . Then

$$\varphi : (G/N)/(K/N) \rightarrow G/K, \quad gN(K/N) \mapsto gK$$

is an isomorphism.

4.7 The Correspondence Theorem

Let G be a group and K be a normal subgroup of G . There are two questions we need to answer.

- (1) What are the subgroups of G/K ?
- (2) What are the normal subgroups of G/K ?

The answer to these questions are the following:

- (1) There is a one-to-one correspondence between the set of all subgroups of G that contain K and the set of all subgroups of G/K :
 - (i) If $H \leq G$ and $R \leq G/K$, then $H/K \leq G/K$.
 - (ii) If $R \leq G/K$, then $R = H/K$ for some $H \leq G$ and $K \subseteq H$.
 - (ii) The above correspondence preserves normality:

$$H/K \trianglelefteq G/K \iff H \trianglelefteq G, K \subseteq H.$$

4.8 Simple Groups

Definition (Simple Groups). Let G be an algebraic group. We say that G is **simple** if

- (1) $G \neq \{1_G\}$
- (2) The only normal subgroups of G are G itself and $\{1_G\}$.

Cannot make the group much simpler; that is, we cannot quotient the group out by a normal subgroup even further.

Remark (Observation). Many-to-one functions help us simplify/organize the domain of the function. An example is the quotient map $\pi : G \rightarrow G/N$ defined by $g \mapsto gN$. This organizes the domain of π into cosets.

Remark. The following two statements are equivalent:

- (1) G is not simple
- (2) There exists a many-to-one homomorphism of G onto a group G' other than the homomorphism sending all elements to 1.

That is, G' can be viewed as a simpler version of G (or, at any rate, not more complicated than G).

Remark. • A **simple Lie group** is not usually defined as a Lie group that is simple in the sense of abstract group theory.

- The most common definition is that a Lie group G is called **simple** if its corresponding Lie algebra is a simple Lie algebra.
- As we shall see, with this definition, a simple Lie group may not be simple as an abstract group.

4.9 Direct Product of Groups

Definition (Direct Product of Groups). If G_1 and G_2 are two groups, the **direct product** of G_1 and G_2 is the set $G_1 \times G_2$ with the group structure defined by the following multiplication rule:

$$(g_1, g_2) \cdot (g'_1, g'_2) = (g_1 g'_1, g_2 g'_2)$$

and the identity is

$$1_{G_1 \times G_2} = (1_{G_1}, 1_{G_2}).$$

If G_1 and G_2 are Lie groups, then $G_1 \times G_2$ is a Lie group.

- (1) $G_1 \times G_2$ is an algebraic group (previous slide)
- (2) G_1 is a smooth manifold and G_2 is a smooth manifold, from a month ago, we have $G_1 \times G_2$ is a smooth manifold.
- (3) multiplication map: $(g_1, g_2) \cdot (g'_1, g'_2) = (g_1 g'_1, g_2 g'_2)$. The multiplication in the first and second components are smooth operations. Hence, the multiplication in the direct product is also smooth.
- (4) Inversion: $(g_1, g_2) \mapsto (g_1^{-1}, g_2^{-1})$.

Example. We know that $S^1 = U(1) = \text{SO}(2)$ is a Lie group. It is, indeed, a compact and connected. Thus, $\underbrace{S^1 \times \cdots \times S^1}_{k \text{ times}}$ is also a Lie group. Note that this set is

- (1) connected
- (2) compact
- (3) abelian

Why is S^1 closed? We know that $S^1 = f^{-1}(1)$ where $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, $f(x, y) = x^2 + y^2$ and that because f is continuous, we know that the preimage of the closed set $\{1\}$ is closed. Clearly, we also see that S^1 is bounded. Since each set is compact, and there are k number of them, we see that because there are a finite product of compact sets, we know that the finite product must also be compact. Note that it is also true that a finite product of connected topological spaces is connected.

In fact, the only k -dimensional, where $k \geq 1$, Lie group that is compact, connected, and abelian is T^k . That is, if G is a k -dimensional compact connected abelian Lie group, then $G \cong T^k$ (where $\cong$ is a Lie group homomorphism). Here are some core ideas of the proof:

- (1) Use the fact G is abelian to show that the Lie Algebra of G is isomorphic to $\mathbb{R}^k$ with $[x, y] = 0$.
- (2) Use the facts that G is compact and connected to show that

$\exp : \mathfrak{g} \rightarrow G$ is surjective.

- (3) Use the fact that G is abelian to show that $\exp : \mathfrak{g} \rightarrow G$ is indeed a group homomorphism; that is, $\mathfrak{g}/\ker(\exp) \cong G$.
- (4) Use the fact that G is compact and that $\mathfrak{g}/\ker(\exp) \cong \mathbb{Z}^k$.
- (5) Conclude that

$$G \cong \mathfrak{g}/\ker(\exp) \cong \mathbb{R}^k/\mathbb{Z}^k \cong (S^1)^k = T^k.$$

End of Week 8

5 Week 9

In mathematics it is often helpful, when studying an object X (such as an algebraic structure, a metric space, or a normed space) to identify certain subsets/subobjects/substructures $E \subseteq X$ that are easier to work with or possess nice properties.

Example (Analysis). $X = C([0, 1], \mathbb{R})$ and $E = \mathbb{P}$. The following fact is true:

$\forall f \in X$ there exists $(P_n) \subseteq E$ such that $p_n \rightarrow f$ uniformly.

Example (Linear Algebra). Let $X = M(n, \mathbb{R})$ and $E = \text{Diagonal } n \times n \text{ real matrices}$. The connection here is that if $A \in X$ has n linearly independent eigenvectors $A = PDP^{-1}$ where P is invertible and $D \in E$. Another connection is that if $A \in X$ is symmetric, then $A = PDP^T$ where P is orthogonal and $D \in E$. (We say that every symmetric matrix is orthogonally diagonalizable)

Example (Complex Matrices). Let $X = M(n, \mathbb{C})$ and $E = \text{diagonal } n \times n \text{ matrices in } X$. If A is unitary (Hermitian) in X , then $PDP^* = A$ for some unitary matrix P and $D \in E$.

Remark (Linear Algebra). If $A \in M(n, \mathbb{C})$, then A is unitarily diagonalizable if and only if $A^*A = AA^*$ where such matrices are called normal matrices. This tells us that hermitian and unitary matrices are normal in this case.

Definition (Torus Subgroup). Let G be a Lie group. A Lie subgroup H of G is said to be a **Torus subgroup** of dimension $k \geq 1$ if

$$H \cong T^k$$

where $\cong$ is a Lie group isomorphism. That is, H is a k -dimensional Torus subgroup if H has dimension k and it is compact, connected, and abelian.

Definition (Maximal Torus). Let G be a Lie group and H be a Torus subgroup of G . We say that H is a **maximal torus** if H is not contained in any larger Torus subgroup of G .

Example ($\text{SO}(2) = \text{U}(1) = S^1$). Clearly, $\text{SO}(2)$ is its own maximal Torus.

Example ($\text{SO}(3)$). Note that

$$H = \left\{ R_\theta^z = \begin{pmatrix} \cos \theta & \sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} : \theta \in \mathbb{R} \right\}$$

is a copy of $\text{SO}(2) = S^1 = T^1$ inside $\text{SO}(3)$, so it is a Torus subgroup. We claim that H is maximal torus in $\text{SO}(3)$. The reason is as follows: Let T be a Torus subgroup of $\text{SO}(3)$ that contains H . It suffices to show that $T = H$ (and also H is maximal). Since any Torus subgroup is abelian, any $A \in T$ commutes with all $R_\theta^z \in H$:

$$\forall A \in T \text{ and } \forall \theta \in \mathbb{R} \quad AR_\theta^z = R_\theta^z A.$$

In what follows, we will show that if $A \in \text{SO}(3)$ satisfies (*), then $A \in H$. Let e_1, e_2, e_3 be our standard basis vectors. In order to show that $A \in \text{SO}(3)$ that satisfies (*) is indeed H , it suffices to show that:

- (1) $A|_{(e_1, e_2)\text{-plane}}$ is a Linear isometry of the (e_1, e_2) plane.
- (2) $\det A > 0$.

Proof. (1) It is enough to show that $Ae_1 \in (e_1, e_2)\text{-plane}$ and $Ae_2 \in (e_1, e_2)\text{-plane}$. Let $Ae_1 = \alpha_1 e_1 + \alpha_2 e_2 + \alpha_3 e_3$. Our goal is to show that $\alpha_3 = 0$. Note that, by assumption, A commutes with all R_θ^z , in particular, it commutes with

$$R_\pi^z = \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

Therefore,

$$AR_\pi^z(e_1) = R_\pi^z A(e_1) \iff AR_\pi^z(e_1) = A$$

■

Main point of this lecture is that not all Lie groups have a Torus subgroup.

End of Tuesday Lecture

Remark. We can travel from the Lie Algebra to the Lie group using the $\exp : \text{Lie}(G) \rightarrow G$ (where $\text{Lie}(G)$ is a Lie Algebra of the Lie group). This tells us that every compact Lie group has a Torus subgroup. Indeed, if x is any nonzero vector in $\text{Lie}(G)$, then the closure of $\{\exp(tx) : t \in \mathbb{R}\}$ is a compact connected abelian Lie subgroup of G . Note that H is a subgroup of $\text{Lie}(G)$ which is based on the two observations:

- (1) $e^{tx}e^{sx} = e^{(t+s)x} = e^{sx} \cdot e^{tx}$
- (2) $e^{tx}e^{-tx} = e^{0x} = 1_G$.

Note that

- (1) If H is a subgroup of Lie group G , then $\overline{H}$ is also a subgroup of G . Furthermore, if this is also abelian (which it is), then the closure is also abelian.
- (2) Closure of a connected set is connected (in any topological space).
- (3) A closed subset of a compact is compact.

We will prove (1) in this case.

Let G be a Lie group. Let H be an algebraic subgroup of G . Then the closure of H , denoted by $\overline{H}$, is a closed subgroup of G (and so it is a Lie subgroup of G). Moreover, if H is abelian, $\overline{H}$ is abelian also.

Proof. Note that clearly, $\overline{H}$ is closed. It suffices to show that $\overline{H}$ is closed under multiplication. Suppose a and b are in $\overline{H}$. Our goal is to show that $ab \in \overline{H}$. Note that $a \in \overline{H}$, there exists a sequence (a_n) in H such that $a_n \rightarrow a$ in G . Since $b \in \overline{H}$, there exists a sequence (b_n) in H such that $b_n \rightarrow b$ in G . Therefore,

$$(a_n, b_n) \rightarrow (a, b) \text{ in } G \times G.$$

Let's denote multiplication in G by $m : G \times G \rightarrow G$. We know $m : G \times G \rightarrow G$ is continuous. Then

$$m(a_n, b_n) = m(a, b) \implies a_n b_n \rightarrow ab \in G.$$

So, ab is the limit of a sequence of H . Thus, $ab \in \overline{H}$.

Also, $\overline{H}$ contains the inverse of each of its elements. Suppose $a \in \overline{H}$. Our goal is to show that $a^{-1} \in \overline{H}$. We have

$$a \in \overline{H} \implies \exists (a_n) \subseteq H \text{ such that } a_n \rightarrow a \in G.$$

Since inversion is a continuous map, we have

$$a_n^{-1} \rightarrow a^{-1} \in G.$$

Thus, a^{-1} is the limit of a sequence in H . Thus, $a^{-1} \in \overline{H}$.

Now, we want to prove that $\overline{H}$ is abelian. Suppose $a, b \in \overline{H}$. Our goal is to show that $ab = ba$. Similarly to what we saw above, there exists $(a_n) \subseteq H$ such that $a_n \rightarrow a$ in G and there exists a sequence $(b_n) \subseteq H$ such that $b_n \rightarrow b$ in G . So, we have $a_n b_n \rightarrow ab$ and $b_n a_n \rightarrow ba$. Then $ab = ba$. ■

Remark (Fact Used in the Above Proof From Analysis). If $z_n \rightarrow z$ and f is continuous, then $f(z_n) \rightarrow f(z)$.

6 Week 10

6.1 Tangent Spaces

Let M be a smooth manifold. What do we mean by a **smooth path** in M ?

Definition (Smooth Path). A **smooth path** in M is a smooth function $f : I \rightarrow M$ with $t \mapsto f(t)$ where $I \subseteq \mathbb{R}$ is a nonempty open interval. (We refer to this path as the path f or the path $f(t)$).

- Each component of the function is a smooth path in $\mathbb{R}$.
- Path connected implies continuous paths between two points a and b .
- We can also extend this notion to smooth paths between a and b .

Example (Examples of Smooth Paths). • $r : (-1, 1) \rightarrow \mathbb{R}^2$ defined by $r(t) = (t, t^2)$ is a smooth path in $M = \mathbb{R}^2$.

- $A : (0, \pi) \rightarrow \text{SO}(2)$, where

$$A(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}$$

is a **smooth path** in $M = \text{SO}(2)$. Consider the following two situations: Let $f : (0, \pi) \rightarrow \mathbb{R}^4$

where $t \mapsto (\cos t, -\sin t, \sin t, \cos t)$ and $A : (0, \pi) \rightarrow \text{SO}(2)$ where

$$t \mapsto \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}.$$

Note from these two examples, the implicit assumption is $\text{SO}(2) \subseteq \text{GL}(2, \mathbb{R}) \subseteq \mathbb{R}^4$.

- (An example of a non-smooth map) Let $A : (-\pi/2, \pi/2) \rightarrow \text{SO}(2)$, where

$$A(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}$$

is not a smooth path in $\text{SO}(2)$. (However, it is a continuous path in $\text{SO}(2)$).

Theorem (Manifold Theory (Relating to the 2nd Example Above)). Suppose N, M, L are three smooth manifolds. Suppose M is an embedded sub-manifold of L . Suppose $F : N \rightarrow M$. Then $F : N \rightarrow M$ is smooth if and only if $i \circ F : N \rightarrow L$ where i is the inclusion map ($i : M \rightarrow L$)

In our example (2), $N = (0, 1)$ and $M = \text{SO}(2)$ and $L = \mathbb{R}^4$. It is clear that $A : N \rightarrow L$ is smooth (because each smooth was smooth). So, by the above claim $A : N \rightarrow M$ is also smooth.

Note that derivative of a path can be thought of in the following way:

- If $r(t) = (r_1(t), r_2(t), \dots, r_n(t))$ is a smooth path in $\mathbb{R}^n$, then

$$r'(t) = \lim_{\delta t \rightarrow 0} \frac{r(t + \delta t) - r(t)}{\delta t} = (r'_1(t), r'_2(t), \dots, r'_n(t)).$$

Remark. Let M be an embedded submanifold of $\mathbb{R}^k$. Let $p \in M$. As we know the tangent space to M at p , $T_p M$, can be viewed as a vector space associated with $p \in M$. What is the precise definition of $T_p M$?

The answer is: There are multiple ways to define $T_p M$. Here is one of them

- $v \in T_p M$ if and only if there exists a path $A(t)$ in M such that $A(0) = p$ and $A'(0) = v$.
- More precisely, $v \in T_p M$ if and only if there exists $\varepsilon > 0$ and a smooth path $A : (-\varepsilon, \varepsilon) \rightarrow M$ such that $A(0) = p$ and $A'(0) = v$.
- Note that $A'(0)$ is sometimes referred to as the velocity vector of the path $A(t)$ at the point p .

Note that $T_p M$ consists of the velocity vectors of all smooth paths in M through $p \in M$.

Example. Prove that

$$(1) T_I(\text{O}(2)) = \{A \in M(2, \mathbb{R}) : A^T = -A\} = \left\{ \begin{pmatrix} 0 & -\theta \\ \theta & 0 \end{pmatrix} : \theta \in \mathbb{R} \right\}.$$

$$(2) T_I(\text{SO}(2)) = \{A \in M(2, \mathbb{R}) : A^T = -A\}.$$

Proof. Here we will prove (2). The proof of (1) is the same.

- (I) We claim that $T_I(\text{SO}(2)) \subseteq \text{skew}(2, \mathbb{R})$. Let $X \in T_I(\text{SO}(2))$. This tells us that there exists a smooth path $A : (-\varepsilon, \varepsilon) \rightarrow \text{SO}(2)$ such that $A(0) = I$ and $A'(0) = X$. Note that for all $t \in (-\varepsilon, \varepsilon)$ $A(t) \in \text{SO}(2)$. Then for all $t \in (-\varepsilon, \varepsilon)$ $A(t)^T A(t) = I$. Hence, we have

$$\frac{d}{dt}[A(t)^T A(t)] = 0 \implies A'(t)^T A(t) + A(t)^T A'(t) = 0.$$

In particular, if $t = 0$, we get

$$\begin{aligned} A'(0)^T A(0) + A(0)^T A'(0) &= 0 \\ \implies X^T I + I^T X &= 0 \\ \implies X^T + X &= 0 \\ \implies X^T &= -X. \end{aligned}$$

Hence, $X \in \text{Skew}(2, \mathbb{R})$.

(II) Now, we show that $\text{Skew}(2, \mathbb{R}) \subseteq T_I \text{SO}(2)$. Let $\begin{pmatrix} 0 & -\theta \\ \theta & 0 \end{pmatrix}$ be an arbitrary elements of $\text{Skew}(2, \mathbb{R})$.

We need to show that this matrix is in $T_I \text{SO}(2)$. That is, we need to show that there exists a smooth path $A(t)$ in $\text{SO}(2)$ such that $A(0) = I$ and $A'(0) = \begin{pmatrix} 0 & -\theta \\ \theta & 0 \end{pmatrix}$. Consider the path $A : (-1, 1) \rightarrow \text{SO}(2)$ given by

$$A(t) = \begin{pmatrix} \cos(\theta t) & -\sin(\theta t) \\ \sin(\theta t) & \cos(\theta t) \end{pmatrix}.$$

Notice that, clearly, $A(0) = I$. Also,

$$A'(t) = \begin{pmatrix} -\theta \sin(\theta t) & -\theta \cos(\theta t) \\ \theta \cos(\theta t) & -\theta \sin(\theta t) \end{pmatrix}$$

and so we have

$$A'(0) = \begin{pmatrix} 0 & -\theta \\ \theta & 0 \end{pmatrix}.$$

■

Remark. (i) As we saw in part 1 of the proof above, one can often find a condition that tangent vectors at the identity of a classical matrix Lie group G must satisfy by differentiating the defining equation of G .

(ii) Once this condition is identified, the next question is whether every vector satisfying it actually arises as a tangent vector at the identity. In other words, we need to check whether there exists smooth paths in G whose tangent at the identity matches the condition (this was part 2 of our solution).

(iii) It is at this stage, when constructing smooth paths in G , that a particularly useful tool, **the matrix exponential**, comes to our aid.

Definition (Matrix Exponential). Let $A \in M(n, \mathbb{C})$. We define the exponential of A , denoted by $\exp(A)$ or e^A , as follows:

$$\exp(A) = \sum_{m=0}^{\infty} \frac{A^m}{m!} = I + A + \frac{A^2}{2!} + \cdots$$

Example. • Let $D = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$. Compute e^D . Recall that if D is any diagonal matrix, then for

every natural number m , D^m is just raising the diagonal entries by m . Then

$$\begin{aligned} e^D &= \sum_{m=0}^{\infty} \frac{D^m}{m!} = \sum_{m=0}^{\infty} \frac{1}{m!} \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}^m \\ &= \sum_{m=0}^{\infty} \frac{1}{m!} \begin{pmatrix} 2^m & 0 \\ 0 & 3^m \end{pmatrix} \\ &= \end{aligned}$$