

# Homework 7

Lance Remigio

April 29, 2025

**Problem 1.** Let  $(V, \|\cdot\|)$  be an infinite dimensional normed space.

(i) Assume that  $(V, \|\cdot\|)$  is Banach.

**Problem 2 (Extra Credit).** Let  $(V, \|\cdot\|)$  be a normed space in which for any sequence  $(v_n)$  in  $V$

$$\sum_{n=1}^{\infty} \|v_n\| < \infty \implies \sum_{n=1}^{\infty} v_n \text{ converges in } V.$$

Prove that  $(V, \|\cdot\|)$  is Banach.

**Proof.** Suppose that every absolutely convergent series is convergent. Our goal is to show that  $(V, \|\cdot\|)$  is a Banach space. To do this, we will show that every Cauchy sequence in  $V$  converges. Let  $(v_n)$  be a Cauchy sequence in  $V$ . From here, our strategy is to find a subsequence  $(v_{n_k})$  of  $(v_n)$  such that  $(v_{n_k})$  converges in  $V$  (by the lemma). By definition,  $(v_n)$  being Cauchy implies that for all  $\varepsilon > 0$ , there exists an  $N \in \mathbb{N}$  such that for any  $n > m > N$ , we have

$$\|v_n - v_m\| < \varepsilon.$$

For  $\varepsilon = 1$ , there exists an  $n_1 \in \mathbb{N}$  such that for any  $n > m > n_1$ , we have

$$\|v_n - v_m\| < 1.$$

Furthermore, if  $\varepsilon = \frac{1}{2}$ . So, there exists an  $n_2 > n_1$  by the Archimedean Property such that for any  $n > m > n_2$ , we have

$$\|v_n - v_m\| < \frac{1}{2}.$$

In particular, if  $\varepsilon = \frac{1}{2^{k-1}}$  for all  $k \in \mathbb{N}$ , then we can find an  $n_k \in \mathbb{N}$  such that for any  $n > m > n_k$ , we have

$$\|v_n - v_m\| < \frac{1}{2^{k-1}}.$$

Moreover, by the Archimedean Property we can find an  $n_{k+1} \in \mathbb{N}$  such that  $n_{k+1} > n_k > n_{k-1}$ . Hence, it follows that  $(v_{n_k})$  is a subsequence in  $V$  such that

$$0 \leq \|v_{n_{k+1}} - v_{n_k}\| < \frac{1}{2^{k-1}}. \quad (*)$$

Note that since  $\sum_{k=1}^{\infty} \frac{1}{2^{k-1}}$  is a geometric series it follows from the Comparison Test that

$$\sum_{k=1}^{\infty} \|v_{n_{k+1}} - v_{n_k}\|$$

converges to some  $v \in V$ . By assumption, this tells us that

$$\sum_{k=1}^{\infty} (v_{n_{k+1}} - v_{n_k})$$

---

converges to some  $v$  in  $V$ . Now, observe that

$$\begin{aligned} v_{n_1} + \sum_{j=1}^{k-1} (v_{n_{j+1}} - v_{n_j}) &= v_{n_1} + (v_{n_2} - v_{n_1}) + (v_{n_3} - v_{n_2}) + \cdots + (v_{n_k} - v_{n_{k-1}}) \\ &= v_{n_k}. \end{aligned}$$

Taking the limit on both sides of the above equality, we see that

$$\begin{aligned} \lim_{k \rightarrow \infty} v_{n_k} &= \lim_{k \rightarrow \infty} \left[ v_{n_1} + \sum_{j=1}^{k-1} (v_{n_{j+1}} - v_{n_j}) \right] \\ &= v_{n_1} + \lim_{k \rightarrow \infty} \sum_{j=1}^{k-1} (v_{n_{j+1}} - v_{n_j}) \\ &= v_{n_1} + v. \end{aligned}$$

Thus, we now see that  $(v_{n_k})$  converges in  $V$  which tells us that  $(v_n)$  is a converges in  $V$ . Hence, ■

**Lemma.** Let  $(V, \|\cdot\|)$  be a normed space. Suppose  $(v_n)$  is a Cauchy sequence, and some subsequence  $(v_{n_k})$  converges to a point  $v$  in  $V$ . Then  $(v_n)$  converges to  $v$  in  $V$ .

**Proof.** Let  $n > m$ . Since  $(v_n)$  is a Cauchy sequence in  $V$ , it follows that

$$\|v_n - v_m\| \rightarrow 0$$

as  $n, m \rightarrow \infty$ . Also,  $(v_{n_k})$  converges to some  $v \in V$ . So, for  $k \rightarrow \infty$ , we have

$$\|v_{n_k} - v\| \rightarrow 0.$$

Using the triangle inequality, it follows that

$$0 \leq \|v_n - v\| \leq \|v_n - v_{n_k}\| + \|v_{n_k} - v\| \rightarrow 0.$$

Using the Squeeze Theorem, we have

$$\|v_n - v\| \rightarrow 0$$

as  $n \rightarrow \infty$  and we are done. ■