

Math 234A: Homework 1

Lance Remigio

September 3, 2024

1. (i) **(Parallelogram identity)** Let $z, w \in \mathbb{C}$. Show that

$$|z + w|^2 + |z - w|^2 = 2(|z|^2 + |w|^2).$$

Proof. Let $z, w \in \mathbb{C}$ with $z = x + iy$ and $w = u + iv$ with $x, y \in \mathbb{R}$ and $u, v \in \mathbb{R}$. Our goal is to show that

$$|z + w|^2 + |z - w|^2 = 2(|z|^2 + |w|^2).$$

Consider $|z - w|^2$ and notice that

$$z - w = (x - u) + i(y - v).$$

By definition of the modulus, we have

$$\begin{aligned} |z - w|^2 &= (z - w)\overline{z - w} \\ &= ((x - u) + i(y - v))((x - u) - i(y - v)) \\ &= (x - u)^2 + (y - v)^2 \\ &= x^2 - 2xu + u^2 + y^2 - 2yv + v^2 \\ &= (x^2 + y^2) - 2(xu + yv) + (u^2 + v^2) \\ &= |z|^2 - 2(xu + yv) + |w|^2. \end{aligned}$$

Note that

$$z + w = (x + u) + i(y + v).$$

$$\begin{aligned} |z + w|^2 &= (z + w)\overline{z + w} \\ &= ((x + u) + i(y + v))((x + u) - i(y + v)) \\ &= (x + u)^2 + (y + v)^2 \\ &= x^2 + 2xu + u^2 + y^2 + 2yv + v^2 \\ &= |z|^2 + 2(xu + yv) + |w|^2. \end{aligned}$$

Adding these two moduli together gives us

$$|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2 = 2(|z|^2 + |w|^2)$$

which is our desired result. ■

- (ii) **(Binomial Expansion):** Let $z, w \in \mathbb{C}$ and n be a positive integer. Show that

$$(z + w)^n = \sum_{k=0}^n \binom{n}{k} z^k w^{n-k}$$

where $\binom{n}{k} = \frac{n!}{k!(n-k)!}$.

Proof. Let $z, w \in \mathbb{C}$. We proceed via induction on $n \in \mathbb{Z}^+$ to show that

$$(z + w)^n = \sum_{k=0}^n \binom{n}{k} z^k w^{n-k}$$

where $\binom{n}{k} = \frac{n!}{k!(n-k)!}$. Let $n = 1$ be our base case. Then we have

$$\begin{aligned} \sum_{k=0}^1 \binom{1}{k} z^k w^{1-k} &= \binom{1}{0} z^0 w^1 + \binom{1}{1} z^1 w^0 \\ &= (z + w)^1, \end{aligned}$$

which tells us that the result holds in our base case. Now, suppose the result holds for n th case. We will show the result holds for the $n + 1$ case. By our induction hypothesis, we see that

$$\begin{aligned} (z + w)^{n+1} &= (z + w)(z + w)^n \\ &= (z + w) \sum_{k=0}^n \binom{n}{k} z^k w^{n-k} \\ &= \sum_{k=0}^n \binom{n}{k} z^{k+1} w^{n-k} + w \sum_{k=0}^n \binom{n}{k} z^k w^{n-k+1}. \end{aligned}$$

Reordering indices in the first summation by setting $m = k + 1$, we have

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k} z^{k+1} w^{n-k} + w \sum_{k=0}^n \binom{n}{k} z^k w^{n-k+1} &= \sum_{m=1}^{n+1} \binom{n}{m-1} z^m w^{(n+1)-m} \\ &\quad + \sum_{k=0}^n \binom{n}{k} z^k w^{(n+1)-k}. \end{aligned}$$

Then separating the first and last term of each summation, respectively, we have

$$\begin{aligned} (z + w)^{n+1} &= \binom{n}{0} z^n w + \sum_{m=1}^n \binom{n}{m-1} z^m w^{(n+1)-m} + \sum_{k=1}^n \binom{n}{k} z^k w^{n-k+1} + \binom{n}{n} w^{n+1} \\ &= \binom{n}{n} z^n w + \sum_{k=1}^n \left[\binom{n}{k} + \binom{n}{k-1} \right] z^k w^{(n+1)-k} + \binom{n}{0} w^{n+1}. \end{aligned}$$

Using the fact that **(Need to show this!)**

$$\binom{n}{k} + \binom{n}{k-1} = \binom{n+1}{k}$$

and collecting the first and last terms of the summation, we see that

$$\begin{aligned} (z + w)^{n+1} &= \binom{n}{0} z^0 w^{n+1} + \sum_{k=1}^n \binom{n+1}{k} z^k w^{(n+1)-k} + \binom{n}{n} z^{n+1} w^0 \\ &= \sum_{k=0}^{n+1} \binom{n+1}{k} z^k w^{(n+1)-k} \end{aligned}$$

which completes our induction argument. ■

2. For $z, w \in \mathbb{C}$. Define $\langle z, w \rangle = \Re(z\bar{w})$. (If we think of \mathbb{C} as two dimensional real vector space, then $\langle \cdot, \cdot \rangle$ defines an inner product on \mathbb{C}).

- (i) Cauchy Schwarz Inequality: Show that $|\langle z, w \rangle|^2 \leq |z|^2 |w|^2$ for all $z, w \in \mathbb{C}$.

Proof. First, we would like to show that for any $z, w \in \mathbb{C}$

$$\langle z, w \rangle^2 + \langle iz, w \rangle^2 = |z|^2 |w|^2.$$

Let $z, w \in \mathbb{C}$. By definition of $\langle z, w \rangle$, we see that

$$\langle z, w \rangle^2 + \langle iz, w \rangle^2 = (xu + yv)^2 + (xv - uy)^2. \quad (1)$$

Observe that

$$(xu + yv)^2 = (xu)^2 + 2xuyv + (yv)^2 \quad (2)$$

and

$$(xv - uy)^2 = (xv)^2 - 2xuyv + (uy)^2. \quad (3)$$

Then adding (2) and (3) results in

$$\begin{aligned} \langle z, w \rangle^2 + \langle iz, w \rangle^2 &= (xu)^2 + (yv)^2 + (xv)^2 + (uy)^2 \\ &= x^2 u^2 + y^2 v^2 + x^2 v^2 + u^2 y^2 \\ &= u^2(x^2 + y^2) + v^2(x^2 + y^2) \\ &= (x^2 + y^2)(u^2 + v^2) \\ &= |z|^2 |w|^2. \end{aligned}$$

Now, we need to show that $|\langle z, w \rangle|^2 \leq |z|^2 |w|^2$. By using the result above, we can write

$$|\langle z, w \rangle|^2 \leq |\langle z, w \rangle|^2 + |\langle iz, w \rangle|^2 = |z|^2 |w|^2$$

and we are done. ■

(ii) Triangle Inequalities: Show

$$|z + w| \leq |z| + |w|$$

and

$$||z| - |w|| \leq |z - w|$$

for all $z, w \in \mathbb{C}$.

Proof. Let $z, w \in \mathbb{C}$. We will first show that $|z + w| \leq |z| + |w|$. First, we will show the following results:

$$|z + w|^2 = |z|^2 + 2|\langle z, w \rangle| + |w|^2 \quad (1)$$

and

$$|z - w|^2 = |z|^2 - 2|\langle z, w \rangle| + |w|^2. \quad (2)$$

Let $z = x + iy$ and $w = u + iv$ for $x, y, u, v \in \mathbb{R}$. Observe that

$$z + w = (x + u) + i(y + v)$$

and

$$z - w = (x - u) + i(y - v).$$

Using the definition of the modulus, we see that

$$\begin{aligned} |z + w|^2 &= (z + w)\overline{(z + w)} = ((x + u) + i(y + v))((x + u) - i(y + v)) \\ &= (x + u)^2 + (y + v)^2 \\ &= x^2 + 2xu + u^2 + y^2 + 2yv + v^2 \\ &= (x^2 + y^2) + 2(xu + yv) + (u^2 + v^2) \\ &= |z|^2 + 2\langle z, w \rangle + |w|^2. \end{aligned} \quad \begin{array}{l} \text{(since } \Re(z\bar{w}) = xu + yv) \\ \text{(since } \langle z, w \rangle = \Re(z\bar{w})) \end{array}$$

Similarly, we have

$$\begin{aligned}
|z - w|^2 &= (z - w)\overline{(z - w)} = ((x - u) + i(y - v))((x - u) - i(y - v)) \\
&= (x - u)^2 + (y - v)^2 \\
&= x^2 - 2xu + u^2 + y^2 - 2yv + v^2 \\
&= x^2 + y^2 - 2(xu + yv) + u^2 + v^2 \\
&= (x^2 + y^2) - 2\langle z, w \rangle + (u^2 + v^2) \\
&= |z|^2 - 2\langle z, w \rangle + |w|^2.
\end{aligned}$$

Now, let us prove that $|z + w| \leq |z| + |w|$. Consider $|z + w|^2$. By part (a), we see that

$$\begin{aligned}
|z + w|^2 &= |z|^2 + 2\langle z, w \rangle + |w|^2 \\
&\leq |z|^2 + 2|z||w| + |w|^2 \\
&\leq (|z| + |w|)^2.
\end{aligned}$$

By taking the square root of both sides, we see that

$$|z + w| \leq |z| + |w|.$$

To show the second inequality, consider $|z - w|^2$. Then using part (a) again, we have

$$\begin{aligned}
|z - w|^2 &= |z|^2 - 2\langle z, w \rangle + |w|^2 \\
&\geq |z|^2 - 2|z||w| + |w|^2 \\
&= (|z| - |w|)^2.
\end{aligned}$$

By taking the square root of both sides, we see that

$$|z - w| \geq ||z| - |w||.$$

■

3. **(Lagrange Identity)** Let $z_1, \dots, z_n, w_1, \dots, w_n \in \mathbb{C}$. Show that

$$\left| \sum_{k=1}^n z_k w_k \right|^2 = \sum_{k=1}^n |z_k|^2 \sum_{k=1}^n |w_k|^2 - \sum_{1 \leq i < j \leq n} |z_i \overline{w_j} - z_j \overline{w_i}|^2.$$

Use this to deduce that

$$\left| \sum_{k=1}^n z_k w_k \right|^2 \leq \sum_{k=1}^n |z_k|^2 \sum_{k=1}^n |w_k|^2.$$

Proof. We will show that

$$\sum_{1 \leq i < j \leq n} |z_i \overline{w_j} - z_j \overline{w_i}|^2 = \sum_{k=1}^n |z_k|^2 \sum_{k=1}^n |w_k|^2 - \left| \sum_{k=1}^n z_k w_k \right|^2.$$

Observe that

$$\begin{aligned}
\sum_{1 \leq i < j \leq n} |z_i \overline{w_j} - z_j \overline{w_i}|^2 &= \sum_{i=1}^n \sum_{j=1}^n |z_i \overline{w_j} - z_j \overline{w_i}|^2 \\
&= \sum_{i=1}^n \sum_{j=1}^n (z_i \overline{w_j} - z_j \overline{w_i})(\overline{z_i} w_j - \overline{z_j} w_i) \\
&= \sum_{i=1}^n \sum_{j=1}^n \left[|z_i|^2 |w_j|^2 - 2 z_j \overline{z_i} \overline{w_i} w_j + |z_j|^2 |w_i|^2 \right] \\
&= \sum_{i=1}^n \sum_{j=1}^n |z_i|^2 |w_j|^2 - 2 \sum_{i=1}^n \sum_{j=1}^n z_j \overline{z_i} \overline{w_i} w_j + \sum_{i=1}^n \sum_{j=1}^n |z_j|^2 |w_i|^2 \\
&= \sum_{i=1}^n |z_i|^2 \sum_{j=1}^n |w_i|^2 + \sum_{i=1}^n |w_i|^2 \sum_{j=1}^n |z_j|^2 - 2 \sum_{i=1}^n \sum_{j=1}^n z_j \overline{z_i} \overline{w_i} w_j \\
&= \sum_{i=1}^n |z_i| \sum_{i=1}^n |w_i|^2 - \sum_{i=1}^n \overline{z_i} w_i \sum_{i=1}^n z_i w_i \quad (\text{for } i = j) \\
&= \sum_{i=1}^n |z_i|^2 \sum_{i=1}^n |w_i|^2 - \left| \sum_{i=1}^n z_i w_i \right|^2.
\end{aligned}$$

Thus, we conclude that

$$\begin{aligned}
\left| \sum_{k=1}^n z_k w_k \right|^2 &= \sum_{k=1}^n |z_k|^2 \sum_{k=1}^n |w_k|^2 - \sum_{1 \leq i < j \leq n} |z_i \overline{w_j} - z_j \overline{w_i}|^2 \\
&\leq \sum_{k=1}^n |z_k|^2 \sum_{k=1}^n |w_k|^2.
\end{aligned}$$

4. Express the following complex number in the form $\alpha + i\beta$:

(i) $(1 + i)^{-1}$

Solution. Observe that

$$(1 + i)^{-1} = \frac{1}{1 + i}$$

and that

$$\frac{1}{1 + i} \cdot \frac{(1 - i)}{(1 - i)} = \frac{(1 - i)}{1 - i^2} = \frac{1 - i}{2} = \frac{1}{2} - \frac{1}{2}i.$$

(ii) $(1 + i)/2i$

Solution. Observe that

$$\frac{(1 + i)}{2i} = \frac{1}{2i}(1 + i) = \frac{1}{2i} + \frac{1}{2} = \frac{1}{2} - \frac{1}{2}i.$$

(iii) $(5 + 5i)^{10}$

Solution. Let $z = 1 + i$. Observe that we can write

$$(5 + 5i)^{10} = 5^{10}(1 + i)^{10}.$$

Note that

$$r = \sqrt{1^2 + 1^2} = \sqrt{2}.$$

Furthermore, we have

$$\tan^{-1}(1/1) = \tan^{-1}(1) = \frac{\pi}{4}.$$

Using De Moivre's formula, we can write

$$\begin{aligned} z^{10} &= (\sqrt{2})^{10}(\cos(10\theta) + i\sin(10\theta)) \\ &= (\sqrt{2})^{10}(\cos(5\pi/2) + i\sin(5\pi/2)) = (\sqrt{2})^{10}i. \end{aligned}$$

Then we have

$$(5 + 5i)^{10} = 5^{10}(\sqrt{2})^{10}i = 312500000i.$$

■

(iv) $\left(\frac{2+i}{3-2i}\right)^2$

Solution. Our first step is to get $\frac{2+i}{3-2i}$ in terms of $\alpha + i\beta$. Thus, observe that

$$\frac{2+i}{3-2i} = \frac{2+i}{3-2i} \cdot \frac{3+2i}{3+2i} = \frac{7i+4}{13} = \frac{4}{13} + i\frac{7}{13}.$$

Furthermore, we have

$$\left(\frac{4}{13} + i\frac{7}{13}\right)^2 = \frac{1}{169}(4+7i)^2 = \frac{1}{169}(16+46i-49) = \frac{1}{169}(-33+46i).$$

Thus, we have that

$$\left(\frac{2+i}{3-2i}\right)^2 = \frac{-33}{169} + \frac{46}{169}i$$

■

(v) $\left(\frac{-1+i\sqrt{3}}{2}\right)^3.$

Solution. Denote $z = \frac{-1}{2} + i\frac{\sqrt{3}}{2}$. Then observe that

$$\theta = \tan^{-1}\left(\frac{\sqrt{3}/2}{-1/2}\right) = \tan^{-1}(-\sqrt{3}) = \frac{2\pi}{3}.$$

Furthermore, we have

$$r = \sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} = 1.$$

Using De Moivre's formula, we have that

$$z^3 = 1^3 \cdot \left(\cos\left(3 \cdot \frac{2\pi}{3}\right) + i\sin\left(3 \cdot \frac{2\pi}{3}\right)\right) = \cos(2\pi) + i\sin(2\pi) = 1 + i0 = 1.$$

■

5. Let $z \in \mathbb{C}^*$ and $z = \gamma(\cos \varphi + i\sin \varphi)$ where $n \in \mathbb{Z}^+$ and

$$w = \gamma^{1/n} \left[\cos\left(\frac{\varphi + 2\pi k}{n}\right) + i\sin\left(\frac{\varphi + 2\pi k}{n}\right) \right]$$

where $k \in \mathbb{Z}$. Show that $w^n = z$.

Proof. Note that for any $n \in \mathbb{N}$ that

$$z^n = (\cos \varphi + i\sin \varphi)^n = \cos n\varphi + i\sin n\varphi.$$

Thus, we have

$$\begin{aligned}
 w^n &= \left(\gamma^{1/n} \left[\cos \left(\frac{\varphi + 2\pi k}{n} \right) + i \sin \left(\frac{\varphi + 2\pi k}{n} \right) \right] \right)^n \\
 &= (\gamma^{1/n})^n \left[\cos \left(\frac{\varphi + 2\pi k}{n} \right) + i \sin \left(\frac{\varphi + 2\pi k}{n} \right) \right]^n \\
 &= \gamma \left[\cos \left(n \cdot \frac{\varphi + 2\pi k}{n} \right) + i \sin \left(n \cdot \frac{\varphi + 2\pi k}{n} \right) \right] \\
 &= \gamma (\cos \varphi + i \sin \varphi) \\
 &= z
 \end{aligned}$$

which ends our proof. ■

6. (Computing fourth roots): Find your distinct complex numbers w such that $w^4 = z$ for

(i) $z = i$.

Solution. ■

(ii) $z = -i$.

Solution. ■

(iii) $z = 1$.

Solution. ■

(iv) $z = -1$.

Solution. ■

7. Sketch the following sets in \mathbb{C} .

(i) $\zeta = \{z \in \mathbb{C} : \Re((1+i)z - 2) = 0\}$.

Solution. ■

(ii) Let $a, c \in \mathbb{R}$ and $b \in \mathbb{C}$ with $b\bar{b} - a > 0$ and

$$\zeta = \{z \in \mathbb{C} : a|z|^2 + \bar{b}z + b\bar{z} + c = 0\}.$$

Solution. ■

(iii) $\zeta = \{z \in \mathbb{C} : |z - i| = 2\}$.

Solution. ■

8. Let $z, a \in \mathbb{C}$.

(i) Show that $|1 - z\bar{a}|^2 - |z - a|^2 = (1 - |z|)^2(1 - |a|^2)$.

Proof. ■

(ii) Assume that $|a| < 1$. Show that

$$|z| < 1 \Leftrightarrow \left| \frac{z - a}{1 - \bar{a}z} \right| < 1$$

and

$$|z| = 1 \Leftrightarrow \left| \frac{z - a}{1 - \bar{a}z} \right| = 1.$$

Proof.

