

Recap: minimize  $f(x)$  / maximize  $g(x)$   
 (subject to  $x \in X$  ← constraint set)

TODAY:  $f: \mathbb{R} \rightarrow \mathbb{R}$

Examples:  
 minimize ... costs  
 distance  
 time  
 labor/effort/energy  
 waste  
 risk

Sidenote: We focus on the minimization formulation, as the maximizers of a given function  $g(x)$  are equal to the minimizers of  $-g(x) = f(x)$ .

Goal today: Gentle introduction via consideration of  $f: \mathbb{R} \rightarrow \mathbb{R}$ , function of one variable.

Definition: Let  $I \subseteq \mathbb{R}$  and consider  $f: I \rightarrow \mathbb{R}$  (functional).

A point  $x^* \in I$  is called:

1. a global minimizer of  $f$  on  $I$  if:  $\forall x \in I, f(x^*) \leq f(x)$
2. a strict global min. of  $f$  on  $I$  if:  $\forall x \in I, x \neq x^*, f(x^*) < f(x)$
3. a local minimizer of  $f$ :  
 $\exists \delta > 0, \forall x \in (x^* - \delta, x^* + \delta) \subset I, f(x^*) \leq f(x)$
4. a strict local min. of  $f$ :  
 $\exists \delta > 0, \forall x \in (x^* - \delta, x^* + \delta) \subset I, x \neq x^*, f(x^*) < f(x)$
5. A point  $x^* \in I$  is called a stationary point if  $f'(x^*) = 0$   
 (assuming that  $f'$  is defined at  $x^*$ )

# **Theorem (First-order necessary condition)**

Suppose  $f$  is continuously differentiable on  $I$ .

Proof: Recall:  $f'(x) = \lim_{x \rightarrow x^*} \frac{f(x) - f(x^*)}{x - x^*}$


Since  $x^*$  is a local minimizer of  $f$ :  $\exists \delta > 0, \forall x \in (x^* - \delta, x^* + \delta) \subset I, f(x^*) \leq f(x)$

case 1:  $x < x^* \Leftrightarrow x - x^* < 0 \Rightarrow f'(x^*) \leq 0$

case 2:  $x > x^* \Leftrightarrow x - x^* > 0 \Rightarrow f'(x^*) \geq 0$

By continuity of  $f'$ :  $f'(x^*) = f'(x^*) = 0$ .

If a point  $x^*$  in the interior of  $I$  is a local minimizer of  $f$ , then  $f'(x^*) = 0$ .



Next: Sufficient conditions for when stationary points are in fact minimizers  $\rightarrow$  Taylor expansion (HW1)

Theorem (Second-order sufficient conditions): Suppose  $f, f', f''$  are continuous on  $I$  ( $f \in C^2(I, \mathbb{R})$ ) and  $x^* \in I$  is a stationary point of  $f$ .

1. If  $f''(x) \geq 0 \quad \forall x \in I$ , then  $x^*$  is a global minimizer of  $f$  on  $I$ .
2. If  $f''(x) > 0 \quad \forall x \in I, x \neq x^*$ , then  $x^*$  is a strict global minimizer of  $f$  on  $I$ .
3.  $f''(x^*) > 0$ , then  $x^*$  is a strict local minimizer of  $f$ .

Proof of 2nd-order suff. conditions: If  $x \in I, x \neq x^*$ , Taylor's formula yields

$$f(x) = f(x^*) + f'(x^*)(x-x^*) + \frac{1}{2} f''(\xi)(x-x^*)^2 \quad \text{with } \xi \text{ between } x \text{ and } x^*.$$

1.  $f''(x) \geq 0 \quad \forall x \in I$  implies  $f''(\xi) \geq 0$  also. Moreover,  $x^*$  is a stationary point  $\Leftrightarrow f'(x^*) = 0$

$$\Rightarrow f(x) = f(x^*) + \frac{1}{2} \underbrace{f''(\xi)}_{\geq 0} \underbrace{(x-x^*)^2}_{\geq 0}$$

$$f(x) = f(x^*) + \text{"something"} \geq 0 \Rightarrow f(x^*) \leq f(x) \quad \forall x \in I.$$

2.  $x \neq x^*$  means  $(x-x^*)^2 > 0$ . Moreover  $f''(\xi) > 0$  (assuming  $\xi \neq x^*$ )

$$\Rightarrow f(x) = f(x^*) + \text{"something"} > 0 \Rightarrow f(x^*) < f(x) \quad \forall x \in I, x \neq x^*.$$

3. If  $f''(x^*) > 0$ , the continuity of  $f''$  implies that  $\exists \delta > 0$   $f''(x) > 0 \quad \forall x \in (x^* - \delta, x^* + \delta) \subset I$

$$\Rightarrow f(x) > f(x^*) \quad \forall x \in (x^* - \delta, x^* + \delta) \subset I.$$

Example:  $f(x) = 3x^4 - 4x^3 + 1$ .

- find stationary pt's:  $f'(x) = 12x^3 - 12x^2 = 12x^2(x-1) \Rightarrow x=0$  &  $x=1$  stationary points

- analyze stat. pt's:  $f''(x) = 36x^2 - 24x = 12x(3x-2)$

Case 1:  $x=1$ :  $f''(1) = 12 > 0 \Rightarrow$  strict local min.

Case 2:  $x=0$ :  $f''(0) = 0 \Rightarrow$  Then on 2nd order suff. condition doesn't provide information.