


Convergence analysis of the Gradient Method (GM) when applied to convex costs.

Theorem Let $f \in \mathcal{F}_L'(\mathbb{R}^n)$ with finite minimum $f^* = f(x^*)$ attained at $x^* \in \mathbb{R}^n$.

Then, for any $0 \leq \gamma \leq \frac{1}{L}$, the iterates of GM satisfy

$$(f(x^t) - f(x^*)) \leq \frac{1}{2\gamma t} \|x^0 - x^*\|^2.$$

This means that the $\{f(x^t)\}_{t \in \mathbb{N}_0}$ converges sublinearly to f^* .



"worst case analysis": $f(x^t) - f^* = \frac{1}{2\gamma t} \|x^0 - x^*\|^2 \leadsto f(x^{t+1}) - f^* = \frac{1}{2\gamma(t+1)} \|x^0 - x^*\|^2$

$$\frac{f(x^{t+1}) - f^*}{f(x^t) - f^*} = \dots = \frac{t}{t+1} \xrightarrow{t \rightarrow \infty} 1 \quad (\rho=1, \beta=1 \leadsto \text{sublinear convergence})$$

Proof (Do compare with proof of GM for general f in Lecture 5)

Two key bounds on $f \in \mathcal{F}_L'(\mathbb{R}^n)$:

- Lipschitz continuity of ∇f yields (Lemma in Lecture 6):

$$f(z) \leq f(x) + (\nabla f)(x)^T (z - x) + \frac{L}{2} \|z - x\|^2 \quad \forall z, x \in \mathbb{R}^n \quad (1)$$

- From convexity of f , we have:

$$f(z) \geq f(x) + (\nabla f)(x)^T (z - x) \quad \forall z, x \in \mathbb{R}^n. \quad (2)$$

Consider a single iteration of GM: $x^+ = x - \gamma(\nabla f)(x)$ with $0 \leq \gamma \leq \frac{1}{L}$. Then from (1) with $z = x^+$

$$f(x^+) \leq f(x) + (\nabla f)(x)^T (x^+ - x) + \frac{L}{2} \|x^+ - x\|^2$$

$$\leadsto f(x) + (\nabla f)(x)^T (-\gamma(\nabla f)(x)) + \frac{L}{2} \|\gamma(\nabla f)(x)\|^2$$

$$\stackrel{x^+ - x = -\gamma(\nabla f)(x)}{=} f(x) - \gamma \|\nabla f(x)\|^2 + \frac{L\gamma^2}{2} \|\nabla f(x)\|^2$$

$$\leq f(x) - \frac{\gamma}{2} \|\nabla f(x)\|^2 \quad \stackrel{\frac{L\gamma^2}{2} \leq \frac{\gamma}{2} \text{ with } (*)}{\leadsto}$$

$$L\gamma^2 \leq \gamma$$

$$\text{for } \gamma \leq \frac{1}{L}$$

$$(*) \text{ why? } L\gamma^2 - \gamma = \gamma(L\gamma - 1)$$



$$f(x^+) \leq f(x) - \frac{\gamma}{2} \|\nabla f(x)\|^2 \leadsto f(x^+) \leq f(x)$$

Now apply (2) with $z = x^*$: $f(x^*) \geq f(x) + (\nabla f)(x)^T(x^* - x) \Rightarrow f(x) \leq f(x^*) + (\nabla f)(x)^T(x - x^*)$
 (f is convex!)

$$\begin{aligned} f(x^+) &\leq f(x) - \frac{\gamma}{2} \|\nabla f(x)\|^2 \\ &\leq f(x^*) + (\nabla f)(x)^T(x - x^*) - \frac{\gamma}{2} \|\nabla f(x)\|^2 \\ &= f(x^*) + \frac{1}{2\gamma} \left(\|x - x^*\|^2 - \|x - x^*\|^2 + 2\gamma (\nabla f)(x)^T(x - x^*) - \gamma^2 \|\nabla f(x)\|^2 \right) \\ &= f(x^*) + \frac{1}{2\gamma} \left(\|x - x^*\|^2 - \underbrace{\|(x - x^*) - \gamma(\nabla f)(x)\|^2}_{\text{completion of squares}} \right) \quad \left| \begin{array}{l} - (a-b)^2 = -a^2 + 2ab - b^2 \end{array} \right. \\ &\quad \quad \quad = x^+ - x^* \text{ by def. of } x^+ = x - \gamma(\nabla f)(x) \\ &= \underline{f(x^*)} + \frac{1}{2\gamma} (\|x - x^*\|^2 - \|x^+ - x^*\|^2). \end{aligned}$$

Consider $x = x^{k-1}$, $x^+ = x^k$ and add the inequalities for $k=1, 2, \dots, t$:

$$\sum_{k=1}^t (f(x^k) - f(x^*)) \leq \frac{1}{2\gamma} \sum_{k=1}^t (\|x^{k-1} - x^*\|^2 - \|x^k - x^*\|^2) \underset{\text{telescoping sum}}{=} \frac{1}{2\gamma} (\|x^0 - x^*\|^2 - \|x^t - x^*\|^2) \leq \frac{1}{2\gamma} \|x^0 - x^*\|^2.$$

We saw $f(x^+) \leq f(x)$, so that $\underline{f(x^k)}$ is nonincreasing, which implies:

$$\underbrace{\frac{1}{t} \sum_{k=1}^t (f(x^k) - f(x^*))}_{\underline{f(x^t) - f(x^*)}} \leq \frac{1}{t} \sum_{k=1}^t (f(x^k) - f(x^*)) \leq \underline{\frac{1}{2\gamma t} \|x^0 - x^*\|^2}$$
