

Homework 3 Solutions

1. (a) Since α^k is a minimizer of
 $h(\alpha) = f(x^k + \alpha d^k)$,

it necessarily follows from the necessary condition
 that α^k is a critical point of h , i.e. $h'(\alpha^k) = 0$

$$h'(\alpha) = (\nabla f)(x^k + \alpha d^k)^T d^k$$

so

$$\begin{aligned} & \underbrace{(\nabla f)(x^k + \alpha^k d^k)^T d^k}_{= \langle d^k, (\nabla f)(x^k + \alpha^k d^k) \rangle} = 0 \end{aligned}$$

(b)

$d^k = -(\nabla f)(x^k)$
 $x^{k+1} = x^k + \alpha^k d^k$ with $d^k = -(\nabla f)(x^k)$
 $(\nabla f)(x^{k+1}) = (\nabla f)(x^k + \alpha^k d^k)$

Using the result from (a) with $\underline{d^k = -(\nabla f)(x^k)}$
 and $x^{k+1} = x^k + \alpha^k d^k$, the
 claim of $\langle (\nabla f)(x^k), (\nabla f)(x^{k+1}) \rangle = 0$ follows.
 can done inductively since via linear argument

Solution

Observe first that if we write $x = (x_1, x_2)^T$ then $f(x) = \frac{1}{2}x^T Q x - c^T x + a$ where

$$Q = \begin{bmatrix} 10 & 4 \\ 4 & 2 \end{bmatrix}, \quad c = \begin{bmatrix} 14 \\ 6 \end{bmatrix}, \quad \text{and } a = 20$$

Note that $Q \succ 0$, and recall that $\nabla f(x) = Qx - c$. The steepest descent step is $x^{k+1} = x^k + \alpha^k d^k$ where $d^k = -\nabla f(x^k)$, hence the algorithm is:

Step 0: Initial guess x^0 , threshold $\epsilon > 0$, counter $k = 0$.

Step 1: Choose direction $d^k = -\nabla f(x^k) = -Qx^k + c$. if $\|d^k\| < \epsilon$ stop.

Step 2: Compute step size

$$\alpha^k = \frac{(d^k)^T d^k}{(d^k)^T Q d^k}$$



Step 3: Set $x^{k+1} = x^k + \alpha^k d^k$, and increment counter $k = k + 1$. Go to step 1.

```
clc;clear all;close all
Q=[10 4; 4 2]; c=[14;6]; a=20; %define objective function
ep=10^(-6); %tolerance
X=[]; D=[]; A=[]; F=[]; %declare state, direction, step, function value

X(:,1)=[0;10]; %initialize state
D(:,1)=-Q*X(:,1)+c; %initial direction
A(1,1)=D(:,1)'*D(:,1)/(D(:,1)'*Q*D(:,1)); %initial step size
F(1,1)=X(:,1)'*Q*X(:,1)/2 - c'*X(:,1) + a;
for k=1:1000 %allow 1000 steps to converge
if(norm(D(:,k))<ep) break; end; %quit if tolerance reached
X(:,k+1)=X(:,k)+A(1,k)*D(:,k); %take the step
D(:,k+1)=-Q*X(:,k+1)+c; %descent direction
A(1,k+1)=D(:,k+1)'*D(:,k+1)/(D(:,k+1)'*Q*D(:,k+1)); %step size
F(1,k+1)=X(:,k+1)'*Q*X(:,k+1)/2 - c'*X(:,k+1) + a; %function value
end
%Contour plot
[xx,yy]=meshgrid(-10:.2:10);
zz=(Q(1,1)*xx.^2+(Q(2,1)+Q(1,2))*xx.*yy+Q(2,2)*yy.^2)/2-c(1)*xx-c(2)*yy+a;

figure
[C,h]=contour(xx,yy,zz,400); hold on, plot(X(1,:),X(2:),'k',X(1,:),X(2:),'r*')
hold off, xlabel('x'), ylabel('y')
% set(h,'ShowText','on','TextStep',get(h,'LevelStep')), colormap cool
set(h,'TextStep',get(h,'LevelStep')), colormap jet
axis([-5 5 -2 10])

figure
mesh(xx,yy,zz)
zlim([0 1200])
grid on
```

k	x_1^k	x_2^k	d_1^k	d_2^k	$\ d^k\ $	α^k	$f(x^k)$
1	0.000000	10.000000	-26.000000	-14.000000	29.529646	0.086645	60.000000
2	-2.252782	8.786963	1.379968	-2.562798	2.910712	2.180000	22.222576
3	0.755548	3.200064	-6.355739	-3.422321	7.218567	0.086645	12.987827
4	0.204852	2.903535	0.337335	-0.626480	0.711528	2.180000	10.730379
5	0.940243	1.537809	-1.553670	-0.836592	1.764590	0.086645	10.178542
6	0.805625	1.465322	0.082462	-0.153144	0.173934	2.180000	10.043645
7	0.985392	1.131468	-0.379797	-0.204506	0.431357	0.086645	10.010669
8	0.952485	1.113749	0.020158	-0.037436	0.042518	2.180000	10.002608
9	0.996429	1.032138	-0.092842	-0.049992	0.105446	0.086645	10.000638
10	0.988385	1.027806	0.004928	-0.009151	0.010394	2.180000	10.000156
11	0.999127	1.007856	-0.022695	-0.012221	0.025776	0.086645	10.000038
12	0.997161	1.006797	0.001205	-0.002237	0.002541	2.180000	10.000009
13	0.999787	1.001920	-0.005548	-0.002987	0.006301	0.086645	10.000002
14	0.999306	1.001662	0.000294	-0.000547	0.000621	2.180000	10.000001
15	0.999948	1.000469	-0.001356	-0.000730	0.001540	0.086645	10.000000
16	0.999830	1.000406	0.000072	-0.000134	0.000152	2.180000	10.000000
17	0.999987	1.000115	-0.000332	-0.000179	0.000377	0.086645	10.000000
18	0.999959	1.000099	0.000018	-0.000033	0.000037	2.180000	10.000000
19	0.999997	1.000028	-0.000081	-0.000044	0.000092	0.086645	10.000000
20	0.999990	1.000024	0.000004	-0.000008	0.000009	2.180000	10.000000
21	0.999999	1.000007	-0.000020	-0.000011	0.000023	0.086645	10.000000
22	0.999998	1.000006	0.000001	-0.000002	0.000002	2.180000	10.000000
23	1.000000	1.000002	-0.000005	-0.000003	0.000006	0.086645	10.000000
24	0.999999	1.000001	0.000000	0.000000	0.000001	2.180000	10.000000

- (b) Solve (a) with MATLAB optimization solver “fminunc” by setting the same threshold, i.e., the stopping criterion ε . Print out the first 10 and the last 10 iterations.

Solution: Here is the results obtained by starting from $x_0 = \begin{bmatrix} 0 \\ 10 \end{bmatrix}$.

First 10 iterations:

Iteration	Func-count	$f(x)$	Step-size	First-order optimality
0	3	60		26
1	6	33.9053	0.0384615	13.8
2	12	22.0956	0.1	3.57
3	18	20.7477	0.176204	3.55
4	24	19.55	0.158109	3.17
5	30	18.4858	0.176204	3.16
6	36	17.5401	0.158109	2.82
7	42	16.6999	0.176204	2.8
8	48	15.9533	0.158109	2.5
9	54	15.2899	0.176204	2.49
10	60	14.7004	0.158109	2.22

Last 10 iterations:

Iteration	Func-count	$f(x)$	Step-size	First-order optimality
192	1152	10	0.15485	4.65e-05
193	1158	10	0.181212	4.74e-05
194	1164	10	0.155063	4.14e-05
195	1170	10	0.181965	4.23e-05
196	1176	10	0.15361	3.65e-05
197	1182	10	0.183124	3.75e-05
198	1188	10	0.152404	3.23e-05
199	1194	10	0.18358	3.34e-05
200	1200	10	0.15122	2.86e-05
201	1206	10	0.183269	2.97e-05
202	1212	10	0.152357	2.56e-05

Code for reference:

```
clear
close all
opt = optimoptions(@fminunc,'Display','iter-detailed','Algorithm',...
    'quasi-newton','HessUpdate','steepdesc','MaxFunctionEvaluations',2000);
fun = @(x) 5*x(1)^2 + x(2)^2 + 4*x(1)*x(2) - 14*x(1) -6*x(2) + 20;
x0 = [0 10]';
[x,fval,exitflag,output] = fminunc(fun,x0,opt);
```


3

$$(a) \quad f(x_1, x_2) = \frac{x_1^4}{4} - x_1^2 + 2x_1 + \overbrace{(x_2^2 - 2x_2 + 1)}^{x_2^2 - 2x_2 + 1}$$

Newton's Method: $x^{k+1} = x^k - (Hf)(x^k)^{-1} (\nabla f)(x^k)$
So one step: $x^1 = x^0 - (Hf)(x^0)^{-1} (\nabla f)(x^0)$

$$(\nabla f)(x) = \begin{pmatrix} x_1^3 - 2x_1 + 2 \\ 2x_2 - 2 \end{pmatrix}$$

$$(Hf)(x) = \begin{pmatrix} 3x_1^2 - 2 & 0 \\ 0 & 2 \end{pmatrix}$$

$(Hf)(x)$ not invertible when $3x_1^2 - 2 = 0 \Leftrightarrow x_1 = \pm\sqrt{\frac{2}{3}}$

→ Newton's Method breaks down
when $x^0 = \begin{pmatrix} \pm\sqrt{\frac{2}{3}} \\ y \end{pmatrix}, y \in \mathbb{R}$.

3(b)

```
function newtons_method()
    % initialize
    x0 = [-1; 0];
    tolerance = 1e-6;
    max_iter = 10000;
    iter = 0;
    x = x0;

    path = x';
    fprintf('%8e %10e %10e %10e %10e %10e\n', 'Iter(k)', 'x1', 'x2', 'd1', 'd2', 'f(x)');

    while iter < max_iter
        H = hessian_f(x);
        g = gradient_f(x);

        if rank(H) < 2
            error('The hessian matrix is singular.');
```

end

```
        d = -H \ g;
        x = x + d;

        path = [path; x'];
        f_value = f(x);

        fprintf('%8d %10.4f %10.4f %10.4f %10.4f %10.4f\n', iter, x(1), x(2), d(1), d(2), f_value);

        if norm(d) < tolerance
            break;
        end

        iter = iter + 1;
    end

    [X1, X2] = meshgrid(linspace(-5, 2, 100), linspace(-2, 2, 100));
    Z = (X1.^4)/4 - X1.^2 + 2*X1 + (X2 - 1).^2;
    contour(X1, X2, Z, 50);
    hold on;
    plot(path(:,1), path(:,2), 'r-o', 'LineWidth', 1, 'MarkerSize', 5);
    title('Newton Method Path');
    xlabel('x_1');
    ylabel('x_2');
    hold off;
end

function f_val = f(x)
    f_val = (x(1)^4) / 4 - x(1)^2 + 2*x(1) + (x(2) - 1)^2;
end

function g = gradient_f(x)
    g = [x(1)^3 - 2*x(1) + 2; 2*(x(2) - 1)];
end

function H = hessian_f(x)
    H = [3*x(1)^2 - 2, 0; 0, 2];
end
```