

# Critical Metrology

## Parameter Estimation limits near Quantum Phase Transitions in the Quantum Rabi Model

Lander Zulueta Loizaga

Faculty of Science and Technology  
University of the Basque Country (UPV/EHU)

*Advisors: Iñigo L. Egusquiza & Jagoba Apellaniz*

July 2026

- ① Introduction to Critical Metrology
- ② Analytical Approach
- ③ Numerical Methods across the Phase Transition
- ④ Conclusions & Future Outlook

# What is Metrology?

Metrology is the science of measurement. We focus on **parameter estimation** in the *asymptotic limit* (large sample size case).

## Classical Parameter Estimation

Precision bounded by the **Shot-noise limit**:

$$(\Delta g)^2 \sim \frac{1}{\nu N}$$

where  $N$  is the size of the sample and  $\nu$  the number of repetitions.

## Quantum Parameter Estimation

Exploiting quantum resources allows reaching the **Heisenberg limit**:

$$(\Delta g)^2 \sim \frac{1}{\nu N^2}$$

A quadratic improvement over the classical case.


# Critical Metrology & Quantum Phase Transitions

## Critical Metrology:

- Exploits **extreme susceptibility** near a phase transition for enhanced parameter estimation.

## Quantum Phase Transition (QPT):

- Occurs at  $T = 0$ .
- Driven by **quantum fluctuations**.
- Signalled by the **closing of the energy gap**.
- Strictly requires the **thermodynamic limit** ( $N \rightarrow \infty$ ).



Energy gap closing at a QPT vs. avoided level-crossing.

# Precision Metric: Quantum Fisher Information

How do we quantify estimation precision?

## Quantum Cramér–Rao Bound

The variance of *any* unbiased estimator of  $g$  is bounded by:

$$(\Delta g)^2 \geq \frac{1}{\nu F_Q(\varrho_g)}$$

where  $\nu$  is the number of independent measurements and  $F_Q$  is the **Quantum Fisher Information** (QFI).

- **Our goal:** Show that  $F_Q$  diverges at the critical coupling  $g_c$  (w/o considering critical slowing down)
- For **pure states**:

$$F_Q = 4 \left[ \langle \partial_g \psi | \partial_g \psi \rangle - |\langle \partial_g \psi | \psi \rangle|^2 \right]$$

# The Quantum Rabi Model (QRM)

A single two-level atom coupled to one cavity mode:

$$H_{\text{QRM}} = \omega a^\dagger a + \Delta \sigma_z + g \sigma_x (a + a^\dagger)$$

- $\omega$ : cavity frequency,  $2\Delta$ : qubit splitting,  $g$ : coupling strength.

## Classical Oscillator Limit

The QRM achieves critical behavior with a *single* atom when

$$\frac{\Delta}{\omega} \longrightarrow \infty.$$

The frequency ratio plays the role of a *thermodynamic limit*. Exhibits a *Superradiant* QPT with order parameter  $\langle a^\dagger a \rangle \frac{\omega}{\Delta}$ .

# Analytical Approach: Schrieffer–Wolff Transformation

We decouple spin and bosonic subspaces via a unitary  $U = e^S$  ( $S$  anti-hermitian).

**First-order effective projected Hamiltonian:**

$$H_{\text{SW1}} = \omega a^\dagger a - \Delta - \frac{g^2}{2\Delta} (a + a^\dagger)^2$$

- Applied a mean-field to find the **critical point**:  $g_c = \sqrt{\omega \Delta/2}$
- Recognised GS as a *squeezed vacuum* via a Bogoliubov transformation.

Resulting QFI (Normal Phase):

$$F_Q(\varrho_g) \approx \frac{2g^2}{(\omega \Delta - 2g^2)^2} \xrightarrow{g \rightarrow g_c} \text{diverges}$$

- The **second-order** Schrieffer–Wolff expansion yields the same divergence.

# Overview & Limitations of the Analytic Approach

## Achievements:

$$F_Q(\varrho_g) \approx \frac{2g^2}{(\omega\Delta - 2g^2)^2} \propto (g_c - g)^{-2}$$

- Derived the QFI divergence *analytically*.
- Found a critical exponent  $\nu = -2$ .


## The Superradiant Phase Problem

Beyond  $g_c$  the effective potential inverts, a Bogoliubov is not suitable, *numerical* strategy is required. We will work on that after a quick **cross-check**.

# Numerical Validation (Normal Phase)


Before crossing the critical point we validate the numerics against the analytical prediction.

- Exact diagonalisation using QuTiP (truncation  $N=60$ ).
- **Result:** Numerical QFI (blue) matches the analytical divergence.
- **Note:** Instead of  $g$ , the dimensionless coupling strength  $\gamma = \frac{g}{\sqrt{\Delta\omega}}$  will be used onwards.  $\gamma_c = \frac{1}{\sqrt{2}}$ .



QFI vs.  $\gamma$  ( $\omega/\Delta = 5 \times 10^{-5}$ ,  $N=60$ ).

# Critical Exponent in the Normal Phase



Log-log scaling: critical exponent  $\nu = -2$  in the Normal Phase.

# The Superradiant Collapse


## Problem

Crossing  $\gamma_c$ , the standard Fock-basis computation breaks down.

## Why does it fail?

- 1 The photon number  $\langle a^\dagger a \rangle$  becomes *macroscopic*.
- 2 A Fock basis centred at the vacuum  $|0\rangle$  requires  $N \rightarrow \infty$  to capture the state.

**Consequence:** Fidelity with the true ground state drops to zero.



Fidelity collapse in the standard basis beyond  $\gamma_c$ .

## Solution: Displaced Squeezed Basis

To capture the superradiant ground state, we *adapt* the Fock basis in two steps.

- 1 **Displacement** — shift the origin to the mean-field amplitude  $\alpha_s$  (variational):

$$D(\alpha_s) = \exp(\alpha_s a^\dagger - \alpha_s^* a)$$


- 2 **Squeezing (Bogoliubov rotation)** — basis change to new bosonic operators  $(b, b^\dagger)$ .

### Result

The new  $(b, b^\dagger)$  and displaced basis efficiently hosts the macroscopic photon occupation of the superradiant phase.


# Convergence Restored

Using the displaced squeezed ( $b, b^\dagger$ ) basis, high fidelity is recovered with modest truncation ( $N \approx 60$ ).




# QFI Landscape across the Phase Diagram

Using the displaced squeezed basis, we can map the QFI in *both* phases.



# Entanglement Entropy



Entanglement entropy ( $\omega/\Delta = 5 \cdot 10^{-5}$ ): tendency change at  $\gamma_c$ . Validity of mean-field.

# Overview & Conclusions

- 1 **Finite-size criticality.** The QRM exhibits thermodynamic-like criticality in the  $\Delta/\omega \rightarrow \infty$  limit with a *single* atom.
- 2 **Divergent precision.** The QFI diverges at  $g_c$ , confirming the viability of critical metrology.
- 3 **Methodology.**
  - Analytical (Schrieffer–Wolff) methods yield analytical results in the Normal Phase.
  - Displacement + Squeezing enable stable numerics in the Superradiant Phase.

## Future directions:

- Study *critical slowing down* (dynamical protocols).
- Confirm intuition behind the anomalous branching of QFI in the superradiant phase.

# Thank you!




<https://github.com/Landerzl/Critical-Metrology>

## Questions & Discussion


# Experimental Implementations of the QRM

## Cavity QED (Optical Regime)




- Natural atoms (e.g., Alkali metals)
- Fabry-Pérot optical resonators
- Typically operates in the weak/strong coupling regimes

## Circuit QED (Microwave Regime)



- Artificial atoms (Superconducting circuits)
- Coplanar waveguide (CPW) resonators
- Access to ultrastrong/deep strong coupling

## Backup: Spectral Gap Closure



Ground state ( $E_0$ ) and first excited state ( $E_1$ ) energies versus coupling. The spectral gap closes exactly at the analytical prediction  $g_c = \sqrt{\omega\Delta/2}$ .

## Inspirational Quote

