

## Practice Problems for *Linear Algebra Review*

### Within Exam Scope

---

#### 1. Subspaces and Spans

Suppose  $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4, \mathbf{v}_5$  are vectors in  $\mathbb{R}^3$ . Do you agree with the statement that these vectors span  $\mathbb{R}^3$ ?

#### 2. Orthogonality

Which pairs are orthogonal among the vectors

$$\mathbf{v}_1 = \begin{pmatrix} 1 \\ 2 \\ -2 \\ 1 \end{pmatrix}, \mathbf{v}_2 = \begin{pmatrix} 4 \\ 0 \\ 4 \\ 0 \end{pmatrix}, \mathbf{v}_3 = \begin{pmatrix} 1 \\ -1 \\ -1 \\ -1 \end{pmatrix}$$

#### 3. Linear Independence

If  $\mathbf{v}_1, \mathbf{v}_2$  and  $\mathbf{v}_3$  are independent vectors, would the following vectors be independent?

$$\mathbf{w}_1 = \mathbf{v}_1 + \mathbf{v}_3, \quad \mathbf{w}_2 = \mathbf{v}_1 + \mathbf{v}_2, \quad \mathbf{w}_3 = \mathbf{v}_2 + \mathbf{v}_3.$$

#### 4. Range and Rank

For  $\mathbf{A} \in \mathbb{R}^{n \times n}$ , if  $\text{rank}(\mathbf{A}) = m < n$ , then what is the dimension of the null space?

#### 5. Matrix Inverse

We have matrices  $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$  and  $\mathbf{B} = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ . Which of the following are true?

- (a)  $\mathbf{A}$  is singular
- (b)  $\mathbf{B}$  is invertible
- (c)  $\mathbf{A} + \mathbf{B}$  is invertible.

#### 6. Subspaces

The smallest subspace of  $\mathbb{R}^3$  containing the vectors  $\begin{pmatrix} 0 \\ -3 \\ 6 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ 1 \\ -2 \end{pmatrix}$  is the line whose equations are  $x = a$  and  $z = by$ . What are the values of  $a$  and  $b$ ?

### 7. Matrix Inverse

For  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$  and  $\alpha \in \mathbb{R}$ , assume  $\mathbf{A}$  and  $\mathbf{B}$  are invertible:

- (a) (True/False)  $(\mathbf{A}^{-1})^\top = (\mathbf{A}^\top)^{-1}$ ,
- (b) (True/False)  $(\mathbf{AB})^{-1} = \mathbf{A}^{-1}\mathbf{B}^{-1}$ ,
- (c) (True/False) If  $\det(\mathbf{A}) = 2$ , then  $\det(\mathbf{A}^{-1}) = 2^{-1}$ .
- (d) (True/False) If  $\det(\mathbf{A}) = 2$  and  $\alpha > 1$ , then  $\det(\alpha\mathbf{A}) = 2$ .

### 8. Linear Independence

Let  $\mathbf{u} = \begin{pmatrix} \lambda \\ 1 \\ 0 \end{pmatrix}$ ,  $\mathbf{v} = \begin{pmatrix} 1 \\ \lambda \\ 1 \end{pmatrix}$ ,  $\mathbf{w} = \begin{pmatrix} 0 \\ 1 \\ \lambda \end{pmatrix}$ . What are possible values of  $\lambda$  that make  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  linearly dependent?

### 9. Matrix Product

For  $\mathbf{A} = \begin{bmatrix} 1 & 1/3 \\ x & y \end{bmatrix}$ , find the value of  $x$  and  $y$  such that  $\mathbf{A}^2 = \mathbf{0}$ .

### 10. Inner Product

Consider the space of all matrices in  $\mathbb{R}^{2 \times 2}$ . Define inner product as  $\langle \mathbf{A}, \mathbf{B} \rangle = \text{tr}(\mathbf{A}^\top \mathbf{B})$  for  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{2 \times 2}$ . Let  $\mathbf{U} = \begin{bmatrix} 1 & 4 \\ -3 & 5 \end{bmatrix}$  and  $\mathbf{V} = \begin{bmatrix} x^2 & x-1 \\ x+1 & -1 \end{bmatrix}$ . Find all values of  $x$  such that  $\mathbf{U} \perp \mathbf{V}$ .

### 11. Matrix Inverse

If a non-singular matrix  $\mathbf{A} \in \mathbb{R}^{n \times n}$  satisfies  $\mathbf{A}^3 - 4\mathbf{A}^2 + 3\mathbf{A} - 2\mathbf{I}_n = \mathbf{0}$ , what is  $\mathbf{A}^{-1}$ ?

### 12. Basis

Let  $\mathbf{u} = \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$ ,  $\mathbf{v} = \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}$  and  $\mathbf{w} = \begin{pmatrix} 1 \\ 2 \\ c \end{pmatrix}$  where  $c \in \mathbb{R}$ . The set  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a basis for  $\mathbb{R}^3$  provided that  $c$  is **not** equal to what value?

### 13. Subspaces

Consider the set of points  $(x, y, z) \in \mathbb{R}^3$ . Which one of the following is a subspace of  $\mathbb{R}^3$ ?

- (a)  $x + 3y - 2z = 3$ .
- (b)  $x + y + z = 0$  and  $x - y - z = 2$ .
- (c)  $\frac{x+1}{2} = \frac{y-2}{4} = \frac{z}{3}$ .
- (d)  $x^2 + y^2 = z$ .
- (e)  $x = -z$  and  $x = z$ .
- (f)  $\frac{x}{3} = \frac{y+1}{2}$ .

## Beyond Exam Scope

---

### 1. Solutions to Linear Equations

Consider the linear system of equations

$$\mathbf{Ax} = \mathbf{b} \tag{1}$$

where  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\mathbf{b} \in \mathbb{R}^m$  are known and  $\mathbf{x} \in \mathbb{R}^n$  is unknown. Let  $\mathbf{b} \in \mathcal{R}(\mathbf{A})$  so that (1) has at least one solution. Let  $\mathbf{x}_1$  be one such solution. Show that *all* solutions to (1) can be written in the form  $\mathbf{x}_1 + \mathbf{n}$  for some  $\mathbf{n} \in \mathcal{N}(\mathbf{A})$  where  $\mathcal{N}(\mathbf{A}) := \{\mathbf{x} \in \mathbb{R}^n : \mathbf{Ax} = \mathbf{0}\}$  is the *nullspace* of  $\mathbf{A}$ . (You can check that  $\mathcal{N}(\mathbf{A})$  is indeed a subspace of  $\mathbb{R}^n$ .)

---

### 2. Adjacency Matrices I (Challenging - Not Examinable!)

If  $\mathbf{A}$  is an adjacency matrix show that  $\mathbf{A}_{ij}^k = \#$  paths from  $i \rightarrow j$  in exactly  $k$  steps. *Hint:* Use induction. (If you're not familiar with induction that's ok and you can skip this question.)

---

### 3. Adjacency Matrices II

Consider the adjacency matrix  $\mathbf{A}$  below with rows and columns ordered according to the node labels  $\{'A', 'B', 'C', 'D', 'E', 'F', 'G'\}$ . Compute  $\mathbf{A}^3$  and confirm that the value in  $\mathbf{A}^3$  corresponding to paths from  $E \rightarrow B$  is correct by explicitly writing out and counting all the paths from  $E \rightarrow B$  that take exactly 3 steps.

$$\mathbf{A} = \begin{pmatrix} 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 1 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

---

### 4. Diagonalization

Suppose an  $n \times n$  matrix  $\mathbf{A}$  can be diagonalized and let  $\mathbf{u}_k := \mathbf{A}^k \mathbf{u}_0$  for some initial vector  $\mathbf{u}_0$ . Show that for any  $k \in \mathbb{N}$  we can write

$$\mathbf{u}_k = c_1 \lambda_1^k \mathbf{x}_1 + \cdots + c_n \lambda_n^k \mathbf{x}_n. \tag{2}$$

- (a) What are the  $c_i$ 's,  $\lambda_i$ 's and  $\mathbf{x}_i$ 's?
- (b) There are many problems of interest where  $\mathbf{u}_k$  represents the state of some system (e.g. an economy, or web-browser in Google's PageRank model) at time  $k$ . Very often, we are interested in the behavior of this system in the *long-run*, i.e. when  $k$  gets very large. How might the representation in (2) be useful for determining this long-run behaviour?

---

## 5. Cash-Flow Dynamics

Multinational companies in the U.S., Japan and Europe have assets of \$4 trillion. At the start \$2 trillion are in the U.S. and \$2 trillion in Europe. Each year  $1/2$  the U.S. money stays home, and  $1/4$  goes to each of Europe and Japan. For Japan and Europe,  $1/2$  stays home and  $1/2$  is sent to the U.S.

- (a) Find the  $3 \times 3$  matrix  $\mathbf{A}$  that gives  $\mathbf{x}_{k+1} = \mathbf{A}\mathbf{x}_k$  where  $\mathbf{x}_k$  is the  $3 \times 1$  vector containing the asset values in the US, Europe and Japan, respectively, at the end of year  $k$ .
- (b) Find the eigenvalues and eigenvectors of  $\mathbf{A}$ .
- (c) Find the cash-flow distribution at the end of year  $k$ .
- (d) Find the limiting cash-flow distribution of the \$4 trillion as the world ends.

(This question is taken from Gilbert Strang's *Linear Algebra and Its Applications*.)

---