

Homework 3

Hsin-Wei, Chen

B12902132 CSIE

Introduction to Analysis II

June 27, 2024

Problem 1. Find the radius and interval of convergence of the following power series.

1. $\sum_{n=3}^{\infty} \frac{x^n}{\log(n)}$
2. $\sum_{n=1}^{\infty} \frac{n!x^n}{n^n}$
3. $\sum_{n=0}^{\infty} n!x^n$

Let the convergence radius be R .

1. $\lim_{n \rightarrow \infty} \frac{\log(n+1)}{\log(n)} = 1 \Rightarrow R = 1$
2. $\lim_{n \rightarrow \infty} \frac{(n+1)!/(n+1)^{n+1}}{n!/n^n} = \lim_{n \rightarrow \infty} \frac{1}{(1+\frac{1}{n})^n} = 1/e \Rightarrow R = e$
3. $\lim_{n \rightarrow \infty} \frac{(n+1)!}{n!} = +\infty \Rightarrow R = 0$

Problem 2.

- (1) Show that if $\sum_{n=1}^{\infty} a_n = A$ ($C, 1$) and $\sum_{n=1}^{\infty} n^2 a_n < \infty$ exist, then the series converges to A as usual sense.
- (2) Show that if $\sum_{n=1}^{\infty} a_n = A$ ($C, 1$) and $\sum_{n=1}^{\infty} na_n^2 < \infty$ exist, then the series converges to A as usual sense.

Proof. (1) Let s_n be the n th partial sum of the series, and let $\sigma_n = \frac{1}{n} \sum_{k=1}^n s_k$, and since $\sum_{k=1}^n a_k$ is Cesaro

summable, suppose $\sigma_n \rightarrow A$ as $n \rightarrow \infty$. Since $\sum_{n=1}^{\infty} n^2 a_n < \infty \Rightarrow (n-1)a_n \rightarrow 0$ as $n \rightarrow \infty$. Given $\epsilon > 0$,

$na_n < \epsilon/2$, for some $N \in \mathbb{N}$. Choose $N' > \max\{\frac{2}{3}(N+1), \frac{2}{\epsilon} \sum_{k=1}^{N-1} (k-1)|a_k|\}$ Since

$$\begin{aligned} |s_n - \sigma_n| &= \left| \frac{\sum_{k=1}^n (s_n - s_k)}{n} \right| = \left| \frac{1}{n} \sum_{k=1}^n (k-1)a_k \right| \leq \sum_{k=1}^{N-1} \frac{k-1}{n} |a_k| + \sum_{k=N}^n \frac{k-1}{n} |a_k| \\ &\leq \sum_{k=1}^{N-1} \frac{k-1}{n} |a_k| + \frac{(N-n+1)}{n} \frac{\epsilon}{2} \\ &\leq \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

if $n > N'$. Hence $s_n = (s_n - \sigma_n) + \sigma_n \rightarrow 0 + A = A$.

(2) Let $\sigma_n \rightarrow A$ as $n \rightarrow \infty$, and $\sum_{k=1}^{\infty} ka_k^2 = M$.

$$\begin{aligned} s_n - \sigma_n &= \frac{\sum_{k=1}^n (s_n - s_k)}{n} = \frac{1}{n} \sum_{k=1}^n (k-1)a_k = \frac{1}{n} \sum_{k=1}^n ka_k - \frac{1}{n} s_n \\ \Rightarrow (1 + \frac{1}{n})s_n - \sigma_n &= \sum_{k=1}^n \frac{k}{n} a_k \\ \Rightarrow |(1 + \frac{1}{n})s_n - \sigma_n|^2 &= (\sum_{k=1}^n \frac{k}{n} a_k)^2 \leq (\sum_{k=1}^n \frac{1}{n^2}) (\sum_{k=1}^n k^2 a_k^2) = \frac{1}{n} \sum_{k=1}^n k^2 a_k^2 \quad \dots(1) \end{aligned}$$

Let $s'_n = \sum_{k=1}^n ka_k^2$ and $\sigma'_n = \frac{1}{n} \sum_{k=1}^n s'_k$

$$\begin{aligned} \sigma'_n &= \frac{1}{n} \sum_{k=1}^n (n+1-k)ka_k^2 = (1 + \frac{1}{n}) \sum_{k=1}^n ka_k^2 - \frac{1}{n} \sum_{k=1}^n k^2 a_k^2 = (1 + \frac{1}{n})s'_n - \frac{1}{n} \sum_{k=1}^n k^2 a_k^2 \\ \Rightarrow \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=1}^n k^2 a_k^2 &= \lim_{n \rightarrow \infty} (1 + \frac{1}{n})s'_n - \sigma'_n = M - M = 0 \end{aligned}$$

Hence $\lim_{n \rightarrow \infty} |(1 + \frac{1}{n})s_n - \sigma_n|^2 = 0 \Rightarrow \lim_{n \rightarrow \infty} (1 + \frac{1}{n})s_n - \sigma_n = 0 \Rightarrow \lim_{n \rightarrow \infty} s_n = A$

□

Problem 3. Let $a_n = \begin{cases} 1 & \text{if } n = 3k + 1 \\ 0 & \text{if } n = 3k + 2 \\ -1 & \text{if } n = 3k \end{cases}$. Find the $(C, 1)$ and the Abel sum of $\sum_{n=1}^{\infty} a_n$.

Proof. 1. Let s_n be the partial sum of the sequence $\{a_n\}$, then $s_n = \begin{cases} 1 & \text{if } n \neq 3k \\ 0 & \text{if } n = 3k \end{cases}$,

$$\sum_{n=0}^m s_n = \begin{cases} 2k+1 & \text{if } m = 3k+1 \\ 2k+2 & \text{if } m = 3k+2 \\ 2k & \text{if } m = 3k \end{cases}$$

. Since $\limsup_{m \rightarrow \infty} \frac{1}{m} \sum_{n=0}^m s_n = \lim_{k \rightarrow \infty} \frac{2k+2}{3k+2} = \frac{2}{3}$ and that $\liminf_{m \rightarrow \infty} \frac{1}{m} \sum_{n=0}^m s_n = \lim_{k \rightarrow \infty} \frac{2k}{3k} = \frac{2}{3}$,

then $\lim_{m \rightarrow \infty} \frac{1}{m} \sum_{n=0}^m s_n = \frac{2}{3}$. Hence $\sum_{n=1}^{\infty} a_n = \frac{2}{3} (C, 1)$.

2.

$$\sum_{n=1}^{\infty} a_n x^n = \sum_{k=0}^{\infty} x^{3k+1} - x^{3k+3} = \frac{x}{1-x^3} - \frac{x^3}{1-x^3} = \frac{x(1+x)}{1+x+x^2}, \quad x \in [0, 1)$$

then $\lim_{x \rightarrow 1^-} \frac{x(1+x)}{1+x+x^2} = \frac{2}{3}$. Hence Abel sum is $\frac{2}{3}$.

□

Problem 4. (Blancmange function)

Construct the function $g(x)$ by letting $g(x) = |x|$ for $x \in [-\frac{1}{2}, \frac{1}{2}]$ and extending g so that it becomes periodic on \mathbb{R} . Define

$$f(x) = \sum_{k=0}^{\infty} \frac{g(4^k x)}{4^k}$$

- (a) Use Weierstrass-M test to show that $f(x)$ is continuous on \mathbb{R} .
- (b) Show that f is differentiable at no point.

Proof.

- (a) Since $|\frac{g(4^k x)}{4^k}| \leq \frac{1}{2 \cdot 4^k}$, and $\sum_{k=0}^{\infty} \frac{1}{2 \cdot 4^k}$ converges, by Weierstrass M-test, $\sum_{k=1}^n \frac{g(4^k x)}{4^k}$ converges uniformly to $f(x)$ and since $\frac{g(4^k x)}{4^k}$ is continuous on \mathbb{R} , $f(x)$ is continuous on \mathbb{R} .
- (b) Fix a real number x and a positive number m .

Put

$$\delta_m = \pm \frac{1}{4^{m+1}}$$

The sign of δ_m is chosen so that there is no integer contained in $4^m(x + \delta_m)$ and $4^m x$.

Define

$$\gamma_k = \frac{g(4^k(x + \delta_m)) - g(4^k x)}{\delta_m}$$

Then for $k > m$, $4^k \delta_m$ is an integer, therefore $\gamma_k = 0$. When $0 \leq k \leq m$, $|\gamma_k| = 4^k$, then

$$\left| \frac{f(x + \delta_m) - f(x)}{\delta_m} \right| \pmod{2} \equiv \left| \sum_{k=0}^m \left(\frac{1}{4^k} \right) \gamma_k \right| \pmod{2} \equiv (m+1) \pmod{2}$$

Hence $\frac{f(x + \delta_m) - f(x)}{\delta_m}$ oscillates between odd numbers and even numbers as $m \rightarrow \infty$. Therefore $\lim_{m \rightarrow \infty} \frac{f(x + \delta_m) - f(x)}{\delta_m}$ does not exist, and since x is arbitrary, we conclude that f is differentiable at no point.

□