

Forward pass:

$$a_0^{(1)} = \sigma(w_{00}^{(1)} a_0^{(0)} + w_{01}^{(1)} a_1^{(0)} + w_{02}^{(1)}) \quad (1)$$

$$a_1^{(1)} = \sigma(w_{10}^{(1)} a_0^{(0)} + w_{11}^{(1)} a_1^{(0)} + w_{12}^{(1)}) \quad (2)$$

$$a_0^{(2)} = \sigma(w_{00}^{(2)} a_0^{(1)} + w_{01}^{(2)} a_1^{(1)} + w_{02}^{(2)}) \quad (3)$$

$$a_1^{(2)} = \sigma(w_{10}^{(2)} a_0^{(1)} + w_{11}^{(2)} a_1^{(1)} + w_{12}^{(2)}) \quad (4)$$

Or in matrix notation:

$$\vec{a}^{(1)} = \begin{pmatrix} a_0^{(1)} \\ a_1^{(1)} \end{pmatrix} = \sigma \left(\begin{pmatrix} w_{00}^{(1)} a_0^{(0)} + w_{01}^{(1)} a_1^{(0)} + w_{02}^{(1)} \\ w_{10}^{(1)} a_0^{(0)} + w_{11}^{(1)} a_1^{(0)} + w_{12}^{(1)} \end{pmatrix} \right) = \sigma \left(\begin{pmatrix} a_0^{(0)} & a_1^{(0)} & 1 \end{pmatrix} \begin{pmatrix} w_{00}^{(1)} & w_{10}^{(1)} \\ w_{01}^{(1)} & w_{11}^{(1)} \\ w_{02}^{(1)} & w_{12}^{(1)} \end{pmatrix} \right) = \sigma(\vec{A}^{(0)} W^{(1)})$$

$$\vec{a}^{(2)} = \sigma(\vec{A}^{(1)} W^{(2)})$$

Note that $\vec{a}^{(1)} = \begin{pmatrix} a_0^{(1)} \\ a_1^{(1)} \end{pmatrix}$ and $\vec{A}^{(1)} = (a_0^{(1)} \ a_1^{(1)} \ 1)$ both contain the values for the nodes in layer (1). $\vec{A}^{(1)}$ is needed to include the biases.

The costfunction is

$$C = \frac{1}{2}(a_0^{(2)} - T_0)^2 + \frac{1}{2}(a_1^{(2)} - T_1)^2$$

or using (3) and (4)

$$C = \frac{1}{2}(\sigma(w_{00}^{(2)} a_0^{(1)} + w_{01}^{(2)} a_1^{(1)} + w_{02}^{(2)}) - T_0)^2 + \frac{1}{2}(\sigma(w_{10}^{(2)} a_0^{(1)} + w_{11}^{(2)} a_1^{(1)} + w_{12}^{(2)}) - T_1)^2 \quad (5)$$

We can find how much each weight in layer (2) have to change by finding the derivative of C with respect to each weight.

Let's start with finding $\frac{dC}{dw_{00}^{(2)}}$

Only the first term of C in equation (5) include $w_{00}^{(2)}$. This term is a composite function $f \circ g \circ h(w_{00}^{(2)}, w_{01}^{(2)}, w_{02}^{(2)}, w_{10}^{(2)}, w_{11}^{(2)}, w_{12}^{(2)})$ where f, g and h and their derivatives are

$$f(\sigma) = \frac{1}{2}(\sigma - T_0)^2 \quad \frac{df}{d\sigma} = (\sigma - T_0)$$

$$g(h) = \sigma(h) \quad \frac{dg}{dh} = \frac{d\sigma}{dh} = \sigma(h)(1 - \sigma(h))$$

$$h(w_{00}^{(2)}, w_{01}^{(2)}, w_{02}^{(2)}) = w_{00}^{(2)} a_0^{(1)} + w_{01}^{(2)} a_1^{(1)} + w_{02}^{(2)} \quad \frac{dh}{dw_{00}^{(2)}} = a_0^{(1)}$$

By using (3) we note that

$$(\sigma - T_0) = (a_0^{(2)} - T_0)$$

and

$$\sigma(h)(1 - \sigma(h)) = a_0^{(2)}(1 - a_0^{(2)})$$

The chain rule

$$\frac{dC}{dw_{00}^{(2)}} = \frac{df}{d\sigma} \frac{d\sigma}{dh} \frac{dh}{dw_{00}^{(2)}}$$

then gives us $\frac{dC}{dw_{00}^{(2)}} = (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) a_0^{(1)}$, the first entry in the change matrix.

The rest is found in a similar manner:

$$\begin{pmatrix} \frac{dC}{dw_{00}^{(2)}} & \frac{dC}{dw_{01}^{(2)}} & \frac{dC}{dw_{02}^{(2)}} \\ \frac{dC}{dw_{10}^{(2)}} & \frac{dC}{dw_{11}^{(2)}} & \frac{dC}{dw_{12}^{(2)}} \end{pmatrix} = \begin{pmatrix} (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) a_0^{(1)} & (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) a_1^{(1)} & (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) 1 \\ (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) a_0^{(1)} & (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) a_1^{(1)} & (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) 1 \end{pmatrix}$$

We note that each entry in the upper row contains $\delta_0^{(2)} = (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)})$ and that each entry in the lower row contains $\delta_1^{(2)} = (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)})$.

Therefore

$$\frac{dC}{dW^{(2)}} = \begin{pmatrix} \frac{dC}{dw_{00}^{(2)}} & \frac{dC}{dw_{01}^{(2)}} & \frac{dC}{dw_{02}^{(2)}} \\ \frac{dC}{dw_{10}^{(2)}} & \frac{dC}{dw_{11}^{(2)}} & \frac{dC}{dw_{12}^{(2)}} \end{pmatrix} = \begin{pmatrix} \delta_0^{(2)} a_0^{(1)} & \delta_0^{(2)} a_1^{(1)} & \delta_0^{(2)} \cdot 1 \\ \delta_1^{(2)} a_0^{(1)} & \delta_1^{(2)} a_1^{(1)} & \delta_1^{(2)} \cdot 1 \end{pmatrix} = \delta^{(2)} A^{(1)}$$

where $\delta^{(2)} = \begin{pmatrix} \delta_0^{(2)} \\ \delta_1^{(2)} \end{pmatrix}$ and we had $A^{(1)} = (a_0^{(1)} a_1^{(1)} 1)$.

To find how much each weight in layer (1) have to change we can use (1) and (2) in (5)

$$C = \frac{1}{2} \left(\sigma \left(w_{00}^{(2)} \sigma(w_{00}^{(1)} a_0^{(0)} + w_{01}^{(1)} a_1^{(0)} + w_{02}^{(1)}) + w_{01}^{(2)} \sigma(w_{10}^{(1)} a_0^{(0)} + w_{11}^{(1)} a_1^{(0)} + w_{12}^{(1)}) + w_{02}^{(2)} \right) - T_0 \right)^2 + \frac{1}{2} \left(\sigma \left(w_{10}^{(2)} \sigma(w_{00}^{(1)} a_0^{(0)} + w_{01}^{(1)} a_1^{(0)} + w_{02}^{(1)}) + w_{11}^{(2)} \sigma(w_{10}^{(1)} a_0^{(0)} + w_{11}^{(1)} a_1^{(0)} + w_{12}^{(1)}) + w_{12}^{(2)} \right) - T_1 \right)^2 \quad (6)$$

We want to find $\frac{dC}{dw_{00}^{(1)}}$ and note that $dw_{00}^{(1)}$ occur in both terms of (6). The composit function has been a little more complicated, but nothing new really happens.

A little work will show that

$$\begin{aligned} \frac{dC}{dw_{00}^{(1)}} &= (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) w_{00}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_0^{(0)} + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{10}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_0^{(0)} \\ \frac{dC}{dw_{01}^{(1)}} &= (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) w_{00}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_1^{(0)} + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{10}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_1^{(0)} \\ \frac{dC}{dw_{02}^{(1)}} &= (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) w_{00}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) \cdot 1 + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{10}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) \cdot 1 \\ \frac{dC}{dw_{10}^{(1)}} &= (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) w_{01}^{(2)} a_1^{(1)} (1 - a_1^{(1)}) a_0^{(0)} + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{11}^{(2)} a_1^{(1)} (1 - a_1^{(1)}) a_0^{(0)} \\ \frac{dC}{dw_{11}^{(1)}} &= (a_0^{(2)} - T_0) a_1^{(2)} (1 - a_1^{(2)}) w_{01}^{(2)} a_0^{(1)} (1 - a_1^{(1)}) a_1^{(0)} + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{11}^{(2)} a_1^{(1)} (1 - a_1^{(1)}) a_1^{(0)} \\ \frac{dC}{dw_{12}^{(1)}} &= (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) w_{01}^{(2)} a_1^{(1)} (1 - a_1^{(1)}) \cdot 1 + (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)}) w_{11}^{(2)} a_1^{(1)} (1 - a_1^{(1)}) \cdot 1 \end{aligned}$$

If we again use $\delta_0^{(2)} = (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)})$ and $\delta_1^{(2)} = (a_1^{(2)} - T_1) a_1^{(2)} (1 - a_1^{(2)})$ we can write

$$\frac{dC}{dw_{00}^{(1)}} = \delta_0^{(2)} w_{00}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_0^{(0)} + \delta_1^{(2)} w_{10}^{(2)} a_0^{(1)} (1 - a_0^{(1)}) a_0^{(0)}$$

and by letting $\delta_0^{(1)} = a_0^{(1)}(1 - a_0^{(1)}) \sum_{i=0}^1 \delta_i^{(2)} w_{i0}^{(2)}$ and $\delta_1^{(1)} = a_1^{(1)}(1 - a_1^{(1)}) \sum_{i=0}^1 \delta_i^{(2)} w_{i1}^{(2)}$ we have

$$\frac{dC}{dw_{00}^{(1)}} = \delta_0^{(1)} a_0^{(0)}$$

so the change matrix for layer (1) is

$$\frac{dC}{dW^{(1)}} = \begin{pmatrix} \frac{dC}{dw_{00}^{(1)}} & \frac{dC}{dw_{01}^{(1)}} & \frac{dC}{dw_{02}^{(1)}} \\ \frac{dC}{dw_{10}^{(1)}} & \frac{dC}{dw_{11}^{(1)}} & \frac{dC}{dw_{12}^{(1)}} \end{pmatrix} = \begin{pmatrix} \delta_0^{(1)} a_0^{(0)} & \delta_0^{(1)} a_1^{(0)} & \delta_0^{(1)} \cdot 1 \\ \delta_1^{(1)} a_0^{(0)} & \delta_1^{(1)} a_1^{(0)} & \delta_1^{(2)} \cdot 1 \end{pmatrix} = \delta^{(1)} A^{(0)}$$

where $\delta^{(1)} = \begin{pmatrix} \delta_0^{(1)} \\ \delta_1^{(1)} \end{pmatrix}$ and $A^{(0)} = (a_0^{(0)} a_1^{(0)} 1)$.

By introducing the matrices

$$c = \begin{pmatrix} a_0^{(2)} - T^0 \\ a_1^{(2)} - T^1 \end{pmatrix}$$

$$w_2 = \begin{pmatrix} w_{00}^{(2)} & w_{01}^{(2)} \\ w_{10}^{(2)} & w_{11}^{(2)} \end{pmatrix}$$

$$D_2 = \begin{pmatrix} a_0^{(2)}(1 - a_0^{(2)}) & 0 \\ 0 & a_1^{(2)}(1 - a_1^{(2)}) \end{pmatrix}$$

$$D_1 = \begin{pmatrix} a_0^{(1)}(1 - a_0^{(1)}) & 0 \\ 0 & a_0^{(1)}(1 - a_0^{(1)}) \end{pmatrix}$$

$$\text{we can write } \delta_2 = \begin{pmatrix} \delta_0^{(2)} \\ \delta_1^{(2)} \end{pmatrix} = D_{(2)} \cdot c = \begin{pmatrix} a_0^{(2)}(1 - a_0^{(2)}) & 0 \\ 0 & a_1^{(2)}(1 - a_1^{(2)}) \end{pmatrix} \begin{pmatrix} a_0^{(2)} - T^0 \\ a_1^{(2)} - T^1 \end{pmatrix} = \begin{pmatrix} (a_0^{(2)} - T_0) a_0^{(2)} (1 - a_0^{(2)}) \\ (a_1^{(2)} - T_0) a_1^{(2)} (1 - a_1^{(2)}) \end{pmatrix}$$