R Code for Calculating the Relative Importance of Each Component in a PD Observed Score

Lara Stas

First, read in the following function by running the following lines.

Now, this function can be used to calculate the relative importance of each PD SRM component in a purely dyadic (observed) score. First, specify the estimated variance of each component (you can find this in the lavaan output under the header "Variances")

Legend:

- fam = family component
- i = individual component
- d = dyadic component
- roles: mo = mother; fa = father; oc = oldest child; yc = youngest child.

```
# Estimated variance of each component

fam <- 0.473
i.mo <- 0.129
i.fa <- 0.417
i.oc <- 0.099
i.yc <- 0
d.mofa <- 0.261
d.mooc <- 0.168
d.moyc <- 0.063
d.faoc <- 0.045
d.fayc <- 0.094
d.ycoc <- 0.473
```

Than, tell R by which components the PD scores are defined. For example, as can be seen in Figure 3 of the manuscript. The PD score of the mother-father dyad is a function of the family component, the individual component of the mother-father dyadic component.

```
# mother-father PD score
relvar(c1 = fam, c2 = i.mo, c3 = i.fa, c4 = d.mofa)

## % of variance explained by fam = 36.95
## % of variance explained by i.mo = 10.08
## % of variance explained by i.fa = 32.58
```

```
## % of variance explained by d.mofa = 20.39
# mother-oldest child PD score
relvar(c1 = fam, c2 = i.mo, c3 = i.oc, c4 = d.mooc)
## \% of variance explained by fam = 54.43
## % of variance explained by i.mo = 14.84
## \% of variance explained by i.oc = 11.39
## \% of variance explained by d.mooc = 19.33
# mother-youngest child PD score
relvar(c1 = fam, c2 = i.mo, c3 = i.yc, c4 = d.moyc)
## \% of variance explained by fam = 71.13
## \% of variance explained by i.mo = 19.4
## % of variance explained by i.yc = 0
## % of variance explained by d.moyc = 9.47
# father-oldest child PD score
relvar(c1 = fam, c2 = i.fa, c3 = i.oc, c4 = d.faoc)
## % of variance explained by fam = 45.74
## % of variance explained by i.fa = 40.33
## \% of variance explained by i.oc = 9.57
## \% of variance explained by d.faoc = 4.35
# father-youngest child PD score
relvar(c1 = fam, c2 = i.fa, c3 = i.yc, c4 = d.fayc)
## % of variance explained by fam = 48.07
## \% of variance explained by i.fa = 42.38
## % of variance explained by i.yc = 0
## % of variance explained by d.fayc = 9.55
# youngest-oldest child PD score
relvar(c1 = fam, c2 = i.yc, c3 = i.oc, c4 = d.ycoc)
## % of variance explained by fam = 45.26
## % of variance explained by i.yc = 0
## \% of variance explained by i.oc = 9.47
## % of variance explained by d.ycoc = 45.26
```