

**ANNEX I**

**SUMMARY OF PRODUCT CHARACTERISTICS**

## **1. NAME OF THE MEDICINAL PRODUCT**

Emtricitabine/Tenofovir disoproxil Zentiva 200 mg / 245 mg film-coated tablets

## **2. QUALITATIVE AND QUANTITATIVE COMPOSITION**

Each film-coated tablet contains 200 mg of emtricitabine and 245 mg of tenofovir disoproxil (equivalent to 291.5 mg of tenofovir disoproxil phosphate or 136 mg of tenofovir).

For the full list of excipients, see section 6.1.

## **3. PHARMACEUTICAL FORM**

Film-coated tablet.

Blue, oval biconvex film-coated tablet without half-score with dimensions approx. 19.35 x 9.75 mm.

## **4. CLINICAL PARTICULARS**

### **4.1 Therapeutic indications**

#### Treatment of HIV-1 infection

Emtricitabine/Tenofovir disoproxil Zentiva is indicated in antiretroviral combination therapy for the treatment of HIV-1 infected adults (see section 5.1).

Emtricitabine/Tenofovir disoproxil Zentiva is also indicated for the treatment of HIV-1 infected adolescents, with NRTI resistance or toxicities precluding the use of first line agents (see sections 4.2, 4.4 and 5.1).

#### Pre-exposure prophylaxis (PrEP)

Emtricitabine/Tenofovir disoproxil Zentiva is indicated in combination with safer sex practices for pre-exposure prophylaxis to reduce the risk of sexually acquired HIV-1 infection in adults and adolescents at high risk (see sections 4.2, 4.4 and 5.1).

### **4.2 Posology and method of administration**

Emtricitabine/Tenofovir disoproxil Zentiva should be initiated by a physician experienced in the management of HIV infection.

#### Posology

*Treatment of HIV in adults and adolescents aged 12 years and older, weighing at least 35 kg*  
One tablet, once daily.

*Prevention of HIV in adults and adolescents aged 12 years and older, weighing at least 35 kg*  
One tablet, once daily.

Separate preparations of emtricitabine and tenofovir disoproxil are available for treatment of HIV-1 infection if it becomes necessary to discontinue or modify the dose of one of the components of Emtricitabine/Tenofovir disoproxil Zentiva. Please refer to the Summary of Product Characteristics for these medicinal products.

If a dose of Emtricitabine/Tenofovir disoproxil Zentiva is missed within 12 hours of the time it is usually taken, Emtricitabine/Tenofovir disoproxil Zentiva should be taken as soon as possible and the normal dosing schedule should be resumed. If a dose of Emtricitabine/Tenofovir disoproxil Zentiva is missed by more than 12 hours and it is almost time for the next dose, the missed dose should not be taken and the usual dosing schedule should be resumed.

If vomiting occurs within 1 hour of taking Emtricitabine/Tenofovir disoproxil Zentiva, another tablet should be taken. If vomiting occurs more than 1 hour after taking Emtricitabine/Tenofovir disoproxil Zentiva a second dose should not be taken.

### Special populations

#### *Elderly*

No dose adjustment is required (see section 5.2).

#### *Renal impairment*

Emtricitabine and tenofovir are eliminated by renal excretion and the exposure to emtricitabine and tenofovir increases in individuals with renal dysfunction (see sections 4.4 and 5.2).

#### Adults with renal impairment

Emtricitabine/tenofovir disoproxil should only be used in individuals with creatinine clearance (CrCl) <80 ml/min if the potential benefits are considered to outweigh the potential risks. See Table 1.

**Table 1: Dosing recommendations in adults with renal impairment**

	<b>Treatment of HIV-1 infection</b>	<b>Pre-exposure prophylaxis</b>
Mild renal impairment (CrCl 50-80 ml/min)	Limited data from clinical studies support once daily dosing (see section 4.4).	Limited data from clinical studies support once daily dosing in HIV-1 uninfected individuals with CrCl 60-80 ml/min. Use is not recommended in HIV-1 uninfected individuals with CrCl < 60 ml/min as it has not been studied in this population (see sections 4.4 and 5.2).
Moderate renal impairment (CrCl 30-49 ml/min)	Administration every 48 hours is recommended based on modelling of single-dose pharmacokinetic data for emtricitabine and tenofovir disoproxil in non-HIV infected subjects with varying degrees of renal impairment (see section 4.4).	Not recommended for use in this population.
Severe renal impairment (CrCl < 30 ml/min) and haemodialysis patients	Not recommended because appropriate dose reductions cannot be achieved with the combination tablet.	Not recommended for use in this population.

#### Paediatrics with renal impairment

Not recommended for use in individuals under the age of 18 years with renal impairment (see section 4.4).

#### *Hepatic impairment*

No dose adjustment is required in patients with hepatic impairment (see sections 4.4 and 5.2).

#### *Paediatric population*

The safety and efficacy of emtricitabine/tenofovir disoproxil in children under the age of 12 years have not been established (see section 5.2).

#### Method of administration

Oral administration. It is preferable that Emtricitabine/Tenofovir disoproxil Zentiva is taken with food. Emtricitabine/Tenofovir disoproxil Zentiva film-coated tablet can be disintegrated in approximately 100 ml of water, orange juice or grape juice and taken immediately.

### **4.3 Contraindications**

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

Use for pre-exposure prophylaxis in individuals with unknown or positive HIV-1 status.

### **4.4 Special warnings and precautions for use**

#### Patients with HIV-1 harbouring mutations

Emtricitabine/Tenofovir disoproxil Zentiva should be avoided in antiretroviral-experienced patients with HIV-1 harbouring the K65R mutation (see section 5.1).

#### Overall HIV-1 infection prevention strategy

Emtricitabine/Tenofovir disoproxil Zentiva is not always effective in preventing the acquisition of HIV-1. The time to onset of protection after commencing emtricitabine/tenofovir disoproxil is unknown.

Emtricitabine/Tenofovir disoproxil Zentiva should only be used for pre-exposure prophylaxis as part of an overall HIV-1 infection prevention strategy including the use of other HIV-1 prevention measures (e.g. consistent and correct condom use, knowledge of HIV-1 status, regular testing for other sexually transmitted infections).

#### *Risk of resistance with undetected HIV-1 infection*

Emtricitabine/Tenofovir disoproxil Zentiva should only be used to reduce the risk of acquiring HIV-1 in individuals confirmed to be HIV negative (see section 4.3). Individuals should be re-confirmed to be HIV negative at frequent intervals (e.g. at least every 3 months) using a combined antigen/antibody test while taking Emtricitabine/Tenofovir disoproxil Zentiva for pre-exposure prophylaxis.

Emtricitabine/Tenofovir disoproxil Zentiva alone does not constitute a complete regimen for the treatment of HIV-1 and HIV-1 resistance mutations have emerged in individuals with undetected HIV-1 infection who are only taking emtricitabine/tenofovir disoproxil.

If clinical symptoms consistent with acute viral infection are present and recent (< 1 month) exposures to HIV-1 are suspected, use of Emtricitabine/Tenofovir disoproxil Zentiva should be delayed for at least one month and HIV-1 status reconfirmed before starting Emtricitabine/Tenofovir disoproxil Zentiva for pre-exposure prophylaxis.

#### *Importance of adherence*

The effectiveness of emtricitabine/tenofovir disoproxil in reducing the risk of acquiring HIV-1 is strongly correlated with adherence as demonstrated by measurable drug levels in blood (see section 5.1). HIV-1 uninfected individuals should be counselled at frequent intervals to strictly adhere to the recommended Emtricitabine/Tenofovir disoproxil Zentiva daily dosing schedule.

## Patients with hepatitis B or C virus infection

HIV-1 infected patients with chronic hepatitis B or C treated with antiretroviral therapy are at an increased risk for severe and potentially fatal hepatic adverse reactions. Physicians should refer to current HIV treatment guidelines for the management of HIV infection in patients co-infected with hepatitis B virus (HBV) or hepatitis C virus (HCV).

The safety and efficacy of emtricitabine/tenofovir disoproxil for pre-exposure prophylaxis in patients with HBV or HCV infection has not been established.

In case of concomitant antiviral therapy for hepatitis B or C, please refer also to the relevant Summary of Product Characteristics for these medicinal products. See also under Use with ledipasvir and sofosbuvir or sofosbuvir and velpatasvir below.

Tenofovir disoproxil is indicated for the treatment of HBV and emtricitabine has shown activity against HBV in pharmacodynamic studies but the safety and efficacy of emtricitabine/tenofovir disoproxil have not been specifically established in patients with chronic HBV infection.

Discontinuation of Emtricitabine/Tenofovir disoproxil Zentiva therapy in patients infected with HBV may be associated with severe acute exacerbations of hepatitis. Patients infected with HBV who discontinue Emtricitabine/Tenofovir disoproxil Zentiva should be closely monitored with both clinical and laboratory follow-up for at least several months after stopping treatment. If appropriate, resumption of hepatitis B therapy may be warranted. In patients with advanced liver disease or cirrhosis, treatment discontinuation is not recommended since post-treatment exacerbation of hepatitis may lead to hepatic decompensation.

## Liver disease

The safety and efficacy of emtricitabine/tenofovir disoproxil have not been established in patients with significant underlying liver disorders. The pharmacokinetics of tenofovir has been studied in patients with hepatic impairment and no dose adjustment is required. The pharmacokinetics of emtricitabine has not been studied in patients with hepatic impairment. Based on minimal hepatic metabolism and the renal route of elimination for emtricitabine, it is unlikely that a dose adjustment would be required for Emtricitabine/Tenofovir disoproxil Zentiva in patients with hepatic impairment (see sections 4.2 and 5.2).

HIV-1 infected patients with pre-existing liver dysfunction, including chronic active hepatitis, have an increased frequency of liver function abnormalities during combination antiretroviral therapy (CART) and should be monitored according to standard practice. If there is evidence of worsening liver disease in such patients, interruption or discontinuation of treatment must be considered.

## Renal and bone effects in adults

### *Renal effects*

Emtricitabine and tenofovir are primarily excreted by the kidneys by a combination of glomerular filtration and active tubular secretion. Renal failure, renal impairment, elevated creatinine, hypophosphataemia and proximal tubulopathy (including Fanconi syndrome) have been reported with the use of tenofovir disoproxil (see section 4.8).

### *Renal monitoring*

Prior to initiating Emtricitabine/Tenofovir disoproxil Zentiva for the treatment of HIV-1 infection or for use in pre-exposure prophylaxis, it is recommended that creatinine clearance is calculated in all individuals.

In individuals without risk factors for renal disease, it is recommended that renal function (creatinine clearance and serum phosphate) is monitored after two to four weeks of use, after three months of use and every three to six months thereafter.

In individuals at risk for renal disease more frequent monitoring of renal function is required.

See also under *Co-administration of other medicinal products* below.

#### *Renal management in HIV-1 infected patients*

If serum phosphate is < 1.5 mg/dl (0.48 mmol/l) or creatinine clearance is decreased to < 50 ml/min in any patient receiving Emtricitabine/Tenofovir disoproxil Zentiva, renal function should be re-evaluated within one week, including measurements of blood glucose, blood potassium and urine glucose concentrations (see section 4.8, proximal tubulopathy). Consideration should be given to interrupting treatment with Emtricitabine/Tenofovir disoproxil Zentiva in patients with creatinine clearance decreased to < 50 ml/min or decreases in serum phosphate to < 1.0 mg/dl (0.32 mmol/l). Interrupting treatment with Emtricitabine/Tenofovir disoproxil Zentiva should also be considered in case of progressive decline of renal function when no other cause has been identified.

Renal safety with emtricitabine/tenofovir disoproxil has only been studied to a very limited degree in HIV-1 infected patients with impaired renal function (creatinine clearance < 80 ml/min). Dose interval adjustments are recommended for HIV-1 infected patients with creatinine clearance 30-49 ml/min (see section 4.2). Limited clinical study data suggest that the prolonged dose interval is not optimal and could result in increased toxicity and possibly inadequate response. Furthermore, in a small clinical study, a subgroup of patients with creatinine clearance between 50 and 60 ml/min who received tenofovir disoproxil in combination with emtricitabine every 24 hours had a 2-4-fold higher exposure to tenofovir and worsening of renal function (see section 5.2). Therefore, a careful benefit-risk assessment is needed when Emtricitabine/Tenofovir disoproxil Zentiva is used in patients with creatinine clearance < 60 ml/min, and renal function should be closely monitored. In addition, the clinical response to treatment should be closely monitored in patients receiving Emtricitabine/Tenofovir disoproxil Zentiva at a prolonged dosing interval. The use of Emtricitabine/Tenofovir disoproxil Zentiva is not recommended in patients with severe renal impairment (creatinine clearance < 30 ml/min) and in patients who require haemodialysis since appropriate dose reductions cannot be achieved with the combination tablet (see sections 4.2 and 5.2).

#### *Renal management in pre-exposure prophylaxis*

Emtricitabine/tenofovir disoproxil has not been studied in HIV-1 uninfected individuals with creatinine clearance < 60 ml/min and is therefore not recommended for use in this population. If serum phosphate is < 1.5 mg/dl (0.48 mmol/l) or creatinine clearance is decreased to < 60 ml/min in any individual receiving Emtricitabine/Tenofovir disoproxil Zentiva for pre-exposure prophylaxis, renal function should be re-evaluated within one week, including measurements of blood glucose, blood potassium and urine glucose concentrations (see section 4.8, proximal tubulopathy). Consideration should be given to interrupting use of Emtricitabine/Tenofovir disoproxil Zentiva in individuals with creatinine clearance decreased to < 60 ml/min or decreases in serum phosphate to < 1.0 mg/dl (0.32 mmol/l). Interrupting use of Emtricitabine/Tenofovir disoproxil Zentiva should also be considered in case of progressive decline of renal function when no other cause has been identified.

#### *Bone effects*

Bone abnormalities such as osteomalacia which can manifest as persistent or worsening bone pain and, which can infrequently contribute to fractures, may be associated with tenofovir disoproxil-induced proximal renal tubulopathy (see section 4.8).

If bone abnormalities are suspected or detected then appropriate consultation should be obtained.

#### *Treatment of HIV-1 infection*

Reductions of bone mineral density (BMD) have been observed with tenofovir disoproxil in randomized controlled clinical trials of duration up to 144 weeks in HIV or HBV-infected patients. These BMD decreases generally improved after treatment discontinuation.

In other studies (prospective and cross-sectional), the most pronounced decreases in BMD were seen in patients treated with tenofovir disoproxil as part of a regimen containing a boosted protease inhibitor. Overall in view of the bone abnormalities associated with tenofovir disoproxil and the

limitations of long-term data on the impact of tenofovir disoproxil on bone health and fracture risk, alternative treatment regimens should be considered for patients with osteoporosis or with a history of bone fractures.

#### *Pre-exposure prophylaxis*

In clinical studies of HIV-1 uninfected individuals, small decreases in BMD were observed. In a study of 498 men, the mean changes from baseline to week 24 in BMD ranged from 0.4% to 1.0% across hip, spine, femoral neck and trochanter in men who received daily emtricitabine/tenofovir disoproxil prophylaxis (n = 247) vs. placebo (n = 251).

#### Renal and bone effects in the paediatric population

There are uncertainties associated with the long-term renal and bone effects of tenofovir disoproxil during the treatment of HIV-1 infection in the paediatric population and the long-term renal and bone effects of emtricitabine/tenofovir disoproxil when used for pre-exposure prophylaxis in uninfected adolescents (see section 5.1). Moreover, the reversibility of renal toxicity after cessation of tenofovir disoproxil for treatment of HIV-1 or after cessation of emtricitabine/tenofovir disoproxil for pre-exposure prophylaxis cannot be fully ascertained.

A multidisciplinary approach is recommended to weigh the benefit/risk balance of the use of Emtricitabine/Tenofovir disoproxil Zentiva for the treatment of HIV-1 infection or for pre-exposure prophylaxis, decide the appropriate monitoring during treatment (including decision for treatment withdrawal) and consider the need for supplementation on a case by case basis.

When using Emtricitabine/Tenofovir disoproxil Zentiva for pre-exposure prophylaxis individuals should be reassessed at each visit to ascertain whether they remain at high risk of HIV-1 infection. The risk of HIV-1 infection should be balanced against the potential for renal and bone effects with long-term use of Emtricitabine/Tenofovir disoproxil Zentiva.

#### *Renal effects*

Renal adverse reactions consistent with proximal renal tubulopathy have been reported in HIV-1 infected paediatric patients aged 2 to < 12 years in clinical study GS-US-104-0352 (see sections 4.8 and 5.1).

#### *Renal monitoring*

Renal function (creatinine clearance and serum phosphate) should be evaluated prior to initiating Emtricitabine/Tenofovir disoproxil Zentiva for treatment of HIV-1 or for pre-exposure prophylaxis, and should be monitored during use as in adults (see above).

#### *Renal management*

If serum phosphate is confirmed to be < 3.0 mg/dl (0.96 mmol/l) in any paediatric patient receiving emtricitabine/tenofovir disoproxil, renal function should be re-evaluated within one week, including measurements of blood glucose, blood potassium and urine glucose concentrations (see section 4.8, proximal tubulopathy). If renal abnormalities are suspected or detected then consultation with a nephrologist should be obtained to consider interruption of Emtricitabine/Tenofovir disoproxil Zentiva use. Interrupting use of Emtricitabine/Tenofovir disoproxil Zentiva should also be considered in case of progressive decline of renal function when no other cause has been identified.

#### *Co-administration and risk of renal toxicity*

The same recommendations apply as in adults (see Co-administration of other medicinal products below).

### *Renal impairment*

The use of emtricitabine/tenofovir disoproxil is not recommended in individuals under the age of 18 years with renal impairment (see section 4.2).

Emtricitabine/Tenofovir disoproxil Zentiva should not be initiated in paediatric patients with renal impairment and should be discontinued in paediatric patients who develop renal impairment during Emtricitabine/Tenofovir disoproxil Zentiva use.

### *Bone effects*

Use of tenofovir disoproxil may cause a reduction in BMD. The effects of tenofovir disoproxil-associated changes in BMD on long-term bone health and future fracture risk are uncertain (see section 5.1).

If bone abnormalities are detected or suspected during use of Emtricitabine/Tenofovir disoproxil Zentiva in any paediatric patient, consultation with an endocrinologist and/or nephrologist should be obtained.

### Weight and metabolic parameters

An increase in weight and in levels of blood lipids and glucose may occur during antiretroviral therapy. Such changes may in part be linked to disease control and life style. For lipids, there is in some cases evidence for a treatment effect, while for weight gain there is no strong evidence relating this to any particular treatment. For monitoring of blood lipids and glucose reference is made to established HIV treatment guidelines. Lipid disorders should be managed as clinically appropriate.

### Mitochondrial dysfunction following exposure *in utero*

Nucleos(t)ide analogues may impact mitochondrial function to a variable degree, which is most pronounced with stavudine, didanosine and zidovudine. There have been reports of mitochondrial dysfunction in HIV negative infants exposed *in utero* and/or postnatally to nucleoside analogues; these have predominantly concerned treatment with regimens containing zidovudine. The main adverse reactions reported are haematological disorders (anaemia, neutropenia) and metabolic disorders (hyperlactatemia, hyperlipasemia). These events have often been transitory. Late onset neurological disorders have been reported rarely (hypertonia, convulsion, abnormal behaviour). Whether such neurological disorders are transient or permanent is currently unknown. These findings should be considered for any child exposed *in utero* to nucleos(t)ide analogues, who present with severe clinical findings of unknown etiology, particularly neurologic findings. These findings do not affect current national recommendations to use antiretroviral therapy in pregnant women to prevent vertical transmission of HIV.

### Immune Reactivation Syndrome

In HIV infected patients with severe immune deficiency at the time of institution of CART, an inflammatory reaction to asymptomatic or residual opportunistic pathogens may arise and cause serious clinical conditions, or aggravation of symptoms. Typically, such reactions have been observed within the first few weeks or months of initiation of CART. Relevant examples are cytomegalovirus retinitis, generalised and/or focal mycobacterial infections, and *Pneumocystis jirovecii* pneumonia. Any inflammatory symptoms should be evaluated and treatment instituted when necessary. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported to occur in the setting of immune reactivation; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment.

### Opportunistic infections

HIV-1 infected patients receiving Emtricitabine/Tenofovir disoproxil Zentiva or any other antiretroviral therapy may continue to develop opportunistic infections and other complications of HIV infection, and therefore should remain under close clinical observation by physicians experienced in the treatment of patients with HIV associated diseases.



### Osteonecrosis

Although the aetiology is considered to be multifactorial (including corticosteroid use, alcohol consumption, severe immunosuppression, higher body mass index), cases of osteonecrosis have been reported particularly in patients with advanced HIV-disease and/or long-term exposure to CART. Patients should be advised to seek medical advice if they experience joint aches and pain, joint stiffness or difficulty in movement.

### Co-administration of other medicinal products

Use of Emtricitabine/Tenofovir disoproxil Zentiva should be avoided with concurrent or recent use of a nephrotoxic medicinal product (see section 4.5). If concomitant use with nephrotoxic agents is unavoidable, renal function should be monitored weekly.

Cases of acute renal failure after initiation of high dose or multiple non-steroidal anti-inflammatory drugs (NSAIDs) have been reported in HIV-1 infected patients treated with tenofovir disoproxil and with risk factors for renal dysfunction. If Emtricitabine/Tenofovir disoproxil Zentiva is co-administered with an NSAID, renal function should be monitored adequately.

A higher risk of renal impairment has been reported in HIV-1 infected patients receiving tenofovir disoproxil in combination with a ritonavir or cobicistat boosted protease inhibitor. Close monitoring of renal function is required in these patients (see section 4.5). In HIV-1 infected patients with renal risk factors, the co-administration of tenofovir disoproxil with a boosted protease inhibitor should be carefully evaluated.

Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly with other medicinal products containing emtricitabine, tenofovir disoproxil, tenofovir alafenamide, or other cytidine analogues, such as lamivudine (see section 4.5). Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly with adefovir dipivoxil.

#### *Use with ledipasvir and sofosbuvir, sofosbuvir and velpatasvir or sofosbuvir, velpatasvir and voxilaprevir*

Co-administration of tenofovir disoproxil with ledipasvir/sofosbuvir, sofosbuvir/velpatasvir or sofosbuvir/velpatasvir/voxilaprevir has been shown to increase plasma concentrations of tenofovir, especially when used together with an HIV regimen containing tenofovir disoproxil and a pharmacokinetic enhancer (ritonavir or cobicistat).

The safety of tenofovir disoproxil when co-administered with ledipasvir/sofosbuvir, sofosbuvir/velpatasvir or sofosbuvir/velpatasvir/voxilaprevir and a pharmacokinetic enhancer has not been established. The potential risks and benefits associated with co-administration should be considered, particularly in patients at increased risk of renal dysfunction. Patients receiving ledipasvir/sofosbuvir, sofosbuvir/velpatasvir or sofosbuvir/velpatasvir/voxilaprevir concomitantly with tenofovir disoproxil and a boosted HIV protease inhibitor should be monitored for adverse reactions related to tenofovir disoproxil.

#### *Co-administration of tenofovir disoproxil and didanosine*

Co-administration is not recommended (see section 4.5).

### Triple nucleoside therapy

There have been reports of a high rate of virological failure and of emergence of resistance at an early stage in HIV-1 infected patients when tenofovir disoproxil was combined with lamivudine and abacavir as well as with lamivudine and didanosine as a once daily regimen. There is close structural similarity between lamivudine and emtricitabine and similarities in the pharmacokinetics and pharmacodynamics of these two agents. Therefore, the same problems may be seen if Emtricitabine/Tenofovir disoproxil Zentiva is administered with a third nucleoside analogue.

## Elderly

Emtricitabine/tenofovir disoproxil has not been studied in individuals over the age of 65 years. Individuals over the age of 65 years are more likely to have decreased renal function, therefore caution should be exercised when administering Emtricitabine/Tenofovir disoproxil Zentiva to older people.

## Excipients

This medicinal product contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

## **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in adults.

As Emtricitabine/Tenofovir disoproxil Zentiva contains emtricitabine and tenofovir disoproxil, any interactions that have been identified with these agents individually may occur with Emtricitabine/Tenofovir disoproxil Zentiva. Interaction studies have only been performed in adults. The steady-state pharmacokinetics of emtricitabine and tenofovir were unaffected when emtricitabine and tenofovir disoproxil were administered together versus each medicinal product dosed alone. *In vitro* and clinical pharmacokinetic interaction studies have shown the potential for CYP450 mediated interactions involving emtricitabine and tenofovir disoproxil with other medicinal products is low.

### Concomitant use not recommended

Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly with other medicinal products containing emtricitabine, tenofovir disoproxil, tenofovir alafenamide or other cytidine analogues, such as lamivudine (see section 4.4). Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly with adefovir dipivoxil.

### *Didanosine*

The co-administration of Emtricitabine/Tenofovir disoproxil Zentiva and didanosine is not recommended (see section 4.4 and Table 2).

### *Renally eliminated medicinal products*

Since emtricitabine and tenofovir are primarily eliminated by the kidneys, co-administration of Emtricitabine/Tenofovir disoproxil Zentiva with medicinal products that reduce renal function or compete for active tubular secretion (e.g. cidofovir) may increase serum concentrations of emtricitabine, tenofovir and/or the co-administered medicinal products.

Use of Emtricitabine/Tenofovir disoproxil Zentiva should be avoided with concurrent or recent use of a nephrotoxic medicinal product. Some examples include, but are not limited to, aminoglycosides, amphotericin B, foscarnet, ganciclovir, pentamidine, vancomycin, cidofovir or interleukin-2 (see section 4.4).

### Other interactions

Interactions between Emtricitabine/Tenofovir disoproxil Zentiva or its individual component(s) and other medicinal products are listed in Table 2 below (increase is indicated as “↑”, decrease as “↓”, no change as “↔”, twice daily as “b.i.d.” and once daily as “q.d.”). If available, 90% confidence intervals are shown in parentheses.

**Table 2: Interactions between Emtricitabine/Tenofovir disoproxil Zentiva or its individual component(s) and other medicinal products**

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
<b>ANTI-INFECTIVES</b>		
<b>Antiretrovirals</b>		
<b>Protease inhibitors</b>		
Atazanavir / ritonavir / tenofovir disoproxil (300 mg q.d./100 mg q.d./245 mg q.d.)	Atazanavir: AUC: ↓ 25% (↓ 42 to ↓ 3) C <sub>max</sub> : ↓ 28% (↓ 50 to ↑ 5) C <sub>min</sub> : ↓ 26% (↓ 46 to ↑ 10)  Tenofovir: AUC: ↑ 37% C <sub>max</sub> : ↑ 34% C <sub>min</sub> : ↑ 29%	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate tenofovir associated adverse events, including renal disorders. Renal function should be closely monitored (see section 4.4).
Atazanavir/ritonavir/emtricitabine	Interaction not studied.	
Darunavir / ritonavir / tenofovir disoproxil (300 mg q.d./100 mg q.d./ 245 mg q.d.)	Darunavir: AUC: ↔ C <sub>min</sub> : ↔  Tenofovir: AUC: ↑ 22% C <sub>min</sub> : ↑ 37%	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate tenofovir associated adverse events, including renal disorders. Renal function should be closely monitored (see section 4.4).
Darunavir/ritonavir/emtricitabine	Interaction not studied.	
Lopinavir/ritonavir / tenofovir disoproxil (400 mg b.i.d. / 100 mg b.i.d / 245 mg q.d.)	Lopinavir/ritonavir: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↔  Tenofovir: AUC: ↑ 32% (↑ 25 to ↑ 38) C <sub>max</sub> : ↔ C <sub>min</sub> : ↑ 51% (↑ 37 to ↑ 66)	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate tenofovir associated adverse events, including renal disorders. Renal function should be closely monitored (see section 4.4).
Lopinavir/ritonavir/emtricitabine	Interaction not studied.	

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
<b>NRTIs</b>		
Didanosine / tenofovir disoproxil	Co-administration of tenofovir disoproxil and didanosine results in a 40-60% increase in systemic exposure to didanosine.	Co-administration of Emtricitabine/Tenofovir disoproxil Zentiva and didanosine is not recommended (see section 4.4).
Didanosine/emtricitabine	Interaction not studied.	Increased systemic exposure to didanosine may increase didanosine related adverse reactions. Rarely, pancreatitis and lactic acidosis, sometimes fatal, have been reported. Co-administration of tenofovir disoproxil and didanosine at a dose of 400 mg daily has been associated with a significant decrease in CD4 cell count, possibly due to an intracellular interaction increasing phosphorylated (i.e. active) didanosine. A decreased dosage of 250 mg didanosine co-administered with tenofovir disoproxil therapy has been associated with reports of high rates of virological failure within several tested combinations for the treatment of HIV-1 infection.
Lamivudine/tenofovir disoproxil	Lamivudine: AUC: ↓ 3% (↓ 8% to ↑ 15) C <sub>max</sub> : ↓ 24% (↓ 44 to ↓ 12) C <sub>min</sub> : NC  Tenofovir: AUC: ↓ 4% (↓ 15 to ↑ 8) C <sub>max</sub> : ↑ 102% (↓ 96 to ↑ 108) C <sub>min</sub> : NC	Lamivudine and Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly (see section 4.4).
Efavirenz/tenofovir disoproxil	Efavirenz: AUC: ↓ 4% (↓ 7 to ↓ 1) C <sub>max</sub> : ↓ 4% (↓ 9 to ↑ 2) C <sub>min</sub> : NC  Tenofovir: AUC: ↓ 1% (↓ 8 to ↑ 6) C <sub>max</sub> : ↑ 7% (↓ 6 to ↑ 22) C <sub>min</sub> : NC	No dose adjustment of efavirenz is required.

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
<b>ANTI-INFECTIVES</b>		
<b>Hepatitis B virus (HBV) antiviral agents</b>		
Adefovir dipivoxil /tenofovir disoproxil	Adefovir dipivoxil: AUC: ↓ 11% (↓ 14 to ↓ 7) C <sub>max</sub> : ↓ 7% (↓ 13 to ↓ 0) C <sub>min</sub> : NC Tenofovir: AUC: ↓ 2% (↓ 5 to ↑ 0) C <sub>max</sub> : ↓ 1% (↓ 7 to ↑ 6) C <sub>min</sub> : NC	Adefovir dipivoxil and Emtricitabine/Tenofovir disoproxil Zentiva should not be administered concomitantly (see section 4.4).
<b>Hepatitis C virus (HCV) antiviral agents</b>		
Ledipasvir/sofosbuvir (90 mg/400 mg q.d.) + atazanavir/ritonavir (300 mg q.d./100 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.) <sup>1</sup>	Ledipasvir: AUC: ↑ 96% (↑ 74 to ↑ 121) C <sub>max</sub> : ↑ 68% (↑ 54 to ↑ 84) C <sub>min</sub> : ↑ 118% (↑ 91 to ↑ 150)  Sofosbuvir: AUC: ↔ C <sub>max</sub> : ↔  GS-331007 <sup>2</sup> : AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↑ 42% (↑ 34 to ↑ 49)  Atazanavir: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↑ 63% (↑ 45 to ↑ 84)  Ritonavir: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↑ 45% (↑ 27 to ↑ 64)  Emtricitabine: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↔  Tenofovir: AUC: ↔ C <sub>max</sub> : ↑ 47% (↑ 37 to ↑ 58) C <sub>min</sub> : ↑ 47% (↑ 38 to ↑ 57)	Increased plasma concentrations of tenofovir resulting from co- administration of tenofovir disoproxil, ledipasvir/sofosbuvir and atazanavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with ledipasvir/sofosbuvir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.  The combination should be used with caution with frequent renal monitoring, if other alternatives are not available (see section 4.4).

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Ledipasvir/sofosbuvir (90 mg/400 mg q.d.) + darunavir/ritonavir (800 mg q.d./100 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.) <sup>1</sup>	<p>Ledipasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Sofosbuvir: AUC: ↓ 27% (↓ 35 to ↓ 18) C<sub>max</sub>: ↓ 37% (↓ 48 to ↓ 25)</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Darunavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Ritonavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↑ 48% (↑ 34 to ↑ 63)</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 50% (↑ 42 to ↑ 59) C<sub>max</sub>: ↑ 64% (↑ 54 to ↑ 74) C<sub>min</sub>: ↑ 59% (↑ 49 to ↑ 70)</p>	<p>Increased plasma concentrations of tenofovir resulting from co-administration of tenofovir disoproxil, ledipasvir/sofosbuvir and darunavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with ledipasvir/sofosbuvir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.</p> <p>The combination should be used with caution with frequent renal monitoring, if other alternatives are not available (see section 4.4).</p>

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Ledipasvir/sofosbuvir (90 mg/400 mg q.d.) + efavirenz/emtricitabine/tenofovir disoproxil (600 mg/200 mg/245 mg q.d.)	<p>Ledipasvir: AUC: ↓ 34% (↓ 41 to ↓ 25) C<sub>max</sub>: ↓ 34% (↓ 41 to ↑ 25) C<sub>min</sub>: ↓ 34% (↓ 43 to ↑ 24)</p> <p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Efavirenz: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 98% (↑ 77 to ↑ 123) C<sub>max</sub>: ↑ 79% (↑ 56 to ↑ 104) C<sub>min</sub>: ↑ 163% (↑ 137 to ↑ 197)</p>	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate adverse reactions associated with tenofovir disoproxil, including renal disorders. Renal function should be closely monitored (see section 4.4).
Ledipasvir/sofosbuvir (90 mg/400 mg q.d.) + emtricitabine/rilpivirine/tenofovir disoproxil (200 mg/25 mg/245 mg q.d.)	<p>Ledipasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Rilpivirine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 40% (↑ 31 to ↑ 50) C<sub>max</sub>: ↔ C<sub>min</sub>: ↑ 91% (↑ 74 to ↑ 110)</p>	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate adverse reactions associated with tenofovir disoproxil, including renal disorders. Renal function should be closely monitored (see section 4.4).

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Ledipasvir/sofosbuvir (90 mg/400 mg q.d.) + dolutegravir (50 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup> AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Ledipasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Dolutegravir AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 65% (↑ 59 to ↑ 71) C<sub>max</sub>: ↑ 61% (↑ 51 to ↑ 72) C<sub>min</sub>: ↑ 115% (↑ 105 to ↑ 126)</p>	No dose adjustment is required. The increased exposure of tenofovir could potentiate adverse reactions associated with tenofovir disoproxil, including renal disorders. Renal function should be closely monitored (see section 4.4).



Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + atazanavir/ritonavir (300 mg q.d./100 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↑ 42% (↑ 37 to ↑ 49)</p> <p>Velpatasvir: AUC: ↑ 142% (↑ 123 to ↑ 164) C<sub>max</sub>: ↑ 55% (↑ 41 to ↑ 71) C<sub>min</sub>: ↑ 301% (↑ 257 to ↑ 350)</p> <p>Atazanavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↑ 39% (↑ 20 to ↑ 61)</p> <p>Ritonavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↑ 29% (↑ 15 to ↑ 44)</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↔ C<sub>max</sub>: ↑ 55% (↑ 43 to ↑ 68) C<sub>min</sub>: ↑ 39% (↑ 31 to ↑ 48)</p>	<p>Increased plasma concentrations of tenofovir resulting from co-administration of tenofovir disoproxil, sofosbuvir/velpatasvir and atazanavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with sofosbuvir/velpatasvir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.</p> <p>The combination should be used with caution with frequent renal monitoring (see section 4.4).</p>

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + darunavir/ritonavir (800 mg q.d./100 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↓ 28% (↓ 34 to ↓ 20) C<sub>max</sub>: ↓ 38% (↓ 46 to ↓ 29)</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Velpatasvir: AUC: ↔ C<sub>max</sub>: ↓ 24% (↓ 35 to ↓ 11) C<sub>min</sub>: ↔</p> <p>Darunavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Ritonavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 39% (↑ 33 to ↑ 44) C<sub>max</sub>: ↑ 55% (↑ 45 to ↑ 66) C<sub>min</sub>: ↑ 52% (↑ 45 to ↑ 59)</p>	<p>Increased plasma concentrations of tenofovir resulting from co administration of tenofovir disoproxil, sofosbuvir/velpatasvir and darunavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with sofosbuvir/velpatasvir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.</p> <p>The combination should be used with caution with frequent renal monitoring (see section 4.4).</p>

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + lopinavir/ritonavir (800 mg/200 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↓ 29% (↓ 36 to ↓ 22) C<sub>max</sub>: ↓ 41% (↓ 51 to ↓ 29)</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Velpatasvir: AUC: ↔ C<sub>max</sub>: ↓ 30% (↓ 41 to ↓ 17) C<sub>min</sub>: ↑ 63% (↑ 43 to ↑ 85)</p> <p>Lopinavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Ritonavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↔ C<sub>max</sub>: ↑ 42% (↑ 27 to ↑ 57) C<sub>min</sub>: ↔</p>	<p>Increased plasma concentrations of tenofovir resulting from co-administration of tenofovir disoproxil, sofosbuvir/velpatasvir and lopinavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with sofosbuvir/velpatasvir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.</p> <p>The combination should be used with caution with frequent renal monitoring (see section 4.4).</p>

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + raltegravir (400 mg b.i.d) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Velpatasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Raltegravir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↓ 21% (↓ 58 to ↑ 48)</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 40% (↑ 34 to ↑ 45) C<sub>max</sub>: ↑ 46% (↑ 39 to ↑ 54) C<sub>min</sub>: ↑ 70% (↑ 61 to ↑ 79)</p>	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate adverse reactions associated with tenofovir disoproxil, including renal disorders. Renal function should be closely monitored (see section 4.4).
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + efavirenz/emtricitabine/tenofovir disoproxil (600 mg/200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↑ 38% (↑ 14 to ↑ 67)</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Velpatasvir: AUC: ↓ 53% (↓ 61 to ↓ 43) C<sub>max</sub>: ↓ 47% (↓ 57 to ↓ 36) C<sub>min</sub>: ↓ 57% (↓ 64 to ↓ 48)</p> <p>Efavirenz: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 81% (↑ 68 to ↑ 94) C<sub>max</sub>: ↑ 77% (↑ 53 to ↑ 104) C<sub>min</sub>: ↑ 121% (↑ 100 to ↑ 143)</p>	Concomitant administration of sofosbuvir/velpatasvir and efavirenz is expected to decrease plasma concentrations of velpatasvir. Co-administration of sofosbuvir/velpatasvir with efavirenz-containing regimens is not recommended.

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir (400 mg/100 mg q.d.) + emtricitabine/rilpivirine/tenofovir disoproxil (200 mg/25 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↔</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Velpatasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Rilpivirine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 40% (↑ 34 to ↑ 46) C<sub>max</sub>: ↑ 44% (↑ 33 to ↑ 55) C<sub>min</sub>: ↑ 84% (↑ 76 to ↑ 92)</p>	No dose adjustment is recommended. The increased exposure of tenofovir could potentiate adverse reactions associated with tenofovir disoproxil, including renal disorders. Renal function should be closely monitored (see section 4.4).

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir/velpatasvir/ voxilaprevir (400 mg/100 mg/100 mg+100 mg q.d.) <sup>3</sup> + darunavir (800 mg q.d.) + ritonavir (100 mg q.d.) + emtricitabine/tenofovir disoproxil (200 mg/245 mg q.d.)	<p>Sofosbuvir: AUC: ↔ C<sub>max</sub>: ↓ 30% C<sub>min</sub>: N/A</p> <p>GS-331007<sup>2</sup>: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: N/A</p> <p>Velpatasvir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Voxilaprevir: AUC: ↑ 143% C<sub>max</sub>: ↑ 72% C<sub>min</sub>: ↑ 300%</p> <p>Darunavir: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↓ 34%</p> <p>Ritonavir: AUC: ↑ 45% C<sub>max</sub>: ↑ 60% C<sub>min</sub>: ↔</p> <p>Emtricitabine: AUC: ↔ C<sub>max</sub>: ↔ C<sub>min</sub>: ↔</p> <p>Tenofovir: AUC: ↑ 39% C<sub>max</sub>: ↑ 48% C<sub>min</sub>: ↑ 47%</p>	<p>Increased plasma concentrations of tenofovir resulting from co-administration of tenofovir disoproxil, sofosbuvir/velpatasvir/voxilaprevir and darunavir/ritonavir may increase adverse reactions related to tenofovir disoproxil, including renal disorders. The safety of tenofovir disoproxil when used with sofosbuvir/velpatasvir/voxilaprevir and a pharmacokinetic enhancer (e.g. ritonavir or cobicistat) has not been established.</p> <p>The combination should be used with caution with frequent renal monitoring (see section 4.4).</p>

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
Sofosbuvir (400 mg q.d.) + efavirenz/emtricitabine/tenofovir disoproxil (600 mg/200 mg/245 mg q.d.)	Sofosbuvir: AUC: ↔ C <sub>max</sub> : ↓ 19% (↓ 40 to ↑ 10)  GS-331007 <sup>2</sup> : AUC: ↔ C <sub>max</sub> : ↓ 23% (↓ 30 to ↑ 16)  Efavirenz: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↔  Emtricitabine: AUC: ↔ C <sub>max</sub> : ↔ C <sub>min</sub> : ↔  Tenofovir: AUC: ↔ C <sub>max</sub> : ↑ 25% (↑ 8 to ↑ 45) C <sub>min</sub> : ↔	No dose adjustment is required.
Ribavirin/tenofovir disoproxil	Ribavirin: AUC: ↑ 26% (↑ 20 to ↑ 32) C <sub>max</sub> : ↓ 5% (↓ 11 to ↑ 1) C <sub>min</sub> : NC	No dose adjustment of ribavirin is required.
<b>Herpes virus antiviral agents</b>		
Famciclovir/emtricitabine	Famciclovir: AUC: ↓ 9% (↓ 16 to ↓ 1) C <sub>max</sub> : ↓ 7% (↓ 22 to ↑ 11) C <sub>min</sub> : NC  Emtricitabine: AUC: ↓ 7% (↓ 13 to ↓ 1) C <sub>max</sub> : ↓ 11% (↓ 20 to ↑ 1) C <sub>min</sub> : NC	No dose adjustment of famciclovir is required.
<b>Antimycobacterials</b>		
Rifampicin/tenofovir disoproxil	Tenofovir: AUC: ↓ 12% (↓ 16 to ↓ 8) C <sub>max</sub> : ↓ 16% (↓ 22 to ↓ 10) C <sub>min</sub> : ↓ 15% (↓ 12 to ↓ 9)	No dose adjustment is required.
<b>ORAL CONTRACEPTIVES</b>		
Norgestimate/ethinyl oestradiol/tenofovir disoproxil	Norgestimate: AUC: ↓ 4% (↓ 32 to ↑ 34) C <sub>max</sub> : ↓ 5% (↓ 27 to ↑ 24) C <sub>min</sub> : NC  Ethinyl oestradiol: AUC: ↓ 4% (↓ 9 to ↑ 0) C <sub>max</sub> : ↓ 6% (↓ 13 to ↑ 0) C <sub>min</sub> : ↓ 2% (↓ 9 to ↑ 6)	No dose adjustment of norgestimate/ethinyl oestradiol is required.

Medicinal product by therapeutic areas	Effects on drug levels Mean percent change in AUC, C <sub>max</sub> , C <sub>min</sub> with 90% confidence intervals if available (mechanism)	Recommendation concerning co-administration with Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine 200 mg, tenofovir disoproxil 245 mg)
<b>IMMUNOSUPPRESSANTS</b>		
Tacrolimus/tenofovir disoproxil /emtricitabine	<p>Tacrolimus: AUC: ↑ 4% (↓ 3 to ↑ 11) C<sub>max</sub>: ↑ 3% (↓ 3 to ↑ 9) C<sub>min</sub>: NC</p> <p>Emtricitabine: AUC: ↓ 5% (↓ 9 to ↓ 1) C<sub>max</sub>: ↓ 11% (↓ 17 to ↓ 5) C<sub>min</sub>: NC</p> <p>Tenofovir: AUC: ↑ 6% (↓ 1 to ↑ 13) C<sub>max</sub>: ↑ 13% (↑ 1 to ↑ 27) C<sub>min</sub>: NC</p>	No dose adjustment of tacrolimus is required.
<b>NARCOTIC ANALGESICS</b>		
Methadone/tenofovir disoproxil	<p>Methadone: AUC: ↑ 5% (↓ 2 to ↑ 13) C<sub>max</sub>: ↑ 5% (↓ 3 to ↑ 14) C<sub>min</sub>: NC</p>	No dose adjustment of methadone is required.

NC = not calculated.

N/A = not applicable.

<sup>1</sup> Data generated from simultaneous dosing with ledipasvir/sofosbuvir. Staggered administration (12 hours apart) provided similar results.

<sup>2</sup> The predominant circulating metabolite of sofosbuvir.

<sup>3</sup> Study conducted with additional voxilaprevir 100 mg to achieve voxilaprevir exposures expected in HCV-infected patients.

## 4.6 Fertility, pregnancy and lactation

### Pregnancy

A large amount of data on pregnant women (more than 1,000 pregnancy outcomes) indicate no malformations or foetal/neonatal toxicity associated with emtricitabine and tenofovir disoproxil. Animal studies on emtricitabine and tenofovir disoproxil do not indicate reproductive toxicity (see section 5.3). Therefore the use of Emtricitabine/Tenofovir disoproxil Zentiva may be considered during pregnancy, if necessary.

### Breast-feeding

Emtricitabine and tenofovir have been shown to be excreted in human milk. There is insufficient information on the effects of emtricitabine and tenofovir in newborns/infants. Therefore Emtricitabine/Tenofovir disoproxil Zentiva should not be used during breast-feeding.

In order to avoid transmission of HIV to the infant it is recommended that women living with HIV do not breast-feed their infants.

### Fertility

No human data on the effect of emtricitabine/tenofovir disoproxil are available. Animal studies do not indicate harmful effects of emtricitabine or tenofovir disoproxil on fertility.



## 4.7 Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed. However, individuals should be informed that dizziness has been reported during treatment with both emtricitabine and tenofovir disoproxil.

## 4.8 Undesirable effects

### Summary of the safety profile

#### *HIV-1 infection*

The most frequently reported adverse reactions considered possibly or probably related to emtricitabine and/or tenofovir disoproxil were nausea (12%) and diarrhoea (7%) in an open-label randomised clinical study in adults (GS-01-934, see section 5.1). The safety profile of emtricitabine and tenofovir disoproxil in this study was consistent with the previous experience with these agents when each was administered with other antiretroviral agents.

#### *Pre-exposure prophylaxis*

No new adverse reactions to emtricitabine/tenofovir disoproxil were identified from two randomised placebo-controlled studies (iPrEx, Partners PrEP) in which 2,830 HIV-1 uninfected adults received emtricitabine/tenofovir disoproxil once daily for pre-exposure prophylaxis. Patients were followed for a median of 71 weeks and 87 weeks, respectively. The most frequent adverse reaction reported in the emtricitabine/tenofovir disoproxil group in the iPrEx study was headache (1%).

### Tabulated summary of adverse reactions

The adverse reactions considered at least possibly related to treatment with the components of emtricitabine/tenofovir disoproxil from clinical study and post-marketing experience in HIV-1 infected patients are listed in Table 3, below, by body system organ class and frequency. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness. Frequencies are defined as very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ) or rare ( $\geq 1/10,000$  to  $< 1/1,000$ ).

**Table 3: Tabulated summary of adverse reactions associated with the individual components of emtricitabine / tenofovir disoproxil based on clinical study and post-marketing experience**

Frequency	Emtricitabine	Tenofovir disoproxil
<i>Blood and lymphatic system disorders</i>		
Common	neutropenia	
Uncommon	anaemia <sup>2</sup>	
<i>Immune system disorders</i>		
Common	allergic reaction	
<i>Metabolism and nutrition disorders</i>		
Very common		hypophosphataemia <sup>1</sup>
Common	hyperglycaemia, hypertriglyceridaemia	
Uncommon		hypokalaemia <sup>1</sup>
Rare		lactic acidosis
<i>Psychiatric disorders</i>		
Common	insomnia, abnormal dreams	

Frequency	Emtricitabine	Tenofovir disoproxil
<i>Nervous system disorders</i>		
Very common	headache	dizziness
Common	dizziness	headache
<i>Gastrointestinal disorders</i>		
Very common	diarrhoea, nausea	diarrhoea, vomiting, nausea
Common	elevated amylase including elevated pancreatic amylase, elevated serum lipase, vomiting, abdominal pain, dyspepsia	abdominal pain, abdominal distension, flatulence
Uncommon		pancreatitis
<i>Hepatobiliary disorders</i>		
Common	elevated serum aspartate aminotransferase (AST) and/or elevated serum alanine aminotransferase (ALT), hyperbilirubinaemia	increased transaminases
Rare		hepatic steatosis, hepatitis
<i>Skin and subcutaneous tissue disorders</i>		
Very common		rash
Common	vesiculobullous rash, pustular rash, maculopapular rash, rash, pruritus, urticaria, skin discolouration (increased pigmentation) <sup>2</sup>	
Uncommon	angioedema <sup>3</sup>	
Rare		angioedema
<i>Musculoskeletal and connective tissue disorders</i>		
Very common	elevated creatine kinase	
Common		bone mineral density decreased
Uncommon		rhabdomyolysis <sup>1</sup> , muscular weakness <sup>1</sup>
Rare		osteomalacia (manifested as bone pain and infrequently contributing to fractures) <sup>1,3</sup> , myopathy <sup>1</sup>
<i>Renal and urinary disorders</i>		
Uncommon		increased creatinine, proteinuria, proximal renal tubulopathy including Fanconi syndrome
Rare		renal failure (acute and chronic), acute tubular necrosis, nephritis (including acute interstitial nephritis) <sup>3</sup> , nephrogenic diabetes insipidus

Frequency	Emtricitabine	Tenofovir disoproxil
<i>General disorders and administration site conditions</i>		
Very common		asthenia
Common	pain, asthenia	

<sup>1</sup> This adverse reaction may occur as a consequence of proximal renal tubulopathy. It is not considered to be causally associated with tenofovir disoproxil in the absence of this condition.

<sup>2</sup> Anaemia was common and skin discolouration (increased pigmentation) was very common when emtricitabine was administered to paediatric patients.

<sup>3</sup> This adverse reaction was identified through post-marketing surveillance but not observed in randomised controlled clinical studies in adults or paediatric HIV clinical studies for emtricitabine or in randomised controlled clinical studies or the tenofovir disoproxil expanded access program for tenofovir disoproxil. The frequency category was estimated from a statistical calculation based on the total number of patients exposed to emtricitabine in randomised controlled clinical studies (n = 1,563) or tenofovir disoproxil in randomised controlled clinical studies and the expanded access program (n = 7,319).

### Description of selected adverse reactions

#### *Renal impairment*

As emtricitabine / tenofovir disoproxil may cause renal damage monitoring of renal function is recommended (see section 4.4). Proximal renal tubulopathy generally resolved or improved after tenofovir disoproxil discontinuation. However, in some HIV-1 infected patients, declines in creatinine clearance did not completely resolve despite tenofovir disoproxil discontinuation. Patients at risk of renal impairment (such as patients with baseline renal risk factors, advanced HIV disease, or patients receiving concomitant nephrotoxic medications) are at increased risk of experiencing incomplete recovery of renal function despite tenofovir disoproxil discontinuation (see section 4.4).

#### *Lactic acidosis*

Cases of lactic acidosis have been reported with tenofovir disoproxil alone or in combination with other antiretrovirals. Patients with predisposing factors such as patients with decompensated liver disease, or patients receiving concomitant medications known to induce lactic acidosis are at increased risk of experiencing severe lactic acidosis during tenofovir disoproxil treatment, including fatal outcomes.

#### *Metabolic parameters*

Weight and levels of blood lipids and glucose may increase during antiretroviral therapy (see section 4.4).

#### *Immune Reactivation Syndrome*

In HIV infected patients with severe immune deficiency at the time of initiation of CART, an inflammatory reaction to asymptomatic or residual opportunistic infections may arise. Autoimmune disorders (such as Graves' disease and autoimmune hepatitis) have also been reported; however, the reported time to onset is more variable and these events can occur many months after initiation of treatment (see section 4.4).

#### *Osteonecrosis*

Cases of osteonecrosis have been reported, particularly in patients with generally acknowledged risk factors, advanced HIV disease or long-term exposure to CART. The frequency of this is unknown (see section 4.4).

### Paediatric population

Assessment of adverse reactions related to emtricitabine is based on experience in three paediatric studies (n = 169) where treatment-naïve (n = 123) and treatment-experienced (n = 46) paediatric HIV infected patients aged 4 months to 18 years were treated with emtricitabine in combination with other antiretroviral agents. In addition to the adverse reactions reported in adults, anaemia (9.5%) and skin

discolouration (31.8%) occurred more frequently in clinical trials in paediatric patients than in adults (see section 4.8, Tabulated summary of adverse reactions).

Assessment of adverse reactions related to tenofovir disoproxil is based on two randomised trials (studies GS-US 104-0321 and GS-US-104-0352) in 184 HIV-1 infected paediatric patients (aged 2 to < 18 years) who received treatment with tenofovir disoproxil (n = 93) or placebo/active comparator (n = 91) in combination with other antiretroviral agents for 48 weeks (see section 5.1). The adverse reactions observed in paediatric patients who received treatment with tenofovir disoproxil were consistent with those observed in clinical studies of tenofovir disoproxil in adults (see section 4.8 Tabulated summary of adverse reactions and 5.1).

Reductions in BMD have been reported in paediatric patients. In HIV-1 infected adolescents (aged 12 to < 18 years), the BMD Z-scores observed in subjects who received tenofovir disoproxil were lower than those observed in subjects who received placebo. In HIV-1 infected children (aged 2 to 15 years), the BMD Z-scores observed in subjects who switched to tenofovir disoproxil were lower than those observed in subjects who remained on their stavudine- or zidovudine-containing regimen (see sections 4.4 and 5.1).

In study GS-US-104-0352, 89 HIV-1 infected paediatric patients with a median age of 7 years (range 2 to 15 years) were exposed to tenofovir disoproxil for a median of 331 weeks. Eight of the 89 patients (9.0%) discontinued study drug due to renal adverse events. Five subjects (5.6%) had laboratory findings clinically consistent with proximal renal tubulopathy, 4 of whom discontinued tenofovir disoproxil therapy. Seven patients had estimated glomerular filtration rate (GFR) values between 70 and 90 ml/min/1.73 m<sup>2</sup>. Among them, 3 patients experienced a clinically meaningful decline in estimated GFR during therapy which improved after discontinuation of tenofovir disoproxil.

#### Other special populations

##### *Individuals with renal impairment*

Since tenofovir disoproxil can cause renal toxicity, close monitoring of renal function is recommended in any adults with renal impairment receiving Emtricitabine/Tenofovir disoproxil Zentiva (see sections 4.2, 4.4 and 5.2). The use of Emtricitabine/Tenofovir disoproxil Zentiva is not recommended in individuals under the age of 18 years with renal impairment (see sections 4.2 and 4.4).

##### *HIV/HBV or HCV co-infected patients*

The adverse reaction profile of emtricitabine and tenofovir disoproxil in a limited number of HIV-infected patients in study GS-01-934 who were co-infected with HBV (n = 13) or HCV (n = 26) was similar to that observed in patients infected with HIV without co-infection. However, as would be expected in this patient population, elevations in AST and ALT occurred more frequently than in the general HIV infected population.

##### *Exacerbations of hepatitis after discontinuation of treatment*

In HBV infected patients, clinical and laboratory evidence of hepatitis have occurred after discontinuation of treatment (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

## 4.9 Overdose

If overdose occurs the individual must be monitored for evidence of toxicity (see section 4.8), and standard supportive treatment applied as necessary.

Up to 30% of the emtricitabine dose and approximately 10% of the tenofovir dose can be removed by haemodialysis. It is not known whether emtricitabine or tenofovir can be removed by peritoneal dialysis.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antivirals for systemic use, antivirals for treatment of HIV infections, combinations, ATC code: J05AR03

#### Mechanism of action

Emtricitabine is a nucleoside analogue of cytidine. Tenofovir disoproxil is converted *in vivo* to tenofovir, a nucleoside monophosphate (nucleotide) analogue of adenosine monophosphate. Both emtricitabine and tenofovir have activity that is specific to human immunodeficiency virus (HIV-1 and HIV-2) and hepatitis B virus.

Emtricitabine and tenofovir are phosphorylated by cellular enzymes to form emtricitabine triphosphate and tenofovir diphosphate, respectively. *In vitro* studies have shown that both emtricitabine and tenofovir can be fully phosphorylated when combined together in cells. Emtricitabine triphosphate and tenofovir diphosphate competitively inhibit HIV-1 reverse transcriptase, resulting in DNA chain termination.

Both emtricitabine triphosphate and tenofovir diphosphate are weak inhibitors of mammalian DNA polymerases and there was no evidence of toxicity to mitochondria *in vitro* and *in vivo*.

#### Antiviral activity in vitro

Synergistic antiviral activity was observed with the combination of emtricitabine and tenofovir *in vitro*. Additive to synergistic effects were observed in combination studies with protease inhibitors, and with nucleoside and non-nucleoside analogue inhibitors of HIV reverse transcriptase.

#### Resistance

##### *In vitro*

Resistance has been seen *in vitro* and in some HIV-1 infected patients due to the development of the M184V/I mutation with emtricitabine or the K65R mutation with tenofovir. Emtricitabine-resistant viruses with the M184V/I mutation were cross-resistant to lamivudine, but retained sensitivity to didanosine, stavudine, tenofovir and zidovudine. The K65R mutation can also be selected by abacavir or didanosine and results in reduced susceptibility to these agents plus lamivudine, emtricitabine and tenofovir. Tenofovir disoproxil should be avoided in patients with HIV-1 harbouring the K65R mutation. In addition, a K70E substitution in HIV-1 reverse transcriptase has been selected by tenofovir and results in low-level reduced susceptibility to abacavir, emtricitabine, lamivudine and tenofovir. HIV-1 expressing three or more thymidine analogue associated mutations (TAMs) that included either the M41L or L210W reverse transcriptase mutation showed reduced susceptibility to tenofovir disoproxil.

##### *In vivo – treatment of HIV-1*

In an open-label randomised clinical study (GS-01-934) in antiretroviral-naïve patients, genotyping was performed on plasma HIV-1 isolates from all patients with confirmed HIV RNA > 400 copies/ml at weeks 48, 96 or 144 or at the time of early study drug discontinuation. As of week 144:

- The M184V/I mutation developed in 2/19 (10.5%) isolates analysed from patients in the emtricitabine/tenofovir disoproxil/efavirenz group and in 10/29 (34.5%) isolates analysed from the lamivudine/zidovudine/efavirenz group (p-value < 0.05, Fisher's Exact test comparing the emtricitabine/tenofovir disoproxil group to the lamivudine/zidovudine group among all patients).
- No virus analysed contained the K65R or K70E mutation.
- Genotypic resistance to efavirenz, predominantly the K103N mutation, developed in virus from 13/19 (68%) patients in the emtricitabine/tenofovir disoproxil/efavirenz group and in virus from 21/29 (72%) patients in the comparative group.

#### *In vivo – pre-exposure prophylaxis*

Plasma samples from 2 clinical studies of HIV-1 uninfected subjects, iPrEx and Partners PrEP, were analysed for 4 HIV-1 variants expressing amino acid substitutions (i.e. K65R, K70E, M184V, and M184I) that potentially confer resistance to tenofovir or emtricitabine. In the iPrEx clinical study, no HIV-1 variants expressing K65R, K70E, M184V, or M184I were detected at the time of seroconversion among subjects who became infected with HIV-1 after enrollment in the study. In 3 of 10 subjects who had acute HIV infection at study enrollment, M184I and M184V mutations were detected in the HIV of 2 of 2 subjects in the emtricitabine/tenofovir disoproxil group and 1 of 8 subjects in the placebo group.

In the Partners PrEP clinical study, no HIV-1 variants expressing K65R, K70E, M184V, or M184I were detected at the time of seroconversion among subjects who became infected with HIV-1 during the study. In 2 of 14 subjects who had acute HIV infection at study enrollment, the K65R mutation was detected in the HIV of 1 of 5 subjects in the tenofovir disoproxil 245 mg group and the M184V mutation (associated with resistance to emtricitabine) was detected in the HIV of 1 of 3 subjects in the emtricitabine/tenofovir disoproxil group.

#### Clinical data

##### *Treatment of HIV-1 infection*

In an open-label randomised clinical study (GS-01-934), antiretroviral-naïve HIV-1 infected adult patients received either a once daily regimen of emtricitabine, tenofovir disoproxil and efavirenz (n = 255) or a fixed combination of lamivudine and zidovudine administered twice daily and efavirenz once daily (n = 254). Patients in the emtricitabine and tenofovir disoproxil group were given emtricitabine/tenofovir disoproxil and efavirenz from week 96 to week 144. At baseline the randomised groups had similar median plasma HIV-1 RNA (5.02 and 5.00 log<sub>10</sub> copies/ml) and CD4 counts (233 and 241 cells/mm<sup>3</sup>). The primary efficacy endpoint for this study was the achievement and maintenance of confirmed HIV-1 RNA concentrations < 400 copies/ml over 48 weeks. Secondary efficacy analyses over 144 weeks included the proportion of patients with HIV-1 RNA concentrations < 400 or < 50 copies/ml, and change from baseline in CD4 cell count.

The 48-week primary endpoint data showed that the combination of emtricitabine, tenofovir disoproxil and efavirenz provided superior antiviral efficacy as compared with the fixed combination of lamivudine and zidovudine with efavirenz as shown in Table 4. The 144 week secondary endpoint data are also presented in Table 4.

**Table 4: 48- and 144-week efficacy data from study GS-01-934 in which emtricitabine, tenofovir disoproxil and efavirenz were administered to antiretroviral-naïve patients with HIV-1 infection**

	<b>GS-01-934 Treatment for 48 weeks</b>		<b>GS-01-934 Treatment for 144 weeks</b>	
	Emtricitabine / tenofovir disoproxil / efavirenz	Lamivudine / zidovudine / efavirenz	Emtricitabine / tenofovir disoproxil / efavirenz*	Lamivudine / zidovudine / efavirenz
HIV-1 RNA < 400 copies/ml (TLOVR)	84% (206/244)	73% (177/243)	71% (161/227)	58% (133/229)
p-value	0.002**		0.004**	
% difference (95% CI)	11% (4% to 19%)		13% (4% to 22%)	
HIV-1 RNA < 50 copies/ml (TLOVR)	80% (194/244)	70% (171/243)	64% (146/227)	56% (130/231)
p-value	0.021**		0.082**	
% difference (95% CI)	9% (2% to 17%)		8% (-1% to 17%)	
Mean change from baseline in CD4 cell count (cells/mm <sup>3</sup> )	+190	+158	+312	+271
p-value	0.002 <sup>a</sup>		0.089 <sup>a</sup>	
Difference (95% CI)	32 (9 to 55)		41 (4 to 79)	

\* Patients receiving emtricitabine, tenofovir disoproxil and efavirenz were given emtricitabine/tenofovir disoproxil plus efavirenz from week 96 to 144.

\*\* The p-value based on the Cochran-Mantel-Haenszel Test stratified for baseline CD4 cell count.

TLOVR = Time to Loss of Virologic Response

<sup>a</sup> Van Elteren Test

In a randomised clinical study (M02-418), 190 antiretroviral-naïve adults were treated once daily with emtricitabine and tenofovir disoproxil in combination with lopinavir/ritonavir given once or twice daily. At 48 weeks, 70% and 64% of patients demonstrated HIV-1 RNA < 50 copies/ml with the once and twice daily regimens of lopinavir/ritonavir, respectively. The mean changes in CD4 cell count from baseline were +185 cells/mm<sup>3</sup> and +196 cells/mm<sup>3</sup>, respectively.

Limited clinical experience in patients co-infected with HIV and HBV suggests that treatment with emtricitabine or tenofovir disoproxil in antiretroviral combination therapy to control HIV infection results in a reduction in HBV DNA (3 log<sub>10</sub> reduction or 4 – 5 log<sub>10</sub> reduction, respectively) (see section 4.4).

#### *Pre-exposure prophylaxis*

The iPrEx study (CO-US-104-0288) evaluated emtricitabine/tenofovir disoproxil or placebo in 2,499 HIV uninfected men (or transgender women) who have sex with men and who were considered at high risk for HIV infection. Subjects were followed for 4,237 person-years. Baseline characteristics are summarised in Table 5.

**Table 5: Study population from study CO-US-104-0288 (iPrEx)**

	<b>Placebo (n = 1248)</b>	<b>Emtricitabine/tenofovir disoproxil (n = 1251)</b>
<b>Age (Yrs), Mean (SD)</b>	27 (8.5)	27 (8.6)
<b>Race, N (%)</b>		
Black/African American	97 (8)	117 (9)
White	208 (17)	223 (18)
Mixed/Other	878 (70)	849 (68)
Asian	65 (5)	62 (5)
<b>Hispanic/Latino Ethnicity, N (%)</b>	906 (73)	900 (72)
<b>Sexual Risk Factors at Screening</b>		
Number of Partners Previous 12 Weeks, Mean (SD)	18 (43)	18 (35)
URAI Previous 12 Weeks, N (%)	753 (60)	732 (59)
URAI with HIV+ (or unknown status) Partner Previous 6 Mos, N (%)	1009 (81)	992 (79)
Involved in Transactional Sex Last 6 Month, N (%)	510 (41)	517 (41)
Known HIV+ Partner Last 6 Months, N (%)	32 (3)	23 (2)
Syphilis Seroreactivity, N (%)	162/1239 (13)	164/1240 (13)
Serum Herpes Simplex Virus Type 2 Infection, N (%)	430/1243 (35)	458/1241 (37)
Urine Leukocyte Esterase Positive, N (%)	22 (2)	23 (2)

URAI = unprotected receptive anal intercourse

The incidences of HIV seroconversion overall and in the subset reporting unprotected receptive anal intercourse are shown in Table 6. Efficacy was strongly correlated with adherence as assessed by detection of plasma or intracellular drug levels in a case-control study (Table 7).

**Table 6: Efficacy in study CO-US-104-0288 (iPrEx)**

	Placebo	Emtricitabine / tenofovir disoproxil	P-value <sup>a, b</sup>
mITT Analysis			
Seroconversions / N	83 / 1217	48 / 1224	0.002
Relative Risk Reduction (95% CI) <sup>b</sup>	42% (18%, 60%)		
URAI Within 12 Weeks Prior to Screening, mITT Analysis			
Seroconversions / N	72 / 753	34 / 732	0.0349
Relative Risk Reduction (95% CI) <sup>b</sup>	52% (28%, 68%)		

<sup>a</sup> P-values by logrank test. P-values for URAI refer to the null hypothesis that efficacy differed between subgroup strata (URAI, no URAI).

<sup>b</sup> Relative risk reduction calculated for mITT based on incident seroconversion, i.e., occurring post-baseline through first post-treatment visit (approximately 1 month after last study drug dispensation).



**Table 7: Efficacy and adherence in study CO-US-104-0288 (iPrEx, matched case-control analysis)**

<b>Cohort</b>	<b>Drug Detected</b>	<b>Drug Not Detected</b>	<b>Relative Risk Reduction (2-sided 95% CI)<sup>a</sup></b>
HIV-Positive Subjects	4 (8%)	44 (92%)	94% (78%, 99%)
HIV-Negative Matched Control Subjects	63 (44%)	81 (56%)	—

<sup>a</sup> Relative risk reduction calculated on incident (post-baseline) seroconversion from the double-blind treatment period and through the 8-week follow-up period. Only samples from subjects randomized to emtricitabine/tenofovir disoproxil were evaluated for detectable plasma or intracellular tenofovir disoproxil-DP levels.

The Partners PrEP clinical study (CO-US-104-0380) evaluated emtricitabine/tenofovir disoproxil, tenofovir disoproxil 245 mg, or placebo in 4,758 HIV-uninfected subjects from Kenya or Uganda in serodiscordant heterosexual couples. Subjects were followed for 7,830 person-years. Baseline characteristics are summarised in Table 8.

**Table 8: Study population from study CO-US-104-0380 (Partners PrEP)**

	<b>Placebo (n = 1,584)</b>	<b>Tenofovir disoproxil 245 mg (n = 1,584)</b>	<b>Emtricitabine / tenofovir disoproxil (n = 1,579)</b>
<b>Age (Yrs), Median (Q1, Q3)</b>	34 (28, 40)	33 (28, 39)	33 (28, 40)
<b>Gender, N (%)</b>			
Male	963 (61)	986 (62)	1,013 (64)
Female	621 (39)	598 (38)	566 (36)
<b>Key Couple Characteristics, N (%) or Median (Q1, Q3)</b>			
Married to study partner	1,552 (98)	1,543 (97)	1,540 (98)
Years living with study partner	7.1 (3.0, 14.0)	7.0 (3.0, 13.5)	7.1 (3.0, 14.0)
Years aware of discordant status	0.4 (0.1, 2.0)	0.5 (0.1, 2.0)	0.4 (0.1, 2.0)

The incidence of HIV seroconversion is shown in Table 9. The rate of HIV-1 seroconversion in males was 0.24/100 person-years of emtricitabine/tenofovir disoproxil exposure and the rate of HIV-1 seroconversion in females was 0.95/100 person-years of emtricitabine/tenofovir disoproxil exposure. Efficacy was strongly correlated with adherence as assessed by detection of plasma or intracellular drug levels and was higher among substudy participants who received active adherence counselling and as show in Table 10.

**Table 9: Efficacy in study CO-US-104-0380 (Partners PrEP)**

	<b>Placebo</b>	<b>Tenofovir disoproxil 245 mg</b>	<b>Emtricitabine / tenofovir disoproxil</b>
Seroconversions / N <sup>a</sup>	52 / 1,578	17 / 1,579	13 / 1,576
Incidence per 100 person-years (95% CI)	1.99 (1.49, 2.62)	0.65 (0.38, 1.05)	0.50 (0.27, 0.85)
Relative Risk Reduction (95% CI)	—	67% (44%, 81%)	75% (55%, 87%)

<sup>a</sup> Relative risk reduction calculated for mITT cohort based on incident (post-baseline) seroconversion. Comparisons for active study groups are made versus placebo.

**Table 10: Efficacy and adherence in study CO-US-104-0380 (Partners PrEP)**

Study Drug Quantification	Number with Tenofovir Detected/Total Samples (%)		Risk Estimate for HIV-1 Protection: Detection Versus No Detection of Tenofovir	
	Case	Cohort	Relative Risk Reduction (95% CI)	p-value
FTC/tenofovir disoproxil group <sup>a</sup>	3 / 12 (25%)	375 / 465 (81%)	90% (56%, 98%)	0.002
Tenofovir disoproxil group <sup>a</sup>	6 / 17 (35%)	363 / 437 (83%)	86% (67%, 95%)	< 0.001
Adherence Substudy	Adherence Substudy Participants <sup>b</sup>		Relative Risk Reduction (95% CI)	p-value
	Placebo	Tenofovir disoproxil 245 mg + Emtricitabine/tenofovir disoproxil		
Seroconversions / N <sup>b</sup>	14 / 404 (3.5%)	0 / 745 (0%)	100% (87%, 100%)	< 0.001

<sup>a</sup> 'Case' = HIV seroconverter; 'Cohort' = 100 randomly selected subjects from each of the tenofovir disoproxil 245 mg and emtricitabine/tenofovir disoproxil groups. Only Case or Cohort samples from subjects randomised to either tenofovir disoproxil 245 mg or emtricitabine/tenofovir disoproxil were evaluated for detectable plasma tenofovir levels.

<sup>b</sup> Substudy participants received active adherence monitoring, e.g. unannounced home visits and pill counts, and counselling to improve compliance with study drug.

### Paediatric population

The safety and efficacy of emtricitabine/tenofovir disoproxil in children under the age of 12 years have not been established.

#### *Treatment of HIV-1 infection in the paediatric population*

There are no clinical studies conducted with emtricitabine/tenofovir disoproxil in the paediatric population with HIV-1 infection.

Clinical efficacy and safety of emtricitabine/tenofovir disoproxil was established from studies conducted with emtricitabine and tenofovir disoproxil when given as single agents.

#### *Studies with emtricitabine*

In infants and children older than 4 months, the majority of patients taking emtricitabine achieved or maintained complete suppression of plasma HIV-1 RNA through 48 weeks (89% achieved  $\leq 400$  copies/ml and 77% achieved  $\leq 50$  copies/ml).

#### *Studies with tenofovir disoproxil*

In study GS-US-104-0321, 87 HIV-1 infected treatment-experienced patients 12 to < 18 years of age were treated with tenofovir disoproxil (n = 45) or placebo (n = 42) in combination with an optimised background regimen (OBR) for 48 weeks. Due to limitations of the study, a benefit of tenofovir disoproxil over placebo was not demonstrated based on plasma HIV-1 RNA levels at week 24. However, a benefit is expected for the adolescent population based on extrapolation of adult data and comparative pharmacokinetic data (see section 5.2).

In patients who received treatment with tenofovir disoproxil or placebo, mean lumbar spine BMD Z-score was -1.004 and -0.809, and mean total body BMD Z-score was -0.866 and -0.584, respectively, at baseline. Mean changes at week 48 (end of double-blind phase) were -0.215 and -0.165 in lumbar spine BMD Z-score, and -0.254 and -0.179 in total body BMD Z-score for the tenofovir disoproxil and placebo groups, respectively. The mean rate of BMD gain was less in the tenofovir disoproxil group compared to the placebo group. At week 48, six adolescents in the tenofovir disoproxil group and one adolescent in the placebo group had significant lumbar spine BMD loss (defined as > 4% loss).

Among 28 patients receiving 96 weeks of treatment with tenofovir disoproxil, BMD Z-scores declined by -0.341 for lumbar spine and -0.458 for total body.

In study GS-US-104-0352, 97 treatment-experienced patients 2 to < 12 years of age with stable, virologic suppression on stavudine- or zidovudine-containing regimens were randomised to either replace stavudine or zidovudine with tenofovir disoproxil (n = 48) or continue on their original regimen (n = 49) for 48 weeks. At week 48, 83% of patients in the tenofovir disoproxil treatment group and 92% of patients in the stavudine or zidovudine treatment group had HIV-1 RNA concentrations < 400 copies/ml. The difference in the proportion of patients who maintained < 400 copies/ml at week 48 was mainly influenced by the higher number of discontinuations in the tenofovir disoproxil treatment group. When missing data were excluded, 91% of patients in the tenofovir disoproxil treatment group and 94% of patients in the stavudine or zidovudine treatment group had HIV-1 RNA concentrations < 400 copies/ml at week 48.

Reductions in BMD have been reported in paediatric patients. In patients who received treatment with tenofovir disoproxil, or stavudine or zidovudine, mean lumbar spine BMD Z-score was -1.034 and -0.498, and mean total body BMD Z-score was -0.471 and -0.386, respectively, at baseline. Mean changes at week 48 (end of randomised phase) were 0.032 and 0.087 in lumbar spine BMD Z-score, and -0.184 and -0.027 in total body BMD Z-score for the tenofovir disoproxil and stavudine or zidovudine groups, respectively. The mean rate of lumbar spine bone gain at week 48 was similar between the tenofovir disoproxil treatment group and the stavudine or zidovudine treatment group. Total body bone gain was less in the tenofovir disoproxil treatment group compared to the stavudine or zidovudine treatment group. One tenofovir disoproxil treated subject and no stavudine or zidovudine treated subjects experienced significant (> 4%) lumbar spine BMD loss at week 48. BMD Z-scores declined by -0.012 for lumbar spine and by -0.338 for total body in the 64 subjects who were treated with tenofovir disoproxil for 96 weeks. BMD Z-scores were not adjusted for height and weight.

In study GS-US-104-0352, 8 out of 89 paediatric patients (9.0%) exposed to tenofovir disoproxil discontinued study drug due to renal adverse events. Five subjects (5.6%) had laboratory findings clinically consistent with proximal renal tubulopathy, 4 of whom discontinued tenofovir disoproxil therapy (median tenofovir disoproxil exposure 331 weeks).

#### *Pre-exposure prophylaxis in the paediatric population*

The efficacy and safety of emtricitabine/tenofovir disoproxil for pre-exposure prophylaxis in adolescents who adhere to daily dosing is expected to be similar to that in adults at the same level of adherence. The potential renal and bone effects with long-term use of emtricitabine/tenofovir disoproxil for pre-exposure prophylaxis in adolescents are uncertain (see section 4.4).

## **5.2 Pharmacokinetic properties**

### Absorption

The bioequivalence of one emtricitabine/tenofovir disoproxil phosphate film-coated tablet with one emtricitabine 200 mg hard capsule and one tenofovir disoproxil 245 mg film-coated tablet was established following single dose administration to fasting healthy subjects. Following oral administration of emtricitabine/tenofovir disoproxil to healthy subjects, emtricitabine and tenofovir disoproxil are rapidly absorbed and tenofovir disoproxil is converted to tenofovir. Maximum emtricitabine and tenofovir concentrations are observed in serum within 0.5 to 3.0 h of dosing in the fasted state. Administration of emtricitabine/tenofovir disoproxil with food resulted in a delay of approximately three quarters of an hour in reaching maximum tenofovir concentrations and increases in tenofovir AUC and  $C_{max}$  of approximately 35% and 15%, respectively, when administered with a high fat or light meal, compared to administration in the fasted state. In order to optimise the absorption of tenofovir, it is recommended that emtricitabine/tenofovir disoproxil should preferably be taken with food.

## Distribution

Following intravenous administration the volume of distribution of emtricitabine and tenofovir was approximately 1.4 l/kg and 800 ml/kg, respectively. After oral administration of emtricitabine or tenofovir disoproxil, emtricitabine and tenofovir are widely distributed throughout the body. *In vitro* binding of emtricitabine to human plasma proteins was < 4% and independent of concentration over the range of 0.02 to 200 µg/ml. *In vitro* protein binding of tenofovir to plasma or serum protein was less than 0.7 and 7.2%, respectively, over the tenofovir concentration range 0.01 to 25 µg/ml.

## Biotransformation

There is limited metabolism of emtricitabine. The biotransformation of emtricitabine includes oxidation of the thiol moiety to form the 3'-sulphoxide diastereomers (approximately 9% of dose) and conjugation with glucuronic acid to form 2'-O-glucuronide (approximately 4% of dose). *In vitro* studies have determined that neither tenofovir disoproxil nor tenofovir are substrates for the CYP450 enzymes. Neither emtricitabine nor tenofovir inhibited *in vitro* drug metabolism mediated by any of the major human CYP450 isoforms involved in drug biotransformation. Also, emtricitabine did not inhibit uridine-5'-diphosphoglucuronyl transferase, the enzyme responsible for glucuronidation.

## Elimination

Emtricitabine is primarily excreted by the kidneys with complete recovery of the dose achieved in urine (approximately 86%) and faeces (approximately 14%).

13% of the emtricitabine dose was recovered in urine as three metabolites. The systemic clearance of emtricitabine averaged 307 ml/min. Following oral administration, the elimination half-life of emtricitabine is approximately 10 hours.

Tenofovir is primarily excreted by the kidney by both filtration and an active tubular transport system with approximately 70 – 80% of the dose excreted unchanged in urine following intravenous administration. The apparent clearance of tenofovir averaged approximately 307 ml/min. Renal clearance has been estimated to be approximately 210 ml/min, which is in excess of the glomerular filtration rate. This indicates that active tubular secretion is an important part of the elimination of tenofovir. Following oral administration, the elimination half-life of tenofovir is approximately 12 to 18 hours.

## Elderly

Pharmacokinetic studies have not been performed with emtricitabine or tenofovir (administered as tenofovir disoproxil) in the elderly (over 65 years of age).

## Gender

Emtricitabine and tenofovir pharmacokinetics are similar in male and female patients.

## Ethnicity

No clinically important pharmacokinetic difference due to ethnicity has been identified for emtricitabine. The pharmacokinetics of tenofovir (administered as tenofovir disoproxil) have not been specifically studied in different ethnic groups.

## Paediatric population

Pharmacokinetic studies have not been performed with emtricitabine/tenofovir disoproxil in children and adolescents (under 18 years of age). Steady-state pharmacokinetics of tenofovir were evaluated in 8 HIV-1 infected adolescent patients (aged 12 to < 18 years) with body weight ≥ 35 kg and in 23 HIV-1 infected children aged 2 to < 12 years. Tenofovir exposure achieved in these paediatric patients receiving oral daily doses of tenofovir disoproxil 245 mg or 6.5 mg/kg body weight tenofovir

disoproxil up to a maximum dose of 245 mg was similar to exposures achieved in adults receiving once-daily doses of tenofovir disoproxil 245 mg. Pharmacokinetic studies have not been performed with tenofovir disoproxil in children under 2 years.

In general, the pharmacokinetics of emtricitabine in infants, children and adolescents (aged 4 months up to 18 years) are similar to those seen in adults.

The pharmacokinetics of emtricitabine and tenofovir (administered as tenofovir disoproxil) are expected to be similar in HIV-1 infected and uninfected adolescents based on the similar exposures of emtricitabine and tenofovir in HIV-1 infected adolescents and adults, and the similar exposures of emtricitabine and tenofovir in HIV-1 infected and uninfected adults.

### Renal impairment

Limited pharmacokinetic data are available for emtricitabine and tenofovir after co-administration of separate preparations or as emtricitabine/tenofovir disoproxil in patients with renal impairment. Pharmacokinetic parameters were mainly determined following administration of single doses of emtricitabine 200 mg or tenofovir disoproxil 245 mg to non-HIV infected subjects with varying degrees of renal impairment. The degree of renal impairment was defined according to baseline creatinine clearance (CrCl) (normal renal function when CrCl > 80 ml/min; mild impairment with CrCl = 50 – 79 ml/min; moderate impairment with CrCl = 30 – 49 ml/min and severe impairment with CrCl = 10 – 29 ml/min).

The mean (%CV) emtricitabine drug exposure increased from 12 (25%) µg·h/ml in subjects with normal renal function, to 20 (6%) µg·h/ml, 25 (23%) µg·h/ml and 34 (6%) µg·h/ml, in subjects with mild, moderate and severe renal impairment, respectively.

The mean (%CV) tenofovir drug exposure increased from 2,185 (12%) ng·h/ml in subjects with normal renal function, to 3,064 (30%) ng·h/ml, 6,009 (42%) ng·h/ml and 15,985 (45%) ng·h/ml, in subjects with mild, moderate and severe renal impairment, respectively.

The increased dose interval for emtricitabine/tenofovir disoproxil in HIV-1 infected patients with moderate renal impairment is expected to result in higher peak plasma concentrations and lower C<sub>min</sub> levels as compared to patients with normal renal function.

In subjects with end-stage renal disease (ESRD) requiring haemodialysis, between dialysis drug exposures substantially increased over 72 hours to 53 (19%) µg·h/ml of emtricitabine, and over 48 hours to 42,857 (29%) ng·h/ml of tenofovir.

A small clinical study was conducted to evaluate the safety, antiviral activity and pharmacokinetics of tenofovir disoproxil in combination with emtricitabine in HIV infected patients with renal impairment. A subgroup of patients with baseline creatinine clearance between 50 and 60 ml/min, receiving once daily dosing, had a 2 – 4-fold increase in tenofovir exposure and worsening renal function.

The pharmacokinetics of emtricitabine and tenofovir (administered as tenofovir disoproxil) in paediatric patients with renal impairment have not been studied. No data are available to make dose recommendations (see sections 4.2 and 4.4).

### Hepatic impairment

The pharmacokinetics of emtricitabine / tenofovir disoproxil have not been studied in subjects with hepatic impairment.

The pharmacokinetics of emtricitabine have not been studied in non-HBV infected subjects with varying degrees of hepatic insufficiency. In general, emtricitabine pharmacokinetics in HBV infected subjects were similar to those in healthy subjects and in HIV infected patients.

A single 245 mg dose of tenofovir disoproxil was administered to non-HIV infected subjects with varying degrees of hepatic impairment defined according to Child-Pugh-Turcotte (CPT) classification. Tenofovir pharmacokinetics were not substantially altered in subjects with hepatic impairment suggesting that no dose adjustment is required in these subjects. The mean (%CV) tenofovir  $C_{max}$  and  $AUC_{0-\infty}$  values were 223 (34.8%) ng/ml and 2,050 (50.8%) ng·h/ml, respectively, in normal subjects compared with 289 (46.0%) ng/ml and 2,310 (43.5%) ng·h/ml in subjects with moderate hepatic impairment, and 305 (24.8%) ng/ml and 2,740 (44.0%) ng·h/ml in subjects with severe hepatic impairment.

### **5.3 Preclinical safety data**

#### Emtricitabine

Non-clinical data on emtricitabine reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity, carcinogenic potential and toxicity to reproduction and development.

#### Tenofovir disoproxil

Non-clinical safety pharmacology studies on tenofovir disoproxil reveal no special hazard for humans. Repeated dose toxicity studies in rats, dogs and monkeys at exposure levels greater than or equal to clinical exposure levels and with possible relevance to clinical use include renal and bone toxicity and a decrease in serum phosphate concentration. Bone toxicity was diagnosed as osteomalacia (monkeys) and reduced BMD (rats and dogs). The bone toxicity in young adult rats and dogs occurred at exposures  $\geq 5$ -fold the exposure in paediatric or adult patients; bone toxicity occurred in juvenile infected monkeys at very high exposures following subcutaneous dosing ( $\geq 40$ -fold the exposure in patients). Findings in the rat and monkey studies indicated that there was a substance-related decrease in intestinal absorption of phosphate with potential secondary reduction in BMD.

Genotoxicity studies revealed positive results in the *in vitro* mouse lymphoma assay, equivocal results in one of the strains used in the Ames test, and weakly positive results in an UDS test in primary rat hepatocytes. However, it was negative in an *in vivo* mouse bone marrow micronucleus assay.

Oral carcinogenicity studies in rats and mice only revealed a low incidence of duodenal tumours at an extremely high dose in mice. These tumours are unlikely to be of relevance to humans.

Reproductive toxicity studies in rats and rabbits showed no effects on mating, fertility, pregnancy or foetal parameters. However, tenofovir disoproxil reduced the viability index and weight of pups in a peri and postnatal toxicity study at maternally toxic doses.

#### Combination of emtricitabine and tenofovir disoproxil

Genotoxicity and repeated dose toxicity studies of one month or less with the combination of these two components found no exacerbation of toxicological effects compared to studies with the separate components.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core

Microcrystalline cellulose  
Croscarmellose sodium  
Talc  
Hydrophobic colloidal silica  
Magnesium stearate

#### Film-coating

Polyvinyl alcohol  
Titanium dioxide  
Macrogol  
Talc  
Indigo carmine aluminum lake (E 132)

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

2 years  
Store below 30°C after first opening.

### **6.4 Special precautions for storage**

This medicinal product does not require any special temperature storage conditions. Store in the original package in order to protect from moisture.  
For storage conditions after first opening of the medicinal product, see section 6.3.

### **6.5 Nature and contents of container**

High density polyethylene (HDPE) bottle with a polypropylene child-resistant cap and induction heat sealing (with aluminium foil) containing 30 film-coated tablets and a silica gel desiccant (in a HDPE container).

The following pack sizes are available:

Outer cartons containing 1 bottle of 30 film-coated tablets and outer cartons containing 90 (3 bottles of 30) film-coated tablets.

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal and other handling**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Open bottles by pressing the child proof cap down and turning it counter-clock wise.

**7. MARKETING AUTHORISATION HOLDER**

Zentiva, k.s.  
U Kabelovny 130  
102 37 Prague 10  
Czech Republic

**8. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1148/001  
EU/1/16/1148/002

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

Date of first authorisation: 9 November 2016  
Date of latest renewal: 16 July 2021

**10. DATE OF REVISION OF THE TEXT**

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu>.



## **ANNEX II**

- A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**
- B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**
- C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**
- D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

## **A. MANUFACTURER(S) RESPONSIBLE FOR BATCH RELEASE**

Name and address of the manufacturer(s) responsible for batch release

S.C. Zentiva S.A.  
50 Theodor Pallady Blvd.  
Bucharest 032266  
Romania

## **B. CONDITIONS OR RESTRICTIONS REGARDING SUPPLY AND USE**

Medicinal product subject to restricted medical prescription (see Annex I: Summary of Product Characteristics, section 4.2).

## **C. OTHER CONDITIONS AND REQUIREMENTS OF THE MARKETING AUTHORISATION**

- **Periodic safety update reports (PSURs)**

The requirements for submission of PSURs for this medicinal product are set out in the list of Union reference dates (EURD list) provided for under Article 107c(7) of Directive 2001/83/EC and any subsequent updates published on the European medicines web-portal.

## **D. CONDITIONS OR RESTRICTIONS WITH REGARD TO THE SAFE AND EFFECTIVE USE OF THE MEDICINAL PRODUCT**

- **Risk management plan (RMP)**

The marketing authorisation holder (MAH) shall perform the required pharmacovigilance activities and interventions detailed in the agreed RMP presented in Module 1.8.2 of the marketing authorisation and any agreed subsequent updates of the RMP.

An updated RMP should be submitted:

- At the request of the European Medicines Agency;
- Whenever the risk management system is modified, especially as the result of new information being received that may lead to a significant change to the benefit/risk profile or as the result of an important (pharmacovigilance or risk minimisation) milestone being reached.

- **Additional risk minimisation measures**

The marketing authorisation holder (MAH) shall ensure that all physicians who are expected to prescribe/use Emtricitabine/Tenofovir disoproxil Zentiva in adults and adolescents for PrEP are provided with a physician educational pack containing the Summary of Product Characteristics and an appropriate educational brochure, as detailed below:

- PrEP educational brochure for prescribers entitled 'Important Safety Information for Prescribers About Emtricitabine/Tenofovir disoproxil Zentiva for a Pre-exposure Prophylaxis (PrEP) Indication'
- PrEP Checklist for prescribers
- PrEP educational brochure for the individual at risk entitled 'Important Information About Emtricitabine/Tenofovir disoproxil Zentiva to Reduce the Risk of getting Human Immunodeficiency Virus (HIV) Infection'
- PrEP reminder card

**PrEP educational brochure for prescribers:**

- Reminder of the key safety information regarding the use of Emtricitabine/Tenofovir disoproxil Zentiva for PrEP in adults and adolescents
- Reminder of factors to help identify individuals at high risk of acquiring HIV-1
- Reminder on the risk of development of HIV-1 drug resistance in undiagnosed HIV-1–Infected individuals
- Provides safety information on adherence, HIV testing, renal, bone and HBV status.

**PrEP Checklist for prescribers:**

- Reminders for evaluations/counselling at the initial visit and follow-up.

**PrEP educational brochure for the individual at risk (to be provided by healthcare provider [HCP]):**

- Reminders on what the individual should know before and while taking Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV infection
- Reminder on the importance of strict adherence to the recommended dosing regimen
- Provides information on how to take Emtricitabine/Tenofovir disoproxil Zentiva
- Provides information on the possible side effects
- Provides information on how to store Emtricitabine/Tenofovir disoproxil Zentiva.

**PrEP reminder card for the individual at risk (to be provided by HCP):**

- Reminders to adhere to the dosing schedule
- Reminder to attend scheduled clinic visits.

**ANNEX III**  
**LABELLING AND PACKAGE LEAFLET**

## **A. LABELLING**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING AND THE IMMEDIATE PACKAGING**

**Outer carton / bottle label**

**1. NAME OF THE MEDICINAL PRODUCT**

Emtricitabine/Tenofovir disoproxil Zentiva 200 mg/245 mg film-coated tablets  
emtricitabine/tenofovir disoproxil

**2. STATEMENT OF ACTIVE SUBSTANCES**

Each film-coated tablet contains 200 mg of emtricitabine and 245 mg of tenofovir disoproxil (equivalent to 291.5 mg of tenofovir disoproxil phosphate or 136 mg of tenofovir).

**3. LIST OF EXCIPIENTS**

**4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

30 film-coated tablets

90 (3 bottles of 30) film-coated tablets

**5. METHOD AND ROUTE OF ADMINISTRATION**

Oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

Store in the original package in order to protect from moisture.  
Store below 30 °C after first opening.

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

Zentiva, k.s.  
U Kabelovny 130  
102 37 Prague 10  
Czech Republic

**12. MARKETING AUTHORISATION NUMBER(S)**

EU/1/16/1148/001 30 film-coated tablets  
EU/1/16/1148/002 90 (3x30) film-coated tablets

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

Emtricitabine/Tenofovir disoproxil Zentiva [outer packaging only]

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included. [outer packaging only]

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC [outer packaging only]  
SN [outer packaging only]  
NN [outer packaging only]

## **B. PACKAGE LEAFLET**



## Package leaflet: Information for the user

### Emtricitabine/Tenofovir disoproxil Zentiva 200 mg/245 mg film-coated tablets emtricitabine/tenofovir disoproxil

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet**

1. What Emtricitabine/Tenofovir disoproxil Zentiva is and what it is used for
2. What you need to know before you take Emtricitabine/Tenofovir disoproxil Zentiva
3. How to take Emtricitabine/Tenofovir disoproxil Zentiva
4. Possible side effects
5. How to store Emtricitabine/Tenofovir disoproxil Zentiva
6. Contents of the pack and other information

#### **1. What Emtricitabine/Tenofovir disoproxil Zentiva is and what it is used for**

**Emtricitabine/Tenofovir disoproxil Zentiva contains two active substances, *emtricitabine* and *tenofovir disoproxil*.** Both of these active substances are *antiretroviral* medicines which are used to treat HIV infection. Emtricitabine is a *nucleoside reverse transcriptase inhibitor* and tenofovir is a *nucleotide reverse transcriptase inhibitor*. However, both are generally known as NRTIs and they work by interfering with the normal working of an enzyme (reverse transcriptase) that is essential for the virus to reproduce itself.

- **Emtricitabine/Tenofovir disoproxil Zentiva is used to treat Human Immunodeficiency Virus 1(HIV-1) infection, in adults.**
- **It is also used to treat HIV in adolescents aged 12 to less than 18 years who weigh at least 35 kg,** and who have already been treated with other HIV medicines that are no longer effective or have caused side effects.
  - Emtricitabine/Tenofovir disoproxil Zentiva should always be used combined with other medicines to treat HIV infection.
  - Emtricitabine/Tenofovir disoproxil Zentiva can be administered in place of emtricitabine and tenofovir disoproxil used separately at the same doses.

**This medicine is not a cure for HIV infection.** While taking Emtricitabine/Tenofovir disoproxil Zentiva you may still develop infections or other illnesses associated with HIV infection.

- **Emtricitabine/Tenofovir disoproxil Zentiva is also used to reduce the risk of getting HIV-1 infection in adults, and adolescents aged 12 years to less than 18 years who weigh at least 35 kg,** when taken daily, together with safer sex practices:  
See section 2 for a list of precautions to take against HIV infection.

## 2. What you need to know before you take Emtricitabine/Tenofovir disoproxil Zentiva

**Do not take Emtricitabine/Tenofovir disoproxil Zentiva to treat HIV or to reduce the risk of getting HIV if you are allergic** to emtricitabine, tenofovir, tenofovir disoproxil or any of the other ingredients of this medicine (listed in section 6).

→ **If this applies to you, tell your doctor immediately.**

### **Before taking Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV:**

Emtricitabine/Tenofovir disoproxil Zentiva can only help reduce your risk of getting HIV **before** you are infected.

- **You must be HIV negative before you start to take Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV.** You must get tested to make sure that you do not already have HIV infection. Do not take Emtricitabine/Tenofovir disoproxil Zentiva to reduce your risk unless you are confirmed to be HIV negative. People who do have HIV must take Emtricitabine/Tenofovir disoproxil Zentiva in combination with other medicines.
- **Many HIV tests can miss a recent infection.** If you get a flu-like illness, it could mean you have recently been infected with HIV. These may be signs of HIV infection:
  - tiredness
  - fever
  - joint or muscle aches
  - headache
  - vomiting or diarrhoea
  - rash
  - night sweats
  - enlarged lymph nodes in the neck or groin

**Tell your doctor about any flu-like illness** – either in the month before starting Emtricitabine/Tenofovir disoproxil Zentiva, or at any time while taking Emtricitabine/Tenofovir disoproxil Zentiva.

### **Warnings and precautions**

#### **While taking Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV:**

- Take Emtricitabine/Tenofovir disoproxil Zentiva every day **to reduce your risk, not just when you think you have been at risk of HIV infection.** Do not miss any doses of Emtricitabine/Tenofovir disoproxil Zentiva, or stop taking it. Missing doses may increase your risk of getting HIV infection.
- Get tested for HIV regularly.
- If you think you were infected with HIV, tell your doctor straight away. They may want to do more tests to make sure you are still HIV negative.
- **Just taking Emtricitabine/Tenofovir disoproxil Zentiva may not stop you getting HIV.**
  - Always practice safer sex. Use condoms to reduce contact with semen, vaginal fluids, or blood.
  - Do not share personal items that can have blood or body fluids on them, such as toothbrushes and razor blades.
  - Do not share or re-use needles or other injection or drug equipment.
  - Get tested for other sexually transmitted infections such as syphilis and gonorrhoea. These infections make it easier for HIV to infect you.

Ask your doctor if you have any more questions about how to prevent getting HIV or spreading HIV to other people.

**While taking Emtricitabine/Tenofovir disoproxil Zentiva to treat HIV or to reduce the risk of getting HIV:**

- **Emtricitabine/Tenofovir disoproxil Zentiva may affect your kidneys.** Before and during treatment, your doctor may order blood tests to measure kidney function. Tell your doctor if you have had kidney disease, or if tests have shown kidney problems. Emtricitabine/Tenofovir disoproxil Zentiva should not be given to adolescents with existing kidney problems. If you have kidney problems, your doctor may advise you to stop taking Emtricitabine/Tenofovir disoproxil Zentiva or, if you already have HIV, to take Emtricitabine/Tenofovir disoproxil Zentiva less frequently. Emtricitabine/Tenofovir disoproxil Zentiva is not recommended if you have severe kidney disease or are on dialysis.
- **Talk to your doctor if you suffer from osteoporosis, have a history of bone fracture or if you have problems with your bones.**

**Bone problems** (manifesting as persistent or worsening bone pain and sometimes resulting in fractures) may also occur due to damage to kidney tubule cells (see section 4, Possible side effects). Tell your doctor if you have bone pain or fractures.

Tenofovir disoproxil may also cause loss of bone mass. The most pronounced bone loss was seen in clinical studies when patients were treated for HIV with tenofovir disoproxil in combination with a boosted protease inhibitor.

Overall, the effects of tenofovir disoproxil on long-term bone health and future fracture risk in adult and paediatric patients are uncertain.

- **Talk to your doctor if you have a history of liver disease, including hepatitis.** Patients infected with HIV who also have liver disease (including chronic hepatitis B or C), who are treated with antiretrovirals, have a higher risk of severe and potentially fatal liver complications. If you have hepatitis B or C, your doctor will carefully consider the best treatment regimen for you.
- **Know your hepatitis B virus (HBV) infection status** before starting Emtricitabine/Tenofovir disoproxil Zentiva. If you have HBV, there is a serious risk of liver problems when you stop taking Emtricitabine/Tenofovir disoproxil Zentiva, whether or not you also have HIV. It is important not to stop taking Emtricitabine/Tenofovir disoproxil Zentiva without talking to your doctor: see section 3, Do not stop taking Emtricitabine/Tenofovir disoproxil Zentiva.
- **Talk to your doctor if you are over 65.** The combination of emtricitabine and tenofovir disoproxil has not been studied in patients over 65 years of age.

**Children and adolescents**

Emtricitabine/Tenofovir disoproxil Zentiva is not for use in children under 12 years of age.

**Other medicines and Emtricitabine/Tenofovir disoproxil Zentiva**

**Do not take Emtricitabine/Tenofovir disoproxil Zentiva** if you are already taking other medicines that contain the components of Emtricitabine/Tenofovir disoproxil Zentiva (emtricitabine and tenofovir disoproxil) or any other antiviral medicines that contain tenofovir alafenamide, lamivudine or adefovir dipivoxil.

**Taking Emtricitabine/Tenofovir disoproxil Zentiva with other medicines that can damage your kidneys:** it is especially important to tell your doctor if you are taking any of these medicines, including

- aminoglycosides (for bacterial infection).
- amphotericin B (for fungal infection).
- foscarnet (for viral infection).

- ganciclovir (for viral infection).
- pentamidine (for infections).
- vancomycin (for bacterial infection).
- interleukin-2 (to treat cancer).
- cidofovir (for viral infection).
- non-steroidal anti-inflammatory drugs (NSAIDs, to relieve bone or muscle pains).

If you are taking another antiviral medicine called a protease inhibitor to treat HIV, your doctor may order blood tests to closely monitor your kidney function.

**It is also important to tell your doctor** if you are taking ledipasvir/sofosbuvir, sofosbuvir/velpatasvir or sofosbuvir/velpatasvir/voxilaprevir to treat hepatitis C infection.

**Taking Emtricitabine/Tenofovir disoproxil Zentiva with other medicines containing didanosine (for treatment of HIV infection):** Taking Emtricitabine/Tenofovir disoproxil Zentiva with other antiviral medicines that contain didanosine can raise the levels of didanosine in your blood and may reduce CD4 cell counts. Rarely, inflammation of the pancreas and lactic acidosis (excess lactic acid in the blood), which sometimes causes death, have been reported when medicines containing tenofovir disoproxil and didanosine were taken together. Your doctor will carefully consider whether to treat you with combinations of tenofovir and didanosine.

**Tell your doctor** if you are taking any of these medicines. Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

#### **Emtricitabine/Tenofovir disoproxil Zentiva with food and drink**

Whenever possible, Emtricitabine/Tenofovir disoproxil Zentiva should be taken with food.

#### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

If you have taken Emtricitabine/Tenofovir disoproxil Zentiva during your pregnancy, your doctor may request regular blood tests and other diagnostic tests to monitor the development of your child.

In children whose mothers took NRTIs during pregnancy, the benefit from the protection against HIV outweighed the risk of side effects.

- **Do not breast-feed during treatment with Emtricitabine/Tenofovir disoproxil Zentiva.** This is because the active substances in this medicine pass into human breast milk.
- Breast-feeding is not recommended in women living with HIV because HIV infection can be passed on to the baby in breast milk.
- If you are breast-feeding, or thinking about breast-feeding, you should **discuss it with your doctor as soon as possible**.

#### **Driving and using machines**

Emtricitabine/Tenofovir disoproxil Zentiva can cause dizziness. If you feel dizzy while taking Emtricitabine/Tenofovir disoproxil Zentiva, **do not drive** and do not use any tools or machines.

#### **Emtricitabine/Tenofovir disoproxil Zentiva contains sodium**

This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

### **3. How to take Emtricitabine/Tenofovir disoproxil Zentiva**

- **Always take this medicine exactly as your doctor has told you.** Check with your doctor or pharmacist if you are not sure.

**The recommended dose of Emtricitabine/Tenofovir disoproxil Zentiva to treat HIV is:**

- **Adults:** one tablet each day, where possible, with food.
- **Adolescents aged 12 to less than 18 years who weigh at least 35 kg:** one tablet each day, whenever possible with food.

**The recommended dose of Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV is:**

- **Adults:** one tablet each day, whenever possible, with food.
- **Adolescents aged 12 to less than 18 years who weigh at least 35 kg:** one tablet each day, whenever possible with food.

If you have difficulty swallowing, you can use the tip of a spoon to crush the tablet. Then mix the powder with about 100 ml (half a glass) of water, orange juice or grape juice, and drink immediately.

- **Always take the dose recommended by your doctor.** This is to make sure that your medicine is fully effective, and to reduce the risk of developing resistance to the treatment. Do not change the dose unless your doctor tells you to.
- If you are being treated for HIV infection, your doctor will prescribe Emtricitabine/Tenofovir disoproxil Zentiva with other antiretroviral medicines. Please refer to the patient information leaflets of the other antiretrovirals for guidance on how to take those medicines.
- If you are taking Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV, take Emtricitabine/Tenofovir disoproxil Zentiva every day, not just when you think you have been at risk of HIV infection.

Ask your doctor if you have any questions about how to prevent getting HIV or prevent spreading HIV to other people.

**If you take more Emtricitabine/Tenofovir disoproxil Zentiva than you should**

If you accidentally take more than the recommended dose of Emtricitabine/Tenofovir disoproxil Zentiva, contact your doctor or nearest emergency department for advice. Keep the tablet bottle with you so that you can easily describe what you have taken.

**If you forget to take Emtricitabine /Tenofovir disoproxil Zentiva**

It is important not to miss a dose of Emtricitabine/Tenofovir disoproxil Zentiva.

- **If you notice within 12 hours** of the time you usually take Emtricitabine/Tenofovir disoproxil Zentiva, take the tablet preferably with food as soon as possible. Then take the next dose at your usual time.
- **If you notice 12 hours or more after** the time you usually take Emtricitabine/Tenofovir disoproxil Zentiva, forget about the missed dose. Wait and take the next dose, preferably with food, at your usual time.

**If you vomit less than 1 hour after taking Emtricitabine/Tenofovir disoproxil Zentiva,** take another tablet. You do not need to take another tablet if you were sick more than 1 hour after taking Emtricitabine/Tenofovir disoproxil Zentiva.

**Do not stop taking Emtricitabine/Tenofovir disoproxil Zentiva**

- If you take Emtricitabine/Tenofovir disoproxil Zentiva for treatment of HIV infection, stopping tablets may reduce the effectiveness of the anti-HIV therapy recommended by your doctor.
- If you are taking Emtricitabine/Tenofovir disoproxil Zentiva to reduce the risk of getting HIV, do not stop taking Emtricitabine/Tenofovir disoproxil Zentiva or miss any doses. Stopping use of Emtricitabine/Tenofovir disoproxil Zentiva, or missing doses, may increase your risk of getting HIV infection.

**Do not stop taking Emtricitabine/Tenofovir disoproxil Zentiva without contacting your doctor.**

- **If you have hepatitis B**, it is especially important not to stop your Emtricitabine/Tenofovir disoproxil Zentiva treatment without talking to your doctor first. You may require blood tests for several months after stopping treatment. In some patients with advanced liver disease or cirrhosis, stopping treatment is not recommended as this may lead to worsening of your hepatitis, which may be life-threatening.

Tell your doctor immediately about new or unusual symptoms after you stop treatment, particularly symptoms you associate with hepatitis B infection.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

#### **4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

##### **Possible serious side effects:**

- **Lactic acidosis** (excess lactic acid in the blood) is a rare but potentially life-threatening side effect. Lactic acidosis occurs more often in women, particularly if they are overweight, and in people with liver disease. The following may be signs of lactic acidosis:
  - deep rapid breathing,
  - drowsiness,
  - feeling sick (nausea), being sick (vomiting),
  - stomach pain.

**If you think you may have lactic acidosis, get medical help immediately.**

- **Any signs of inflammation or infection.** In some patients with advanced HIV infection (AIDS) and a history of opportunistic infections (infections that occur in people with a weak immune system), signs and symptoms of inflammation from previous infections may occur soon after anti-HIV treatment is started. It is thought that these symptoms are due to an improvement in the body's immune response, enabling the body to fight infections that may have been present with no obvious symptoms.
- Autoimmune disorders, when the immune system attacks healthy body tissue, may also occur after you start taking medicines to treat HIV infection. Autoimmune disorders may occur many months after the start of treatment. Look out for any symptoms of infection or other symptoms such as:
  - muscle weakness,
  - weakness beginning in the hands and feet and moving up towards the trunk of the body,
  - palpitations, tremor or hyperactivity.

If you notice these or any symptoms of inflammation or infection, get medical help immediately.

##### **Possible side effects:**

**Very common side effects** (may affect more than 1 in 10 people):

- diarrhoea, being sick (vomiting), feeling sick (nausea),
- dizziness, headache,
- rash,
- feeling weak.

Tests may also show:

- decreases in phosphate in the blood,
- increased creatine kinase.

**Common side effects** (may affect up to 1 in 10 people):

- pain, stomach pain,
- difficulty sleeping, abnormal dreams,
- problems with digestion resulting in discomfort after meals, feeling bloated, flatulence,
- rashes (including red spots or blotches sometimes with blistering and swelling of the skin), which may be allergic reactions, itching, changes in skin colour including darkening of the skin in patches,
- other allergic reactions, such as wheezing, swelling or feeling light-headed,
- loss of bone mass

Tests may also show:

- low white blood cell count (a reduced white blood cell count can make you more prone to infection),
- increased triglycerides (fatty acids), bile or sugar in the blood,
- liver and pancreas problems.

**Uncommon side effects** (may affect up to 1 in 100 people):

- pain in the abdomen (tummy) caused by inflammation of the pancreas,
- swelling of the face, lips, tongue or throat,
- anaemia (low red blood cell count),
- breakdown of muscle, muscle pain or weakness which may occur due to damage to the kidney tubule cells.

Tests may also show:

- decreases in potassium in the blood,
- increased creatinine in your blood,
- changes to your urine.

**Rare side effects** (may affect up to 1 in 1,000 people):

- Lactic acidosis (see Possible serious side effects),
- fatty liver,
- yellow skin or eyes, itching, or pain in the abdomen (tummy) caused by inflammation of the liver,
- inflammation of the kidney, passing a lot of urine and feeling thirsty, kidney failure, damage to kidney tubule cells,
- softening of the bones (with bone pain and sometimes resulting in fractures),
- back pain caused by kidney problems.

Damage to kidney tubule cells may be associated with breakdown of muscle, softening of the bones (with bone pain and sometimes resulting in fractures), muscle pain, muscle weakness and decreases in potassium or phosphate in the blood.

**If you notice any of the side effects listed above or if any of the side effects get serious, talk to your doctor or pharmacist.**

The frequency of the following side effects is not known.

- Bone problems. Some patients taking combination antiretroviral medicines such as Emtricitabine /Tenofovir disoproxil Zentiva may develop a bone disease called osteonecrosis (death of bone tissue caused by loss of blood supply to the bone). Taking this type of medicine for a long time, taking corticosteroids, drinking alcohol, having a very weak immune system, and being overweight, may be some of the many risk factors for developing this disease. Signs of osteonecrosis are:
  - joint stiffness,
  - joint aches and pains (especially of the hip, knee and shoulder),
  - difficulty with movement.

**If you notice any of these symptoms tell your doctor.**

During treatment for HIV there may be an increase in weight and in levels of blood lipids and glucose. This is partly linked to restored health and life style, and in the case of blood lipids sometimes to the HIV medicines themselves. Your doctor will test for these changes.

#### **Other effects in children**

- Children given emtricitabine very commonly experienced changes in skin colour including
  - darkening of the skin in patches
- Children commonly experienced low red blood cell count (anaemia).
  - this may cause the child to be tired or breathless

**If you notice any of these symptoms tell your doctor.**

#### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system](#) listed in [Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

### **5. How to store Emtricitabine/Tenofovir disoproxil Zentiva**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and on the bottle after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special temperature storage conditions. Store in the original package in order to protect from moisture.  
Store below 30°C after first opening.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

### **6. Contents of the pack and other information**

#### **What Emtricitabine/Tenofovir disoproxil Zentiva contains**

- **The active substances are** emtricitabine and tenofovir disoproxil phosphate. Each film-coated tablet contains 200 mg of emtricitabine and 245 mg of tenofovir disoproxil (equivalent to 291.5 mg of tenofovir disoproxil phosphate or 136 mg of tenofovir).
- **The other ingredients are:**  
Tablet core: microcrystalline cellulose, croscarmellose sodium, talc, hydrophobic colloidal silica, magnesium stearate.  
Film coating: polyvinyl alcohol, titanium dioxide, macrogol, talc, indigo carmine aluminum lake (E 132).

#### **What Emtricitabine/Tenofovir disoproxil Zentiva looks like and contents of the pack**

Emtricitabine /Tenofovir disoproxil Zentiva 200 mg/245 mg film-coated tablets are blue, oval biconvex film coated tablet without half-score with dimensions approx. 19.35 x 9.75 mm.

Contents of the pack:

Outer cartons containing 1 bottle of 30 film-coated tablets and outer cartons containing 3 bottles of 30 film-coated tablets.

Not all pack sizes may be marketed.



Open bottles by pressing the child proof cap down and turning it counter-clock wise.

Each bottle contains a silica gel desiccant that must be kept in the bottle to help protect your tablets. The silica gel desiccant is contained in a separate container and should not be swallowed.

### **Marketing Authorisation Holder**

Zentiva, k.s.  
U Kabelovny 130  
102 37 Prague 10  
Czech Republic

### **Manufacturer**

S.C. Zentiva S.A.  
50 Theodor Pallady Blvd.  
Bucharest 032266  
Romania

For any information about this medicine, please contact the local representative of the Marketing Authorisation Holder:

#### **België/Belgique/Belgien**

Zentiva, k.s.  
Tél/Tel: +32 (78) 700 112  
PV-Belgium@zentiva.com

#### **България**

Zentiva, k.s.  
Тел: +35924417136  
PV-Bulgaria@zentiva.com

#### **Česká republika**

Zentiva, k.s.  
Tel: +420 267 241 111  
PV-Czech-Republic@zentiva.com

#### **Danmark**

Zentiva Denmark ApS  
Tlf.: +45 787 68 400  
PV-Denmark@zentiva.com

#### **Deutschland**

Zentiva Pharma GmbH  
Tel: +49 (0) 800 53 53 010  
PV-Germany@zentiva.com

#### **Eesti**

Zentiva, k.s.  
Tel: +372 52 70308  
PV-Estonia@zentiva.com

#### **Ελλάδα**

Zentiva, k.s.  
Τηλ: +30 211 198 7510  
PV-Greece@zentiva.com

#### **España**

Zentiva Spain S.L.U.  
Tel: +34 91 111 58 93  
PV-Spain@zentiva.com

#### **Lietuva**

Zentiva, k.s.  
Tel: +370 52152025  
PV-Lithuania@zentiva.com

#### **Luxembourg/Luxemburg**

Zentiva, k.s.  
Tél/Tel: +352 208 82330  
PV-Luxembourg@zentiva.com

#### **Magyarország**

Zentiva Pharma Kft.  
Tel.: +36 1 299 1058  
PV-Hungary@zentiva.com

#### **Malta**

Zentiva, k.s.  
Tel: +356 2034 1796  
PV-Malta@zentiva.com

#### **Nederland**

Zentiva, k.s.  
Tel: +31 202 253 638  
PV-Netherlands@zentiva.com

#### **Norge**

Zentiva Denmark ApS  
Tlf: +45 787 68 400  
PV-Norway@zentiva.com

#### **Österreich**

Zentiva, k.s.  
Tel: +43 720 778 877  
PV-Austria@zentiva.com

#### **Polska**

Zentiva Polska Sp. z o.o.  
Tel: + 48 22 375 92 00  
PV-Poland@zentiva.com

**France**

Zentiva France  
Tél: +33 (0) 800 089 219  
PV-France@zentiva.com

**Hrvatska**

Zentiva d.o.o.  
Tel: +385 1 6641 830  
PV-Croatia@zentiva.com

**Ireland**

Zentiva, k.s.  
Tel: +353 818 882 243  
PV-Ireland@zentiva.com

**Ísland**

Zentiva Denmark ApS  
Sími: +354 539 5025  
PV-Iceland@zentiva.com

**Italia**

Zentiva Italia S.r.l.  
Tel: +39 800081631  
PV-Italy@zentiva.com

**Κύπρος**

Zentiva, k.s.  
Τηλ: +30 211 198 7510  
PV-Cyprus@zentiva.com

**Latvija**

Zentiva, k.s.  
Tel: +371 67893939  
PV-Latvia@zentiva.com

**Portugal**

Zentiva Portugal, Lda  
Tel: +351210601360  
PV-Portugal@zentiva.com

**România**

ZENTIVA S.A.  
Tel: +4 021.304.7597  
PV-Romania@zentiva.com

**Slovenija**

Zentiva, k.s.  
Tel: +386 360 00 408  
PV-Slovenia@zentiva.com

**Slovenská republika**

Zentiva, a.s.  
Tel: +421 2 3918 3010  
PV-Slovakia@zentiva.com

**Suomi/Finland**

Zentiva Denmark ApS  
Puh/Tel: +358 942 598 648  
PV-Finland@zentiva.com

**Sverige**

Zentiva Denmark ApS  
Tel: +46 840 838 822  
PV-Sweden@zentiva.com

**United Kingdom (Northern Ireland)**

Zentiva, k.s.  
Tel: +44 (0) 800 090 2408  
PV-United-Kingdom@zentiva.com

**This leaflet was last revised in**

Detailed information on this medicine is available on the European Medicines Agency web site:  
<https://www.ema.europa.eu>.