

Introduction to Machine Learning for Social Science

Class 3: Multivariate Regression

Rochelle Terman

Postdoctoral Fellow
Center for International Security Cooperation
Stanford University

January 16th, 2018

Loose Threads

1) Reducible vs. Irreducible error.

- Even if you estimate f perfectly, don't expect perfect predictions.

2) Least squared versus absolute errors.

- Some cases do apply least absolute deviations (L1 regression), which is robust to outliers.
- Mathematical convenience: Easier to compute the derivative of a polynomial than absolute value.
- By Gauss Markov, OLS is BLUE

Questions?





Predicting Election Results

Goal: Predict **Incumbent Vote Share** (create prediction function)

- Use relationship in prior elections to predict future election
- Training data (In sample) \rightsquigarrow Testing data (Out of sample)

Method: Linear Regression (Least squares)

Evaluation (Focus of next lectures):

- 1) In sample fit (training data)
- 2) Out of sample fit (test data)

Time for Change Model (Abramowitz, Linzer)

Predict **Incumbent Vote Share** with political and economic fundamentals

- 1) GDP Growth
- 2) Incumbent Presidential Popularity
- 3) Incumbent Party

Our Estimated Prediction Function

Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \epsilon_i$$

Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \epsilon_i$$


$$\text{Vote}_i = \underbrace{50.76 + 0.16 \times \text{Approval}_i}_{\widehat{\text{Vote}_i}} + \epsilon_i$$





Our Estimated Prediction Function

$$\text{Vote}_i = \hat{\beta}_0 + \hat{\beta}_1 \text{Approval}_i + \epsilon_i$$

$$\text{Vote}_i = \underbrace{50.76 + 0.16 \times \text{Approval}_i}_{\widehat{\text{Vote}_i}} + \epsilon_i$$

R Code!





What is Linear Regression?

Multiple linear regression:

- Assumes a linear relationship between quantitative response Y and **multiple** variables X_1, X_2, \dots, X_p .
- Also called **multivariate regression** because there are > 2 variables.

Multivariate Regression: Function Perspective

Function f such that

$$\text{Vote}_i = f(\text{Approval}_i, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Inc 2nd Term}_i) + \epsilon_i$$

Multivariate Regression: Function Perspective

Function f such that

$$\begin{aligned}\text{Vote}_i &= f(\text{Approval}_i, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Inc 2nd Term}_i) + \epsilon_i \\ &= \beta_0 + \beta_1 \text{Approval}_i + \beta_2 \text{Q1 GDP}_i \\ &\quad + \beta_3 \text{Q2 GDP}_i + \beta_4 \text{Inc 2nd Term}_i + \epsilon_i\end{aligned}$$

Multivariate Regression: Function Perspective

Function f such that

$$\text{Vote}_i = f(\text{Approval}_i, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Inc 2nd Term}_i) + \epsilon_i$$

$$\begin{aligned} &= \beta_0 + \beta_1 \text{Approval}_i + \beta_2 \text{Q1 GDP}_i \\ &\quad + \beta_3 \text{Q2 GDP}_i + \beta_4 \text{Inc 2nd Term}_i + \epsilon_i \end{aligned}$$

$$\begin{aligned} \widehat{\text{Vote}_i} &= \beta_0 + \beta_1 \text{Approval}_i + \beta_2 \text{Q1 GDP}_i \\ &\quad + \beta_3 \text{Q2 GDP}_i + \beta_4 \text{Inc 2nd Term}_i \end{aligned}$$

Multivariate Regression: Function Perspective

Function f such that

$$\text{Vote}_i = f(\text{Approval}_i, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Inc 2nd Term}_i) + \epsilon_i$$

$$\begin{aligned} &= \beta_0 + \beta_1 \text{Approval}_i + \beta_2 \text{Q1 GDP}_i \\ &\quad + \beta_3 \text{Q2 GDP}_i + \beta_4 \text{Inc 2nd Term}_i + \epsilon_i \end{aligned}$$

$$\begin{aligned} \widehat{\text{Vote}_i} &= \beta_0 + \beta_1 \text{Approval}_i + \beta_2 \text{Q1 GDP}_i \\ &\quad + \beta_3 \text{Q2 GDP}_i + \beta_4 \text{Inc 2nd Term}_i \end{aligned}$$

We interpret β_j as the average effect on Y of a one unit increase in X_j , holding all other predictors fixed.

Fitting a Multivariate Regression

Fitting a Multivariate Regression

Choose β 's to minimize:

Fitting a Multivariate Regression

Choose β 's to minimize:

$$\sum_{i=1948}^{2012} \epsilon_i^2 = \sum_{i=1948}^{2012} \left(\text{Vote}_i - \widehat{\text{Vote}}_i \right)^2$$

Fitting a Multivariate Regression

Choose β 's to minimize:

$$\begin{aligned}\sum_{i=1948}^{2012} \epsilon_i^2 &= \sum_{i=1948}^{2012} \left(\text{Vote}_i - \widehat{\text{Vote}_i} \right)^2 \\ &= \sum_{i=1948}^{2012} \left(\text{Vote}_i - \right. \\ &\quad \left. \beta_0 - \beta_1 \text{Approval}_i - \beta_2 \text{Q1 GDP}_i - \beta_3 \text{Q2 GDP}_i - \beta_4 \text{Inc 2nd Term}_i \right)^2\end{aligned}$$

Fitting a Multivariate Regression

Choose β 's to minimize:

$$\begin{aligned}\sum_{i=1948}^{2012} \epsilon_i^2 &= \sum_{i=1948}^{2012} \left(\text{Vote}_i - \widehat{\text{Vote}_i} \right)^2 \\ &= \sum_{i=1948}^{2012} \left(\text{Vote}_i - \right. \\ &\quad \left. \beta_0 - \beta_1 \text{Approval}_i - \beta_2 \text{Q1 GDP}_i - \beta_3 \text{Q2 GDP}_i - \beta_4 \text{Inc 2nd Term}_i \right)^2\end{aligned}$$

to the R code! (Sections 1 and 2)

Multivariate Regression: Our Estimated Prediction Function

$$\text{Vote}_i = \underbrace{51.01 + 0.10 \times \text{Q1 GDP} + 0.57 \times \text{Q2 GDP}}_{\widehat{\text{Vote}_i}} + \underbrace{+0.10 \times \text{Approval} - 4.35 \times \text{Inc 2nd Term}}_{\widehat{\text{Vote}_i}} + \epsilon_i$$

Multivariate Regression: Our Estimated Prediction Function

$$\text{Vote}_i = \underbrace{51.01 + 0.10 \times \text{Q1 GDP} + 0.57 \times \text{Q2 GDP}}_{\widehat{\text{Vote}}_i} + \underbrace{+0.10 \times \text{Approval} - 4.35 \times \text{Inc 2nd Term}}_{\widehat{\text{Vote}}_i} + \epsilon_i$$

Bonus: Why does Approval have a smaller effect in the multivariate regression (.10) compared to the bivariate regression (.16)?
(Hint: Sharks and Ice-cream)

Multivariate Regression: Our Estimated Prediction Function

$$\text{Vote}_i = \underbrace{51.01 + 0.10 \times \text{Q1 GDP} + 0.57 \times \text{Q2 GDP}}_{\widehat{\text{Vote}}_i} + \underbrace{+0.10 \times \text{Approval} - 4.35 \times \text{Inc 2nd Term}}_{\widehat{\text{Vote}}_i} + \epsilon_i$$

Bonus: Why does Approval have a smaller effect in the multivariate regression (.10) compared to the bivariate regression (.16)?
(Hint: Sharks and Ice-cream)

R Code (Section 3)!

Linear Algebra

Vectors

Vector: ordered list of n numbers

- 1
- π
- e
- $(1, 2)$
- $(0, 0)$
- (π, e)
- $(3.1, 4.5, 6.11132)$
- $(\beta_0, \beta_1, \beta_2, \beta_3)$
- Incumbent vote share across elections, Net Approval across election

We will write vectors with bold (β)

Inner Product

Consider two vectors \mathbf{u} and \mathbf{v} and they are the same length. The define their inner product, $\mathbf{u} \cdot \mathbf{v}$, as

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + u_2 v_2 + \dots + u_n v_n$$

Inner Product

Consider two vectors \mathbf{u} and \mathbf{v} and they are the same length. The define their inner product, $\mathbf{u} \cdot \mathbf{v}$, as

$$\begin{aligned}\mathbf{u} \cdot \mathbf{v} &= u_1 v_1 + u_2 v_2 + \dots + u_n v_n \\ &= \sum_{i=1}^N u_i v_i\end{aligned}$$

Rewriting our Prediction

Define:

$$\begin{aligned}\hat{\beta} &= (\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3) \\ &= (51.01, 0.10, 0.57, 0.10, -4.35)\end{aligned}$$

Rewriting our Prediction

Define:

$$\begin{aligned}\hat{\beta} &= (\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3) \\ &= (51.01, 0.10, 0.57, 0.10, -4.35) \\ \mathbf{x}_i &= (1, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Approval}_i, \text{Inc 2nd Term}_i)\end{aligned}$$

Rewriting our Prediction

Define:

$$\begin{aligned}\hat{\beta} &= (\hat{\beta}_0, \hat{\beta}_1, \hat{\beta}_2, \hat{\beta}_3) \\ &= (51.01, 0.10, 0.57, 0.10, -4.35) \\ \mathbf{x}_i &= (1, \text{Q1 GDP}_i, \text{Q2 GDP}_i, \text{Approval}_i, \text{Inc 2nd Term}_i)\end{aligned}$$

Then, we can write our prediction as:

$$\widehat{\text{Vote}}_i = \hat{\beta} \cdot \mathbf{x}_i$$

R Code (Sections 4 5)

Potential Problems

See *Introduction to Statistical Learning* 3.3.3 (p. 92–102)

- 1 Non-linearity of response-predictor relationships:** true relationship is not linear.
- 2 Correlation of error terms:** ϵ_i provides information about ϵ_{i+1} , as in time-series.
- 3 Non-constant variance of error terms (heteroscedasticity):** variances of error terms change with value of response, making a funnel shape.
- 4 Outliers and high-leverage points:** unusual values for Y or X .
- 5 Collinearity:** two or more predictor variables are closely related to one another.

Key Terms:

- Linear Regression, Simple Regression, Multiple Regression
- Cost function
- Sum of Squared Residuals
- In sample, Out of Sample

Key Techniques and R Functions

- Linear algebra operations and terms
 - Inner product
 - Matrix
- `lm`, `plot`, `%*%`

Up next:
Classification and Logistic Regression