


Solving multivariate quadratic systems in practice

Monika Trimoska

Summer school on RWC and privacy
June 30, Dubrovnik, Croatia




Algebraic cryptanalysis




A type of cryptanalytic methods where the problem of finding the secret key (or any attack goal) is **reduced** to the problem of finding a solution to a **nonlinear multivariate polynomial system of equations**.


Algebraic cryptanalysis




Algebraic cryptanalysis



Algebraic cryptanalysis



Algebraic cryptanalysis



The MQ problem (recall)

The MQ problem

Given m multivariate quadratic polynomials f_1, \dots, f_m of n variables over a finite field \mathbb{F}_q , find a tuple $\mathbf{x} = (x_1, \dots, x_n)$ in \mathbb{F}_q^n , such that $f_1(\mathbf{x}) = \dots = f_m(\mathbf{x}) = 0$.

Example.

$$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$$

$$f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$$

$$f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$$

$$f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$$

$$f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$$

$$f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$$

Algebraic cryptanalysis : modelisation example

Example.

Given matrices $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2 \in \mathcal{M}_{n,n}(\mathbb{F}_q)$ (the space of matrices over \mathbb{F}_q of size $n \times n$), find $\mathbf{A}, \mathbf{B} \in \text{GL}_n(\mathbb{F}_q)$ (the space of invertible matrices over \mathbb{F}_q of size $n \times n$), such that

$$\mathbf{D}_1 = \mathbf{A} \mathbf{C}_1 \mathbf{B}$$

$$\mathbf{D}_2 = \mathbf{A} \mathbf{C}_2 \mathbf{B}$$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A}\mathbf{C}_1\mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A}\mathbf{C}_2\mathbf{B}$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

\mathbf{A}

$a_{1,1}$	$a_{1,2}$	$a_{1,3}$	$a_{1,4}$	$a_{1,5}$
$a_{2,1}$	$a_{2,2}$	$a_{2,3}$	$a_{2,4}$	$a_{2,5}$
$a_{3,1}$	$a_{3,2}$	$a_{3,3}$	$a_{3,4}$	$a_{3,5}$
$a_{4,1}$	$a_{4,2}$	$a_{4,3}$	$a_{4,4}$	$a_{4,5}$
$a_{5,1}$	$a_{5,2}$	$a_{5,3}$	$a_{5,4}$	$a_{5,5}$

\mathbf{C}_1

$c_{1,1}$	$c_{1,2}$	$c_{1,3}$	$c_{1,4}$	$c_{1,5}$
$c_{2,1}$	$c_{2,2}$	$c_{2,3}$	$c_{2,4}$	$c_{2,5}$
$c_{3,1}$	$c_{3,2}$	$c_{3,3}$	$c_{3,4}$	$c_{3,5}$
$c_{4,1}$	$c_{4,2}$	$c_{4,3}$	$c_{4,4}$	$c_{4,5}$
$c_{5,1}$	$c_{5,2}$	$c_{5,3}$	$c_{5,4}$	$c_{5,5}$

\mathbf{B}

$b_{1,1}$	$b_{1,2}$	$b_{1,3}$	$b_{1,4}$	$b_{1,5}$
$b_{2,1}$	$b_{2,2}$	$b_{2,3}$	$b_{2,4}$	$b_{2,5}$
$b_{3,1}$	$b_{3,2}$	$b_{3,3}$	$b_{3,4}$	$b_{3,5}$
$b_{4,1}$	$b_{4,2}$	$b_{4,3}$	$b_{4,4}$	$b_{4,5}$
$b_{5,1}$	$b_{5,2}$	$b_{5,3}$	$b_{5,4}$	$b_{5,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A} \mathbf{C}_1 \mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A} \mathbf{C}_2 \mathbf{B}$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

$\mathbf{A} \mathbf{C}_1 \mathbf{B}$

$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A}\mathbf{C}_1\mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A}\mathbf{C}_2\mathbf{B}$

$$d_{1,1} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1} = 0,$$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

$$\mathbf{A}\mathbf{C}_1\mathbf{B}$$

$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A}\mathbf{C}_1\mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A}\mathbf{C}_2\mathbf{B}$

$$d_{1,1} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1} = 0,$$

$$d_{2,1} - \sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1} = 0,$$

$$\mathbf{A}\mathbf{C}_1\mathbf{B}$$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A} \mathbf{C}_1 \mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A} \mathbf{C}_2 \mathbf{B}$

$$d_{1,1} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1} = 0, \quad d_{1,2} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2} = 0, \dots$$

$$d_{2,1} - \sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1} = 0,$$

$\mathbf{A} \mathbf{C}_1 \mathbf{B}$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{D}_1 = \mathbf{A} \mathbf{C}_1 \mathbf{B}$
 $\mathbf{D}_2 = \mathbf{A} \mathbf{C}_2 \mathbf{B}$

$$d_{1,1} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1} = 0, \quad d_{1,2} - \sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2} = 0, \dots$$

$$d_{2,1} - \sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1} = 0, \quad d_{l,p} - \sum_{i=1}^n \sum_{j=1}^n a_{l,j} c_{j,i} b_{i,p} = 0$$

$\mathbf{A} \mathbf{C}_1 \mathbf{B}$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$

=

$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{1,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{2,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{3,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{4,j} c_{j,i} b_{i,5}$
$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,1}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,2}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,3}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,4}$	$\sum_{i=1}^n \sum_{j=1}^n a_{5,j} c_{j,i} b_{i,5}$

Algebraic cryptanalysis : modelisation example



A motivating example: a better idea for modelisation.

Given matrices $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2 \in \mathcal{M}_{n,n}(\mathbb{F}_q)$ (the space of matrices over \mathbb{F}_q of size $n \times n$), find $\mathbf{A}, \mathbf{B} \in \text{GL}_n(\mathbb{F}_q)$ (the space of invertible matrices over \mathbb{F}_q of size $n \times n$), such that

$$\begin{aligned}\mathbf{A}^{-1}\mathbf{D}_1 &= \mathbf{C}_1\mathbf{B} \\ \mathbf{A}^{-1}\mathbf{D}_2 &= \mathbf{C}_2\mathbf{B}\end{aligned}$$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$
 $\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$

\mathbf{A}^{-1}

$a_{1,1}$	$a_{1,2}$	$a_{1,3}$	$a_{1,4}$	$a_{1,5}$
$a_{2,1}$	$a_{2,2}$	$a_{2,3}$	$a_{2,4}$	$a_{2,5}$
$a_{3,1}$	$a_{3,2}$	$a_{3,3}$	$a_{3,4}$	$a_{3,5}$
$a_{4,1}$	$a_{4,2}$	$a_{4,3}$	$a_{4,4}$	$a_{4,5}$
$a_{5,1}$	$a_{5,2}$	$a_{5,3}$	$a_{5,4}$	$a_{5,5}$

\mathbf{D}_1

$d_{1,1}$	$d_{1,2}$	$d_{1,3}$	$d_{1,4}$	$d_{1,5}$
$d_{2,1}$	$d_{2,2}$	$d_{2,3}$	$d_{2,4}$	$d_{2,5}$
$d_{3,1}$	$d_{3,2}$	$d_{3,3}$	$d_{3,4}$	$d_{3,5}$
$d_{4,1}$	$d_{4,2}$	$d_{4,3}$	$d_{4,4}$	$d_{4,5}$
$d_{5,1}$	$d_{5,2}$	$d_{5,3}$	$d_{5,4}$	$d_{5,5}$


=

\mathbf{C}_1

$c_{1,1}$	$c_{1,2}$	$c_{1,3}$	$c_{1,4}$	$c_{1,5}$
$c_{2,1}$	$c_{2,2}$	$c_{2,3}$	$c_{2,4}$	$c_{2,5}$
$c_{3,1}$	$c_{3,2}$	$c_{3,3}$	$c_{3,4}$	$c_{3,5}$
$c_{4,1}$	$c_{4,2}$	$c_{4,3}$	$c_{4,4}$	$c_{4,5}$
$c_{5,1}$	$c_{5,2}$	$c_{5,3}$	$c_{5,4}$	$c_{5,5}$

\mathbf{B}

$b_{1,1}$	$b_{1,2}$	$b_{1,3}$	$b_{1,4}$	$b_{1,5}$
$b_{2,1}$	$b_{2,2}$	$b_{2,3}$	$b_{2,4}$	$b_{2,5}$
$b_{3,1}$	$b_{3,2}$	$b_{3,3}$	$b_{3,4}$	$b_{3,5}$
$b_{4,1}$	$b_{4,2}$	$b_{4,3}$	$b_{4,4}$	$b_{4,5}$
$b_{5,1}$	$b_{5,2}$	$b_{5,3}$	$b_{5,4}$	$b_{5,5}$



Algebraic cryptanalysis : modelisation example

Example. $\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$
 $\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$

$\mathbf{A}^{-1}\mathbf{D}_1$

$\sum_{i=1}^n a_{1,i}d_{i,1}$	$\sum_{i=1}^n a_{1,i}d_{i,2}$	$\sum_{i=1}^n a_{1,i}d_{i,3}$	$\sum_{i=1}^n a_{1,i}d_{i,4}$	$\sum_{i=1}^n a_{1,i}d_{i,5}$
$\sum_{i=1}^n a_{2,i}d_{i,1}$	$\sum_{i=1}^n a_{2,i}d_{i,2}$	$\sum_{i=1}^n a_{2,i}d_{i,3}$	$\sum_{i=1}^n a_{2,i}d_{i,4}$	$\sum_{i=1}^n a_{2,i}d_{i,5}$
$\sum_{i=1}^n a_{3,i}d_{i,1}$	$\sum_{i=1}^n a_{3,i}d_{i,2}$	$\sum_{i=1}^n a_{3,i}d_{i,3}$	$\sum_{i=1}^n a_{3,i}d_{i,4}$	$\sum_{i=1}^n a_{3,i}d_{i,5}$
$\sum_{i=1}^n a_{4,i}d_{i,1}$	$\sum_{i=1}^n a_{4,i}d_{i,2}$	$\sum_{i=1}^n a_{4,i}d_{i,3}$	$\sum_{i=1}^n a_{4,i}d_{i,4}$	$\sum_{i=1}^n a_{4,i}d_{i,5}$
$\sum_{i=1}^n a_{5,i}d_{i,1}$	$\sum_{i=1}^n a_{5,i}d_{i,2}$	$\sum_{i=1}^n a_{5,i}d_{i,3}$	$\sum_{i=1}^n a_{5,i}d_{i,4}$	$\sum_{i=1}^n a_{5,i}d_{i,5}$

=

$\mathbf{C}_1\mathbf{B}$

$\sum_{i=1}^n c_{1,i}b_{i,1}$	$\sum_{i=1}^n c_{1,i}b_{i,2}$	$\sum_{i=1}^n c_{1,i}b_{i,3}$	$\sum_{i=1}^n c_{1,i}b_{i,4}$	$\sum_{i=1}^n c_{1,i}b_{i,5}$
$\sum_{i=1}^n c_{2,i}b_{i,1}$	$\sum_{i=1}^n c_{2,i}b_{i,2}$	$\sum_{i=1}^n c_{2,i}b_{i,3}$	$\sum_{i=1}^n c_{2,i}b_{i,4}$	$\sum_{i=1}^n c_{2,i}b_{i,5}$
$\sum_{i=1}^n c_{3,i}b_{i,1}$	$\sum_{i=1}^n c_{3,i}b_{i,2}$	$\sum_{i=1}^n c_{3,i}b_{i,3}$	$\sum_{i=1}^n c_{3,i}b_{i,4}$	$\sum_{i=1}^n c_{3,i}b_{i,5}$
$\sum_{i=1}^n c_{4,i}b_{i,1}$	$\sum_{i=1}^n c_{4,i}b_{i,2}$	$\sum_{i=1}^n c_{4,i}b_{i,3}$	$\sum_{i=1}^n c_{4,i}b_{i,4}$	$\sum_{i=1}^n c_{4,i}b_{i,5}$
$\sum_{i=1}^n c_{5,i}b_{i,1}$	$\sum_{i=1}^n c_{5,i}b_{i,2}$	$\sum_{i=1}^n c_{5,i}b_{i,3}$	$\sum_{i=1}^n c_{5,i}b_{i,4}$	$\sum_{i=1}^n c_{5,i}b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$
 $\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$

$$\sum_{i=1}^n a_{1,i}d_{i,1} - \sum_{i=1}^n c_{1,i}b_{i,1} = 0,$$

$$\mathbf{A}^{-1}\mathbf{D}_1$$

$\sum_{i=1}^n a_{1,i}d_{i,1}$	$\sum_{i=1}^n a_{1,i}d_{i,2}$	$\sum_{i=1}^n a_{1,i}d_{i,3}$	$\sum_{i=1}^n a_{1,i}d_{i,4}$	$\sum_{i=1}^n a_{1,i}d_{i,5}$
$\sum_{i=1}^n a_{2,i}d_{i,1}$	$\sum_{i=1}^n a_{2,i}d_{i,2}$	$\sum_{i=1}^n a_{2,i}d_{i,3}$	$\sum_{i=1}^n a_{2,i}d_{i,4}$	$\sum_{i=1}^n a_{2,i}d_{i,5}$
$\sum_{i=1}^n a_{3,i}d_{i,1}$	$\sum_{i=1}^n a_{3,i}d_{i,2}$	$\sum_{i=1}^n a_{3,i}d_{i,3}$	$\sum_{i=1}^n a_{3,i}d_{i,4}$	$\sum_{i=1}^n a_{3,i}d_{i,5}$
$\sum_{i=1}^n a_{4,i}d_{i,1}$	$\sum_{i=1}^n a_{4,i}d_{i,2}$	$\sum_{i=1}^n a_{4,i}d_{i,3}$	$\sum_{i=1}^n a_{4,i}d_{i,4}$	$\sum_{i=1}^n a_{4,i}d_{i,5}$
$\sum_{i=1}^n a_{5,i}d_{i,1}$	$\sum_{i=1}^n a_{5,i}d_{i,2}$	$\sum_{i=1}^n a_{5,i}d_{i,3}$	$\sum_{i=1}^n a_{5,i}d_{i,4}$	$\sum_{i=1}^n a_{5,i}d_{i,5}$

=

$$\mathbf{C}_1\mathbf{B}$$

$\sum_{i=1}^n c_{1,i}b_{i,1}$	$\sum_{i=1}^n c_{1,i}b_{i,2}$	$\sum_{i=1}^n c_{1,i}b_{i,3}$	$\sum_{i=1}^n c_{1,i}b_{i,4}$	$\sum_{i=1}^n c_{1,i}b_{i,5}$
$\sum_{i=1}^n c_{2,i}b_{i,1}$	$\sum_{i=1}^n c_{2,i}b_{i,2}$	$\sum_{i=1}^n c_{2,i}b_{i,3}$	$\sum_{i=1}^n c_{2,i}b_{i,4}$	$\sum_{i=1}^n c_{2,i}b_{i,5}$
$\sum_{i=1}^n c_{3,i}b_{i,1}$	$\sum_{i=1}^n c_{3,i}b_{i,2}$	$\sum_{i=1}^n c_{3,i}b_{i,3}$	$\sum_{i=1}^n c_{3,i}b_{i,4}$	$\sum_{i=1}^n c_{3,i}b_{i,5}$
$\sum_{i=1}^n c_{4,i}b_{i,1}$	$\sum_{i=1}^n c_{4,i}b_{i,2}$	$\sum_{i=1}^n c_{4,i}b_{i,3}$	$\sum_{i=1}^n c_{4,i}b_{i,4}$	$\sum_{i=1}^n c_{4,i}b_{i,5}$
$\sum_{i=1}^n c_{5,i}b_{i,1}$	$\sum_{i=1}^n c_{5,i}b_{i,2}$	$\sum_{i=1}^n c_{5,i}b_{i,3}$	$\sum_{i=1}^n c_{5,i}b_{i,4}$	$\sum_{i=1}^n c_{5,i}b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$
 $\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$

$\mathbf{A}^{-1}\mathbf{D}_1$

$\sum_{i=1}^n a_{1,i}d_{i,1}$	$\sum_{i=1}^n a_{1,i}d_{i,2}$	$\sum_{i=1}^n a_{1,i}d_{i,3}$	$\sum_{i=1}^n a_{1,i}d_{i,4}$	$\sum_{i=1}^n a_{1,i}d_{i,5}$
$\sum_{i=1}^n a_{2,i}d_{i,1}$	$\sum_{i=1}^n a_{2,i}d_{i,2}$	$\sum_{i=1}^n a_{2,i}d_{i,3}$	$\sum_{i=1}^n a_{2,i}d_{i,4}$	$\sum_{i=1}^n a_{2,i}d_{i,5}$
$\sum_{i=1}^n a_{3,i}d_{i,1}$	$\sum_{i=1}^n a_{3,i}d_{i,2}$	$\sum_{i=1}^n a_{3,i}d_{i,3}$	$\sum_{i=1}^n a_{3,i}d_{i,4}$	$\sum_{i=1}^n a_{3,i}d_{i,5}$
$\sum_{i=1}^n a_{4,i}d_{i,1}$	$\sum_{i=1}^n a_{4,i}d_{i,2}$	$\sum_{i=1}^n a_{4,i}d_{i,3}$	$\sum_{i=1}^n a_{4,i}d_{i,4}$	$\sum_{i=1}^n a_{4,i}d_{i,5}$
$\sum_{i=1}^n a_{5,i}d_{i,1}$	$\sum_{i=1}^n a_{5,i}d_{i,2}$	$\sum_{i=1}^n a_{5,i}d_{i,3}$	$\sum_{i=1}^n a_{5,i}d_{i,4}$	$\sum_{i=1}^n a_{5,i}d_{i,5}$

=

$\mathbf{C}_1\mathbf{B}$

$\sum_{i=1}^n c_{1,i}b_{i,1}$	$\sum_{i=1}^n c_{1,i}b_{i,2}$	$\sum_{i=1}^n c_{1,i}b_{i,3}$	$\sum_{i=1}^n c_{1,i}b_{i,4}$	$\sum_{i=1}^n c_{1,i}b_{i,5}$
$\sum_{i=1}^n c_{2,i}b_{i,1}$	$\sum_{i=1}^n c_{2,i}b_{i,2}$	$\sum_{i=1}^n c_{2,i}b_{i,3}$	$\sum_{i=1}^n c_{2,i}b_{i,4}$	$\sum_{i=1}^n c_{2,i}b_{i,5}$
$\sum_{i=1}^n c_{3,i}b_{i,1}$	$\sum_{i=1}^n c_{3,i}b_{i,2}$	$\sum_{i=1}^n c_{3,i}b_{i,3}$	$\sum_{i=1}^n c_{3,i}b_{i,4}$	$\sum_{i=1}^n c_{3,i}b_{i,5}$
$\sum_{i=1}^n c_{4,i}b_{i,1}$	$\sum_{i=1}^n c_{4,i}b_{i,2}$	$\sum_{i=1}^n c_{4,i}b_{i,3}$	$\sum_{i=1}^n c_{4,i}b_{i,4}$	$\sum_{i=1}^n c_{4,i}b_{i,5}$
$\sum_{i=1}^n c_{5,i}b_{i,1}$	$\sum_{i=1}^n c_{5,i}b_{i,2}$	$\sum_{i=1}^n c_{5,i}b_{i,3}$	$\sum_{i=1}^n c_{5,i}b_{i,4}$	$\sum_{i=1}^n c_{5,i}b_{i,5}$

Algebraic cryptanalysis : modelisation example

Example. $\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$
 $\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$

$\mathbf{A}^{-1}\mathbf{D}_1$

$\sum_{i=1}^n a_{1,i}d_{i,1}$	$\sum_{i=1}^n a_{1,i}d_{i,2}$	$\sum_{i=1}^n a_{1,i}d_{i,3}$	$\sum_{i=1}^n a_{1,i}d_{i,4}$	$\sum_{i=1}^n a_{1,i}d_{i,5}$
$\sum_{i=1}^n a_{2,i}d_{i,1}$	$\sum_{i=1}^n a_{2,i}d_{i,2}$	$\sum_{i=1}^n a_{2,i}d_{i,3}$	$\sum_{i=1}^n a_{2,i}d_{i,4}$	$\sum_{i=1}^n a_{2,i}d_{i,5}$
$\sum_{i=1}^n a_{3,i}d_{i,1}$	$\sum_{i=1}^n a_{3,i}d_{i,2}$	$\sum_{i=1}^n a_{3,i}d_{i,3}$	$\sum_{i=1}^n a_{3,i}d_{i,4}$	$\sum_{i=1}^n a_{3,i}d_{i,5}$
$\sum_{i=1}^n a_{4,i}d_{i,1}$	$\sum_{i=1}^n a_{4,i}d_{i,2}$	$\sum_{i=1}^n a_{4,i}d_{i,3}$	$\sum_{i=1}^n a_{4,i}d_{i,4}$	$\sum_{i=1}^n a_{4,i}d_{i,5}$
$\sum_{i=1}^n a_{5,i}d_{i,1}$	$\sum_{i=1}^n a_{5,i}d_{i,2}$	$\sum_{i=1}^n a_{5,i}d_{i,3}$	$\sum_{i=1}^n a_{5,i}d_{i,4}$	$\sum_{i=1}^n a_{5,i}d_{i,5}$

=

$\mathbf{C}_1\mathbf{B}$

$\sum_{i=1}^n c_{1,i}b_{i,1}$	$\sum_{i=1}^n c_{1,i}b_{i,2}$	$\sum_{i=1}^n c_{1,i}b_{i,3}$	$\sum_{i=1}^n c_{1,i}b_{i,4}$	$\sum_{i=1}^n c_{1,i}b_{i,5}$
$\sum_{i=1}^n c_{2,i}b_{i,1}$	$\sum_{i=1}^n c_{2,i}b_{i,2}$	$\sum_{i=1}^n c_{2,i}b_{i,3}$	$\sum_{i=1}^n c_{2,i}b_{i,4}$	$\sum_{i=1}^n c_{2,i}b_{i,5}$
$\sum_{i=1}^n c_{3,i}b_{i,1}$	$\sum_{i=1}^n c_{3,i}b_{i,2}$	$\sum_{i=1}^n c_{3,i}b_{i,3}$	$\sum_{i=1}^n c_{3,i}b_{i,4}$	$\sum_{i=1}^n c_{3,i}b_{i,5}$
$\sum_{i=1}^n c_{4,i}b_{i,1}$	$\sum_{i=1}^n c_{4,i}b_{i,2}$	$\sum_{i=1}^n c_{4,i}b_{i,3}$	$\sum_{i=1}^n c_{4,i}b_{i,4}$	$\sum_{i=1}^n c_{4,i}b_{i,5}$
$\sum_{i=1}^n c_{5,i}b_{i,1}$	$\sum_{i=1}^n c_{5,i}b_{i,2}$	$\sum_{i=1}^n c_{5,i}b_{i,3}$	$\sum_{i=1}^n c_{5,i}b_{i,4}$	$\sum_{i=1}^n c_{5,i}b_{i,5}$

$\sum_{i=1}^n a_{1,i}d_{i,1} - \sum_{i=1}^n c_{1,i}b_{i,1} = 0, \quad \sum_{i=1}^n a_{1,i}d_{i,2} - \sum_{i=1}^n c_{1,i}b_{i,2} = 0, \dots$

$\sum_{i=1}^n a_{2,i}d_{i,1} - \sum_{i=1}^n c_{2,i}b_{i,1} = 0,$

Algebraic cryptanalysis : modelisation example

Example.

$$\mathbf{A}^{-1}\mathbf{D}_1 = \mathbf{C}_1\mathbf{B}$$

$$\mathbf{A}^{-1}\mathbf{D}_2 = \mathbf{C}_2\mathbf{B}$$

$$\mathbf{A}^{-1}\mathbf{D}_1$$

$\sum_{i=1}^n a_{1,i}d_{i,1}$	$\sum_{i=1}^n a_{1,i}d_{i,2}$	$\sum_{i=1}^n a_{1,i}d_{i,3}$	$\sum_{i=1}^n a_{1,i}d_{i,4}$	$\sum_{i=1}^n a_{1,i}d_{i,5}$
$\sum_{i=1}^n a_{2,i}d_{i,1}$	$\sum_{i=1}^n a_{2,i}d_{i,2}$	$\sum_{i=1}^n a_{2,i}d_{i,3}$	$\sum_{i=1}^n a_{2,i}d_{i,4}$	$\sum_{i=1}^n a_{2,i}d_{i,5}$
$\sum_{i=1}^n a_{3,i}d_{i,1}$	$\sum_{i=1}^n a_{3,i}d_{i,2}$	$\sum_{i=1}^n a_{3,i}d_{i,3}$	$\sum_{i=1}^n a_{3,i}d_{i,4}$	$\sum_{i=1}^n a_{3,i}d_{i,5}$
$\sum_{i=1}^n a_{4,i}d_{i,1}$	$\sum_{i=1}^n a_{4,i}d_{i,2}$	$\sum_{i=1}^n a_{4,i}d_{i,3}$	$\sum_{i=1}^n a_{4,i}d_{i,4}$	$\sum_{i=1}^n a_{4,i}d_{i,5}$
$\sum_{i=1}^n a_{5,i}d_{i,1}$	$\sum_{i=1}^n a_{5,i}d_{i,2}$	$\sum_{i=1}^n a_{5,i}d_{i,3}$	$\sum_{i=1}^n a_{5,i}d_{i,4}$	$\sum_{i=1}^n a_{5,i}d_{i,5}$

=

$$\mathbf{C}_1\mathbf{B}$$

$$\sum_{i=1}^n a_{1,i}d_{i,1} - \sum_{i=1}^n c_{1,i}b_{i,1} = 0, \quad \sum_{i=1}^n a_{1,i}d_{i,2} - \sum_{i=1}^n c_{1,i}b_{i,2} = 0, \dots$$

$$\sum_{i=1}^n a_{2,i}d_{i,1} - \sum_{i=1}^n c_{2,i}b_{i,1} = 0, \quad \sum_{i=1}^n a_{l,i}d_{i,p} - \sum_{i=1}^n c_{l,i}b_{i,p} = 0$$

$\sum_{i=1}^n c_{1,i}b_{i,1}$	$\sum_{i=1}^n c_{1,i}b_{i,2}$	$\sum_{i=1}^n c_{1,i}b_{i,3}$	$\sum_{i=1}^n c_{1,i}b_{i,4}$	$\sum_{i=1}^n c_{1,i}b_{i,5}$
$\sum_{i=1}^n c_{2,i}b_{i,1}$	$\sum_{i=1}^n c_{2,i}b_{i,2}$	$\sum_{i=1}^n c_{2,i}b_{i,3}$	$\sum_{i=1}^n c_{2,i}b_{i,4}$	$\sum_{i=1}^n c_{2,i}b_{i,5}$
$\sum_{i=1}^n c_{3,i}b_{i,1}$	$\sum_{i=1}^n c_{3,i}b_{i,2}$	$\sum_{i=1}^n c_{3,i}b_{i,3}$	$\sum_{i=1}^n c_{3,i}b_{i,4}$	$\sum_{i=1}^n c_{3,i}b_{i,5}$
$\sum_{i=1}^n c_{4,i}b_{i,1}$	$\sum_{i=1}^n c_{4,i}b_{i,2}$	$\sum_{i=1}^n c_{4,i}b_{i,3}$	$\sum_{i=1}^n c_{4,i}b_{i,4}$	$\sum_{i=1}^n c_{4,i}b_{i,5}$
$\sum_{i=1}^n c_{5,i}b_{i,1}$	$\sum_{i=1}^n c_{5,i}b_{i,2}$	$\sum_{i=1}^n c_{5,i}b_{i,3}$	$\sum_{i=1}^n c_{5,i}b_{i,4}$	$\sum_{i=1}^n c_{5,i}b_{i,5}$

Algebraic cryptanalysis : modelisation example




A motivating example: a better idea for modelisation.

Given matrices $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2 \in \mathcal{M}_{n,n}(\mathbb{F}_q)$ (the space of matrices over \mathbb{F}_q of size $n \times n$), find $\mathbf{A}, \mathbf{B} \in \text{GL}_n(\mathbb{F}_q)$ (the space of invertible matrices over \mathbb{F}_q of size $n \times n$), such that

$$\begin{aligned}\mathbf{A}^{-1}\mathbf{D}_1 &= \mathbf{C}_1\mathbf{B} \\ \mathbf{A}^{-1}\mathbf{D}_2 &= \mathbf{C}_2\mathbf{B}\end{aligned}$$

Algebraic cryptanalysis : modelisation example




A motivating example: a better idea for modelisation.

Given matrices $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2 \in \mathcal{M}_{n,n}(\mathbb{F}_q)$ (the space of matrices over \mathbb{F}_q of size $n \times n$), find $\mathbf{A}, \mathbf{B} \in \text{GL}_n(\mathbb{F}_q)$ (the space of invertible matrices over \mathbb{F}_q of size $n \times n$), such that

$$\begin{aligned}\mathbf{A}^{-1}\mathbf{D}_1 &= \mathbf{C}_1\mathbf{B} \\ \mathbf{A}^{-1}\mathbf{D}_2 &= \mathbf{C}_2\mathbf{B}\end{aligned}$$

→ Results in a **linear** system with the same number of variables and equations.

Algebraic cryptanalysis : modelisation example




A motivating example: a better idea for modelisation.

Given matrices $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2 \in \mathcal{M}_{n,n}(\mathbb{F}_q)$ (the space of matrices over \mathbb{F}_q of size $n \times n$), find $\mathbf{A}, \mathbf{B} \in \text{GL}_n(\mathbb{F}_q)$ (the space of invertible matrices over \mathbb{F}_q of size $n \times n$), such that

$$\begin{aligned}\mathbf{A}^{-1}\mathbf{D}_1 &= \mathbf{C}_1\mathbf{B} \\ \mathbf{A}^{-1}\mathbf{D}_2 &= \mathbf{C}_2\mathbf{B}\end{aligned}$$

- Results in a **linear** system with the same number of variables and equations.
- If $\mathbf{C}_1, \mathbf{C}_2, \mathbf{D}_1, \mathbf{D}_2$ are all full rank, we should have a unique solution.
- We can easily recover \mathbf{A} from \mathbf{A}^{-1} .

Overview of solvers







(Fast) Exhaustive Search

[Bouillaguet, Chen, Cheng, Chou, Niederhagen, Shamir, Yang, 2010]

Exhaustive Search



$$x_1 \cdot x_2 + x_1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$

$$x_2 \cdot x_3 + x_2 \cdot x_4 + x_1 + x_2 + 1 = 0$$

$$x_1 \cdot x_2 + x_2 \cdot x_3 + x_2 \cdot x_4 + x_1 + x_4 = 0$$



$$x_1 \cdot x_4 + x_2 \cdot x_3 + x_2 + x_3 + x_4 = 0$$

Binary search tree

Exhaustive Search



Worst-case complexity: $\mathcal{O}(2^n)$



Binary search tree



$$x_1 \cdot x_2 + x_1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$

$$x_2 \cdot x_3 + x_2 \cdot x_4 + x_1 + x_2 + 1 = 0$$

$$x_1 \cdot x_2 + x_2 \cdot x_3 + x_2 \cdot x_4 + x_1 + x_4 = 0$$

$$x_1 \cdot x_4 + x_2 \cdot x_3 + x_2 + x_3 + x_4 = 0$$

Exhaustive Search





Binary search tree

$$\begin{aligned}1 \cdot 0 + 1 \cdot 0 + 0 \cdot 1 + 0 &= 0 \\0 \cdot 0 + 0 \cdot 1 + 1 + 0 + 1 &= 0 \\1 \cdot 0 + 0 \cdot 0 + 0 \cdot 1 + 1 + 1 &= 0 \\1 \cdot 1 + 0 \cdot 0 + 0 + 0 + 1 &= 0\end{aligned}$$

Exhaustive Search



Worst-case complexity: $\mathcal{O}(2^n)$



Binary search tree

$$\begin{aligned} 1 \cdot 0 + 1 \cdot 0 + 0 \cdot 1 + 0 &= 0 \\ 0 \cdot 0 + 0 \cdot 1 + 1 + 0 + 1 &= 0 \\ 1 \cdot 0 + 0 \cdot 0 + 0 \cdot 1 + 1 + 1 &= 0 \\ 1 \cdot 1 + 0 \cdot 0 + 0 + 0 + 1 &= 0 \end{aligned}$$

Fast Exhaustive Search

* The libFES solver

Gray code

- An ordering of the binary system where two successive values **differ in only one bit**.


Example. $n = 4$

0000	1100
0001	1101
0011	1111
0010	1110
0110	1010
0111	1011
0101	1001
0100	1000

Fast Exhaustive Search

Gray code

0000	1100
0001	1101
0011	1111
0010	1110
0110	1010
0111	1011
0101	1001
0100	1000



$$\begin{aligned}1 \cdot 0 + 1 \cdot 0 + 0 \cdot 1 + 0 &= 0 \\0 \cdot 0 + 0 \cdot 1 + 1 + 0 + 1 &= 0 \\1 \cdot 0 + 0 \cdot 0 + 0 \cdot 1 + 1 + 1 &= 0 \\1 \cdot 1 + 0 \cdot 0 + 0 + 0 + 1 &= 0\end{aligned}$$

Fast Exhaustive Search




Worst-case complexity: $\mathcal{O}(2^n)$

! But, it differs from the depth-first traversal in the polynomial factors


Gray code

0000	1100
0001	1101
0011	1111
0010	1110
0110	1010
0111	1011
0101	1001
0100	1000



$$\begin{aligned}1 \cdot 0 + 1 \cdot 0 + 0 \cdot 1 + 0 &= 0 \\0 \cdot 0 + 0 \cdot 1 + 1 + 0 + 1 &= 0 \\1 \cdot 0 + 0 \cdot 0 + 0 \cdot 1 + 1 + 1 &= 0 \\1 \cdot 1 + 0 \cdot 0 + 0 + 0 + 1 &= 0\end{aligned}$$

Overview of solvers



The background of the slide features three interlocking gears of different sizes, rendered in a light red color against a dark red background. The gears are positioned such that they appear to be meshing together, with one gear at the top and two at the bottom.

SAT solvers

CryptoMiniSat [Soos, Nohl, Castelluccia, 2009], WDSat [T., Dequen, Ionica, 2020]

Simple algorithm

[Bouillaguet, Delaplace, T., 2021]

(SAT solvers)

- **Propositional formula** in Conjunctive Normal Form (**CNF**): a **conjunction of clauses** where each clause is a **disjunction of literals** and where each **literal** is a variable or a negated variable.

Example. $(x_1 \vee \neg x_2) \wedge$
 $(x_2 \vee x_3 \vee x_4) \wedge$
 $(\neg x_1 \vee x_4)$

(SAT solvers)

- **Propositional formula** in Conjunctive Normal Form (**CNF**): a **conjunction of clauses** where each clause is a **disjunction of literals** and where each **literal** is a variable or a negated variable.

Example. $(x_1 \vee \neg x_2) \wedge$
 $(x_2 \vee x_3 \vee x_4) \wedge$
 $(\neg x_1 \vee x_4)$

The SATisfiability problem

Given a propositional formula, determine whether there exists an interpretation (assignment of all variables) such that the formula is satisfied (evaluates to TRUE).

(SAT solvers)

- **Propositional formula** in Conjunctive Normal Form (**CNF**): a **conjunction of clauses** where each clause is a **disjunction of literals** and where each **literal** is a variable or a negated variable.

Example. $(x_1 \vee \neg x_2) \wedge$
 $(x_2 \vee x_3 \vee x_4) \wedge$
 $(\neg x_1 \vee x_4)$


The SATisfiability problem

Given a propositional formula, determine whether there exists an interpretation (assignment of all variables) such that the formula is satisfied (evaluates to TRUE).



SAT solver: a tool for solving the SAT problem.

Partial assignment and conflicts



$$1 \cdot 0 + 1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$


$$0 \cdot x_3 + 0 \cdot x_4 + 1 + 0 + 1 = 0$$

$$1 \cdot 0 + 0 \cdot x_3 + 0 \cdot x_4 + 1 + x_4 = 0$$

$$1 \cdot x_4 + 0 \cdot x_3 + 0 + x_3 + x_4 = 0$$

Partial assignment and conflicts

Which (portion of) branches are missing ??



$$1 \cdot 0 + 1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$

$$0 \cdot x_3 + 0 \cdot x_4 + 1 + 0 + 1 = 0$$


$$1 \cdot 0 + 0 \cdot x_3 + 0 \cdot x_4 + 1 + x_4 = 0$$

$$1 \cdot x_4 + 0 \cdot x_3 + 0 + x_3 + x_4 = 0$$

Partial assignment and conflicts

Which (portion of) branches are missing ??

Worst-case complexity: $\mathcal{O}(2^n)$



$$1 \cdot 0 + 1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$

$$0 \cdot x_3 + 0 \cdot x_4 + 1 + 0 + 1 = 0$$


$$1 \cdot 0 + 0 \cdot x_3 + 0 \cdot x_4 + 1 + x_4 = 0$$

$$1 \cdot x_4 + 0 \cdot x_3 + 0 + x_3 + x_4 = 0$$

Partial assignment and conflicts

Which (portion of) branches are missing ??

Worst-case complexity: $\mathcal{O}(2^n)$



$$1 \cdot 0 + 1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$


$$0 \cdot x_3 + 0 \cdot x_4 + 1 + 0 + 1 = 0$$

$$1 \cdot 0 + 0 \cdot x_3 + 0 \cdot x_4 + 1 + x_4 = 0$$

$$1 \cdot x_4 + 0 \cdot x_3 + 0 + x_3 + x_4 = 0$$

XOR-enabled SAT solvers: take as input XOR constraints as well; perform Gaussian elimination;
*CryptoMiniSat, WDSat

Overview of solvers




Macaulay matrix

Linearisation

Linear systems are easy to solve, nonlinear systems are hard.

Linearisation

Linear systems are easy to solve, nonlinear systems are hard.



Linearisation: for each nonlinear monomial, replace all of its occurrences by a new variable.

Linearisation

Linear systems are easy to solve, nonlinear systems are hard.

↪ **Linearisation:** for each nonlinear monomial, replace all of its occurrences by a new variable.

Example.

$$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$$

$$f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$$

$$f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$$

$$f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$$

$$f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$$

$$f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$$



$$f_1 : y_2 + y_5 + x_1 + x_3 + x_4 = 0$$

$$f_2 : y_4 + y_3 + y_6 + x_1 + x_2 + x_4 = 0$$

$$f_3 : y_5 + y_6 + x_1 + x_3 + 1 = 0$$

$$f_4 : y_1 + y_2 + y_4 + x_3 + x_4 + 1 = 0$$

$$f_5 : y_1 + y_4 + y_3 + x_3 = 0$$

$$f_6 : y_2 + y_3 + y_6 + x_1 + x_2 + x_3 + x_4 = 0$$

Linearisation

Linear systems are easy to solve, nonlinear systems are hard.

↪ **Linearisation:** for each nonlinear monomial, replace all of its occurrences by a new variable.

Example.

$$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$$

$$f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$$

$$f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$$

$$f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$$

$$f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$$

$$f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$$



$$f_1 : y_2 + y_5 + x_1 + x_3 + x_4 = 0$$

$$f_2 : y_4 + y_3 + y_6 + x_1 + x_2 + x_4 = 0$$


$$f_3 : y_5 + y_6 + x_1 + x_3 + 1 = 0$$

$$f_4 : y_1 + y_2 + y_4 + x_3 + x_4 + 1 = 0$$


$$f_5 : y_1 + y_4 + y_3 + x_3 = 0$$

$$f_6 : y_2 + y_3 + y_6 + x_1 + x_2 + x_3 + x_4 = 0$$


Linearisation



Linearisation adds solutions: a *random* quadratic system of m equations in n variables, when $n = m$, is expected to have one solution (probability is $\sim \frac{1}{q}$ for systems over \mathbb{F}_q). The corresponding linearised system has a solution space of dimension $\binom{n+1}{2} - m$.

 $\binom{n}{2}$ quadratic plus n linear monomials

Linearisation



Linearisation adds solutions: a *random* quadratic system of m equations in n variables, when $n = m$, is expected to have one solution (probability is $\sim \frac{1}{q}$ for systems over \mathbb{F}_q). The corresponding linearised system has a solution space of dimension $\binom{n+1}{2} - m$.

 $\binom{n}{2}$ quadratic plus n linear monomials



Loss of information: e.g. assignment $x_1 = 1; x_2 = 0; y_1 = 1$; is part of a valid solution to the linearised system, but $x_1 x_2 \neq y_1$.

Macaulay matrix

Equations
↓

	Monomials →										
	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1
f_1											
f_2											
f_3											
f_4											
f_5											
f_6											

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

Macaulay matrix

Equations
↓

	Monomials →										
	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1
f_1	0	1	0	1	0	1	0	0	1	1	0
f_2	0	0	1	1	1	0	1	1	0	1	0
f_3	0	0	0	1	0	1	0	1	1	0	1
f_4	1	1	0	1	1	0	0	0	1	1	1
f_5	1	0	1	1	1	0	0	0	1	0	0
f_6	0	1	1	1	0	0	1	1	1	1	0

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$




Simple algorithm

[Bouillaguet, Delaplace, T., 2021]

Simple algorithm

- Partial assignment
- Gaussian elimination




$$1 \cdot 0 + 1 \cdot x_3 + x_3 \cdot x_4 + x_3 = 0$$

$$0 \cdot x_3 + 0 \cdot x_4 + 1 + 0 + 1 = 0$$

$$1 \cdot 0 + 0 \cdot x_3 + 0 \cdot x_4 + 1 + x_4 = 0$$

$$1 \cdot x_4 + 0 \cdot x_3 + 0 + x_3 + x_4 = 0$$

Simple algorithm



Guess sufficiently many variables so that the remaining polynomial system can be solved by linearization.

Simple algorithm: complexity

Simple algorithm: complexity

- n - number of variables
- m - number of equations

Simple algorithm: complexity

- n - number of variables
- m - number of equations

 Enumeration ends when:

number of **monomials** \leq number of **equations**

Simple algorithm: complexity

- n - number of variables
- m - number of equations

 Enumeration ends when:

number of **monomials** \leq number of **equations**

$$\binom{n-?}{2} \leq m$$


Simple algorithm: complexity

- n - number of variables
- m - number of equations

 Enumeration ends when:

number of **monomials** \leq number of **equations**

$$\binom{n-?}{2} \leq m$$

 $\mathcal{O}(2^{n-\sqrt{2m}})$


Simple algorithm: complexity

- n - number of variables
- m - number of equations

 Enumeration ends when:


number of **monomials** \leq number of **equations**

$$\binom{n-?}{2} \leq m$$

 $\mathcal{O}(2^{n-\sqrt{2m}})$

 See also: Quantum BDT [Edme, Fouque, Schrottenloher]

Overview of solvers





Gröbner basis algorithms

[Buchberger, 1965]

[Lazard, 1983]

F_4/F_5 [Faugère, 1999/2002]

(XL [Courtois, Klimov, Patarin, Shamir, 2000])

Gröbner basis algorithms (intuition)

*We are essentially describing the XL algorithm.

Gröbner basis algorithms (intuition)

*We are essentially describing the XL algorithm.

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1
f_1	0	1	0	1	0	1	0	0	1	1	0
f_2	0	0	1	1	1	0	1	1	0	1	0
f_3	0	0	0	1	0	1	0	1	1	0	1
f_4	1	1	0	1	1	0	0	0	1	1	1
f_5	1	0	1	1	1	0	0	0	1	0	0
f_6	0	1	1	1	0	0	1	1	1	1	0

Gröbner basis algorithms (intuition)

*We are essentially describing the XL algorithm.

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1
f_1	0	1	0	1	0	1	0	0	1	1	0
f_2	0	0	1	1	1	0	1	1	0	1	0
f_3	0	0	0	1	0	1	0	1	1	0	1
f_4	1	1	0	1	1	0	0	0	1	1	1
f_5	1	0	1	1	1	0	0	0	1	0	0
f_6	0	1	1	1	0	0	1	1	1	1	0

Gröbner basis algorithms (intuition)

*We are essentially describing the XL algorithm.

$D = 3$

$$\begin{aligned} f_1 &: x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0 \\ f_2 &: x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0 \\ f_3 &: x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0 \\ f_4 &: x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0 \\ f_5 &: x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0 \\ f_6 &: x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0 \end{aligned}$$

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1	$x_1x_2x_3$	$x_1x_2x_4$	$x_1x_3x_4$	$x_2x_3x_4$
f_1	0	1	0	1	0	1	0	0	1	1	0				
f_2	0	0	1	1	1	0	1	1	0	1	0				
f_3	0	0	0	1	0	1	0	1	1	0	1				
f_4	1	1	0	1	1	0	0	0	1	1	1				
f_5	1	0	1	1	1	0	0	0	1	0	0				
f_6	0	1	1	1	0	0	1	1	1	1	0				
x_1f_1															
x_2f_1															
...															

Gröbner basis algorithms (intuition)

*We are essentially describing the XL algorithm.

$D = 4$

$$\begin{aligned} f_1 &: x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0 \\ f_2 &: x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0 \\ f_3 &: x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0 \\ f_4 &: x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0 \\ f_5 &: x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0 \\ f_6 &: x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0 \end{aligned}$$

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1	$x_1x_2x_3$	$x_1x_2x_4$	$x_1x_3x_4$	$x_2x_3x_4$	$x_1x_2x_3x_4$
f_1	0	1	0	1	0	1	0	0	1	1	0					
f_2	0	0	1	1	1	0	1	1	0	1	0					
f_3	0	0	0	1	0	1	0	1	1	0	1					
f_4	1	1	0	1	1	0	0	0	1	1	1					
f_5	1	0	1	1	1	0	0	0	1	0	0					
f_6	0	1	1	1	0	0	1	1	1	1	0					
x_1f_1																
x_2f_1																
...																
$x_1x_2f_1$																
$x_1x_3f_1$																

Gröbner basis

Gröbner basis

- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.

Gröbner basis

- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.
- An **ideal** in R is an additive subgroup I such that if $g \in R$ and $f \in I$, then $gf \in I$.

Gröbner basis

- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.
- An **ideal** in R is an additive subgroup I such that if $g \in R$ and $f \in I$, then $gf \in I$.
- The subset $\{f_1, \dots, f_m\} \subset R$ is a **set of generators** for an ideal I if every element $t \in I$ can be written in the form
$$t = \sum_{i=1}^n \text{with } g_i \in R.$$

Gröbner basis

- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.
- An **ideal** in R is an additive subgroup I such that if $g \in R$ and $f \in I$, then $gf \in I$.
- The subset $\{f_1, \dots, f_m\} \subset R$ is a **set of generators** for an ideal I if every element $t \in I$ can be written in the form
$$t = \sum_{i=1}^n \text{with } g_i \in R.$$
- By the **Hilbert basis theorem**: every ideal in R has a **finite** set of generators.

Gröbner basis

- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.
- An **ideal** in R is an additive subgroup I such that if $g \in R$ and $f \in I$, then $gf \in I$.
- The subset $\{f_1, \dots, f_m\} \subset R$ is a **set of generators** for an ideal I if every element $t \in I$ can be written in the form
$$t = \sum_{i=1}^n g_i$$
with $g_i \in R$.
- By the **Hilbert basis theorem**: every ideal in R has a **finite** set of generators.
- The subset of R defined as $V(I) = \{(a_1, \dots, a_n) \in \mathbb{F}_q^n \mid f(a_1, \dots, a_n) = 0 \text{ for all } f \in I\}$ is called an **algebraic variety**. It is the set of all solutions to the system of equations
$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0.$$

Gröbner basis


- Let $R = \mathbb{F}_q[x_1, \dots, x_n]$ be the **polynomial ring** in n variables.
- An **ideal** in R is an additive subgroup I such that if $g \in R$ and $f \in I$, then $gf \in I$.
- The subset $\{f_1, \dots, f_m\} \subset R$ is a **set of generators** for an ideal I if every element $t \in I$ can be written in the form
$$t = \sum_{i=1}^n g_i$$
with $g_i \in R$.
- By the **Hilbert basis theorem**: every ideal in R has a **finite** set of generators.
- The subset of R defined as $V(I) = \{(a_1, \dots, a_n) \in \mathbb{F}_q^n \mid f(a_1, \dots, a_n) = 0 \text{ for all } f \in I\}$ is called an **algebraic variety**. It is the set of all solutions to the system of equations
$$f_1(x_1, \dots, x_n) = \dots = f_m(x_1, \dots, x_n) = 0.$$
- By the **Nullstellensatz**: $\mathbf{I}(V(I)) = I$, where $\mathbf{I}(V)$ denotes the ideal of V , i.e. $\mathbf{I}(V) = \{f \in R \mid f(a) = 0 \text{ for all } a \in V\}$ (Similar to Gauss' fundamental theorem, but for polynomials in many variables).

Gröbner basis

- A **Gröbner basis** of an ideal I is a set of generators with some **nice** (useful) property.

Gröbner basis


- A **Gröbner basis** of an ideal I is a set of generators with some **nice** (useful) property.



For our case, the nice property is that a solution can be extracted easily from the Gröbner basis.

Gröbner basis

- A **Gröbner basis** of an ideal I is a set of generators with some **nice** (useful) property.



For our case, the nice property is that a solution can be extracted easily from the Gröbner basis.

Example. The **shape** of a GB with respect to the lexicographic order

$$f_1 : x_1x_3 + x_1 + x_2x_4 + x_5 + x_6 + 1 = 0$$

$$f_2 : x_1x_4 + x_1 + x_2x_3 + x_2 + x_3x_4 + x_3x_6 + x_4 + x_5 = 0$$

$$f_3 : x_1x_5 + x_1 + x_2 + x_3x_4 + x_6 + 1 = 0$$


$$f_4 : x_1x_2 + x_1x_3 + x_2x_5 + x_3 + x_4 + x_6 + 1 = 0$$

$$f_5 : x_1x_4 + x_2x_3 + x_2x_5 + x_5x_6 + 1 = 0$$

$$f_6 : x_1x_3 + x_1x_4 + x_1 + x_2 + x_3x_6 + x_3 + x_5 = 0$$

Gröbner basis

- A **Gröbner basis** of an ideal I is a set of generators with some **nice** (useful) property.



For our case, the nice property is that a solution can be extracted easily from the Gröbner basis.


Example. The **shape** of a GB with respect to the lexicographic order

$$\begin{aligned}f_1 &: x_1x_3 + x_1 + x_2x_4 + x_5 + x_6 + 1 = 0 \\f_2 &: x_1x_4 + x_1 + x_2x_3 + x_2 + x_3x_4 + x_3x_6 + x_4 + x_5 = 0 \\f_3 &: x_1x_5 + x_1 + x_2 + x_3x_4 + x_6 + 1 = 0 \\f_4 &: x_1x_2 + x_1x_3 + x_2x_5 + x_3 + x_4 + x_6 + 1 = 0 \\f_5 &: x_1x_4 + x_2x_3 + x_2x_5 + x_5x_6 + 1 = 0 \\f_6 &: x_1x_3 + x_1x_4 + x_1 + x_2 + x_3x_6 + x_3 + x_5 = 0\end{aligned}$$




$$\begin{aligned}f'_1 &: x_1 + x_6 = 0 \\f'_2 &: x_2 + x_6 = 0 \\f'_3 &: x_3 + x_6 = 0 \\f'_4 &: x_4 + x_6 + 1 = 0 \\f'_5 &: x_5 = 0\end{aligned}$$

**
*



Gröbner basis

- A **Gröbner basis** of an ideal I is a set of generators with some **nice** (useful) property.




For our case, the nice property is that a solution can be extracted easily from the Gröbner basis.

Example. The **shape** of a GB with respect to the lexicographic order

$$\begin{aligned} f_1 &: x_1x_3 + x_1 + x_2x_4 + x_5 + x_6 + 1 = 0 \\ f_2 &: x_1x_4 + x_1 + x_2x_3 + x_2 + x_3x_4 + x_3x_6 + x_4 + x_5 = 0 \\ f_3 &: x_1x_5 + x_1 + x_2 + x_3x_4 + x_6 + 1 = 0 \\ f_4 &: x_1x_2 + x_1x_3 + x_2x_5 + x_3 + x_4 + x_6 + 1 = 0 \\ f_5 &: x_1x_4 + x_2x_3 + x_2x_5 + x_5x_6 + 1 = 0 \\ f_6 &: x_1x_3 + x_1x_4 + x_1 + x_2 + x_3x_6 + x_3 + x_5 = 0 \end{aligned}$$



$$\begin{aligned} f'_1 &: x_1 + x_6 = 0 \\ f'_2 &: x_2 + x_6 = 0 \\ f'_3 &: x_3 + x_6 = 0 \\ f'_4 &: x_4 + x_6 + 1 = 0 \\ f'_5 &: x_5 = 0 \end{aligned}$$

**
* 

$$V(\langle f_1, \dots, f_6 \rangle) = \{ (0,0,0,1,0,0), (1,1,1,0,0,1) \}$$

Gröbner basis algorithms:

Buchberger, Lazard, F4, F5



Follow the core idea that we described, but combine the equations in an organised way, rather than multiplying them by all possible monomials.

Not covered in this talk:

- Monomial orders
- S-polynomials
- Polynomial long division
- Row reduction in parallel
- Reductions to zero
- Syzygy criterion
- ...

XL / Gröbner basis algorithms: complexity

XL / Gröbner basis algorithms: complexity

$$\mathcal{O} \left(m D_{reg} \binom{n + D_{reg} - 1}{D_{reg}}^{\omega} \right)$$

XL / Gröbner basis algorithms: complexity

$$\mathcal{O} \left(m D_{reg} \binom{n + D_{reg} - 1}{D_{reg}}^{\omega} \right)$$

D_{reg} : degree of regularity



the power of the first non-positive coefficient in the expansion of

$$\frac{(1 - t^2)^m}{(1 - t)^n}$$

XL / Gröbner basis algorithms: complexity

```
▷ m=8
n=7
R.<t> = PowerSeriesRing(ZZ)
hs = ((1-t^2)^m) / (1-t)^n
print(hs)
```


[3] ✓ 0.0s

... 1 + 7*t + 20*t^2 + 28*t^3 + 14*t^4 - 14*t^5 - 28*t^6 - 20*t^7 - 7*t^8 - t^9 + 0(t^20)

→ The number of monomials (columns) **minus** linearly independent equations (rows) at **degree $D = 4$** is 14.

XL / Gröbner basis algorithms: complexity


D_{reg} : degree of regularity



the power of the first non-positive coefficient in the expansion of


$$\frac{(1 - t^2)^m}{(1 - t)^n}$$

```
▷ m=8
  n=7
  R.<t> = PowerSeriesRing(ZZ)
  hs = ((1-t^2)^(m)) / (1-t)^(n)
  print(hs)
[3] ✓ 0.0s
... 1 + 7*t + 20*t^2 + 28*t^3 + 14*t^4 - 14*t^5 - 28*t^6 - 20*t^7 - 7*t^8 - t^9 + 0(t^20)
```



The number of monomials (columns) **minus** linearly independent equations (rows) at **degree $D = 4$** is 14.

Overview of solvers





FXL

[Courtois, Klimov, Patarin, Shamir, 2000]


Hybrid

[Bettale, Faugère, Perret, 2009]

BoolSolve

[Bardet, Faugère, Salvy, Spaenlehauer, 2013]

FXL, Hybrid, BoolSolve



Techniques are already covered in the previous section.

Algorithms will be explained in the summary.



The crossbred algorithm

[Joux, Vitse, 2017]

Crossbred algorithm

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1
f_1	0	1	0	1	0	1	0	0	1	1	0
f_2	0	0	1	1	1	0	1	1	0	1	0
f_3	0	0	0	1	0	1	0	1	1	0	1
f_4	1	1	0	1	1	0	0	0	1	1	1
f_5	1	0	1	1	1	0	0	0	1	0	0
f_6	0	1	1	1	0	0	1	1	1	1	0

Crossbred algorithm

→ Put matrix in reduced row echelon form

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1

...

Crossbred algorithm

→ Take linear subsystem

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1

...



...if we had another 4 equations

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

Crossbred algorithm

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1
...											

Crossbred algorithm

- Subsystem is linear in variables $\{x_1, x_2, x_3\}$.
- Enumerating x_4 will result in a linear subsystem.

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1
...											

Crossbred algorithm

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1
...											

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

Crossbred algorithm

→ Subsystem can be linearised

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1
...											

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

Crossbred algorithm

→ Subsystem can be linearised

	x_1x_2	x_1x_3	x_2x_3	x_1x_4	x_2x_4	x_3x_4	x_1	x_2	x_3	x_4	1
f_1	1	0	0	0	0	0	0	0	0	1	1
f_2	0	1	0	0	0	0	1	1	1	1	0
f_3	0	0	1	0	0	0	1	1	0	1	0
f_4	0	0	0	1	0	0	1	1	1	0	1
f_5	0	0	0	0	1	0	0	1	0	0	0
f_6	0	0	0	0	0	1	1	1	1	0	1
...											

$f_1 : x_1x_3 + x_2x_4 + x_1 + x_3 + x_4 = 0$ $f_2 : x_2x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_4 = 0$ $f_3 : x_2x_4 + x_3x_4 + x_1 + x_3 + 1 = 0$ $f_4 : x_1x_2 + x_1x_3 + x_2x_3 + x_3 + x_4 + 1 = 0$ $f_5 : x_1x_2 + x_2x_3 + x_1x_4 + x_3 = 0$ $f_6 : x_1x_3 + x_1x_4 + x_3x_4 + x_1 + x_2 + x_3 + x_4 = 0$

...if we had another 4 equations, the subsystem would have a unique solution.

Otherwise: check candidate solutions against the other equations.

Crossbred algorithm


Parameters of the algorithm: D, k, d, h

- Enumerate h variables.
- Choose k of the remaining variables.
- Augment system up to degree D (compute degree- D Macaulay matrix).
- Take the subsystem that is at most degree d in the k chosen variables.
- Enumerate all but the k chosen variables.
- Linearise the subsystem and solve it.
- Check if candidate solutions are consistent with the rest of the system.

Crossbred algorithm

Parameters of the algorithm: D, k, d, h

- Enumerate h variables.
- Choose k of the remaining variables.
- Augment system up to degree D (compute degree- D Macaulay matrix).
- Take the subsystem that is at most degree d in the k chosen variables.
- Enumerate all but the k chosen variables.
- Linearise the subsystem and solve it.
- Check if candidate solutions are consistent with the rest of the system.


 The complexity is calculated as the best trade-off between the four parameters.

Crossbred algorithm

	Number of Variables (n)	Seed (0,1,2,3,4)	Date	Contestants	Computational Resource	Data
1	83	0	2023/09/16	Charles Bouillaguet and Julia Sauvage	https://gitlab.lip6.fr/almasty/hpXbred , 3488 AMD EPYC 7J13 cores on the Oracle public cloud	Details
6	74	0	2016/12/17	Antoine Joux	New hybridized XL related algorithm, Heterogeneous cluster of Intel Xeon @ 2.7-3.5 Ghz	Details
7	74	4	2017/11/15	Kai-Chun Ning, Ruben Niederhagen	Parallel Crossbred, 54 GPUs in the Saber cluster	Details
25	66	0	2016/01/22	Tung Chou, Ruben Niederhagen, Bo-Yin Yang	Gray Code enumeration, Rivyera, 128 Spartan 6 FPGAs	Details

Fukuoka MQ challenge record computations ($m = 2n$)

Overview of solvers



Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES

Simple

FXL

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary


(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis



FES

KL

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES

Simple

FXL

F_4 / F_5


SAT solvers

Crossbred

BoolSolve

Hybrid

Summary



Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES

Simple

FXL

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration


Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES



F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES

Simple

FXL

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration

Candidate
solutions

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

	x_1x_2	x_1x_3	x_1x_4	x_1	x_2x_3	x_2x_4	x_2	x_3x_4	x_3	x_4	1	$x_1x_2x_3$	$x_1x_2x_4$	$x_1x_3x_4$	$x_2x_3x_4$	$x_1x_2x_3x_4$
f_1	0	1	0	1	0	1	0	0	1	1	0					
f_2	0	0	1	1	1	0	1	1	0	1	0					
f_3	0	0	0	1	0	1	0	1	1	0	1					
f_4	1	1	0	1	1	0	0	0	1	1	1					
f_5	1	0	1	1	1	0	0	0	1	0	0					
f_6	0	1	1	1	0	0	1	1	1	1	0					
x_1f_1																
x_2f_1																
...																
$x_1x_2f_1$																
$x_1x_3f_1$																

FES

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

FES

Simple

FXL

F_4 / F_5

SAT solvers

Crossbred

BoolSolve

Hybrid

Summary

(Partial)
enumeration

Candidate
solutions
(subsystem)

Conflict search

Extending to
higher degrees

Computing a
Gröbner Basis

$$\begin{aligned}f'_1 &: x_1 + x_6 = 0 \\f'_2 &: x_2 + x_6 = 0 \\f'_3 &: x_3 + x_6 = 0 \\f'_4 &: x_4 + x_6 + 1 = 0 \\f'_5 &: x_5 = 0\end{aligned}$$

**
*



FES

Simple

FXL

F_4 / F_5


SAT solvers

Crossbred


BoolSolve

Hybrid


Summary




Summary




Summary




Summary




Summary




Summary



Summary



Summary



Recap

- ▶ The **MQ problem** is (usually) hard.
- ▶ **Modelisation** can be crucial to how efficient an attack is.
- ▶ We have a variety of **solvers** for (over)determined systems.
- ▶ We can estimate the **complexity** of solving random systems, but for structured systems this requires deeper analysis.

To implement a solver and practice modelisation of different attacks:

Tutorial **Tuesday, July 1**

(install SageMath beforehand: <https://github.com/LarsMath/tutorial-algebraic-cryptanalysis>)

└─ joint with Lars Ran