

PROBLEM #1.

Given the function $f(x)$ find the derivative.

$$f(x) = \frac{x^6}{6} - 3x^4 - \frac{2x^3}{3} + \frac{27x^2}{2} + 18x - 30$$

$$\begin{aligned} f'(x) &= \frac{6x^5}{6} - 3(4)x^3 - 2\frac{(3)}{3}x^2 + \frac{(2)(27)}{2}x + 18 \\ &= x^5 - 12x^3 - 2x^2 + 27x + 18 \end{aligned}$$



PROBLEM #2.

Find the first estimate/iteration of the Newton-Raphson.

$$x_0 = 1.$$

$$x_{i+1} = x_0 - \frac{f(x_0)}{f'(x_0)},$$

$$x_1 = 1 - \frac{\left[\frac{(1)^6}{6} - 3(1)^4 - \frac{2(1)^3}{3} + \frac{27(1)^2}{2} + 18(1) - 30 \right]}{(1)^5 - 12(1)^3 - 2(1)^2 + 27 + 18}$$

$$= 1 - \frac{\left[\frac{1}{6} - 3 - \frac{2}{3} + \frac{27}{2} + 18 - 30 \right]}{1 - 12 - 2 + 27 + 18}$$

$$= 1 - \frac{\left[\frac{1}{6} - \frac{2}{3} - \frac{3}{2} \right]}{32}$$

$$= 1 + \frac{1}{16}$$

PROBLEM #3

Perform Newton-Raphson to find the root around $x = -4$.

```
1 def f(x):
2     return x**6/6 - 3*x**4 - 2*x**3/3 + 27*x**2/2 + 18*x - 30
3
4 def d_f(x):
5     return x**5 - 12*x**3 - 2*x**2 + 27*x + 18 # Complete this line with
6         the derivative you have calculated.
7
8 x = -4.0
9
10 d = {"x": [x], "f(x)": [f(x)]}
11 for i in range(0, 20):
12     x = x - f(x) / d_f(x)
13     d["x"].append(x)
14     d["f(x)"].append(f(x))
15
16 pd.DataFrame(d, columns=['x', 'f(x)'])
```

Run
Reset

| | x | f(x) |
|----|-----------|--------------|
| 0 | -4.000000 | 7.133333e+01 |
| 1 | -3.811287 | 1.223161e+01 |
| 2 | -3.763093 | 6.515564e-01 |
| 3 | -3.760224 | 2.198858e-03 |
| 4 | -3.760214 | 2.531156e-08 |
| 5 | -3.760214 | 1.421085e-13 |
| 6 | -3.760214 | 4.263256e-14 |
| 7 | -3.760214 | 4.263256e-14 |
| 8 | -3.760214 | 4.263256e-14 |
| 9 | -3.760214 | 4.263256e-14 |
| 10 | -3.760214 | 4.263256e-14 |
| 11 | -3.760214 | 4.263256e-14 |
| 12 | -3.760214 | 4.263256e-14 |
| 13 | -3.760214 | 4.263256e-14 |
| 14 | -3.760214 | 4.263256e-14 |
| 15 | -3.760214 | 4.263256e-14 |
| 16 | -3.760214 | 4.263256e-14 |
| 17 | -3.760214 | 4.263256e-14 |
| 18 | -3.760214 | 4.263256e-14 |
| 19 | -3.760214 | 4.263256e-14 |
| 20 | -3.760214 | 4.263256e-14 |

PROBLEM #4.

What would happen if Newton-Raphson was computed for $f(x)$ around point $x_0 = 1.99$.

- ☒ method converges to root nearest $x = -4$.
- ☐ method does not converge, infinitely oscillates
- ☐ none of the other statements are true.
- ☒ method takes over 15 iterations to converge
- ☐ method diverges to infinity
- ☐ method converges to the root nearest $x = 1$.

PROBLEM #5

There are points which oscillate and do not converge - like $x = 3.1$.