


1. The size of a vector is the root of the squared sum of all its components



$$|r| = \sqrt{|r|_1^2 + |r|_2^2}$$

$$= \sqrt{\sum_i |r|_i^2}$$

, more generally this distance (euclidean) holds across any number of dimensions d.

What is the size of
 $v = [1 \ 3 \ 4 \ 2]^T$

$$|v| = \sqrt{(1)^2 + (3)^2 + (4)^2 + (2)^2}$$

$$= \sqrt{30}$$

PROBLEM #2

What is $r = [-5 \ 3 \ 2 \ 8]^T$ dotted with $s = [1 \ 2 \ -1 \ 0]^T$

$$r \cdot s = -5(1) + 3(2) + 2(-1) + 8(0)$$

$$= -1$$

PROBLEM #3

Scalar projection can happen in any dimension

What is the scalar proj. of $r = [3 \ -4 \ 0]^T$ onto $s = [10 \ 5 \ -6]^T$


$$\text{scalar proj: } \frac{r \cdot s}{|r|} = |s| \cos \theta$$

$$r \cdot s = 3(10) + (-4)(5) + 0(-6)$$

$$|r| = \sqrt{3^2 + (-4)^2 + (0)^2}$$


$$\frac{r \cdot s}{|r|} = \frac{10}{5}$$

$$= 2$$



"Want to find the size of the blue vector".
 - scalar proj.

PROBLEM #4.



The vector proj.
is the blue vector.

what is the vector projection of $r = [3 \ -4 \ 0]^T$
 $s = [10 \ 5 \ -6]^T$?

$$\text{vector proj.} = \frac{r}{|r|} \times \text{scalar proj.}$$

$$r \cdot s = 3(10) + (-4)(5) + (0)(-6) \\ = 10$$

$$r \cdot r = (3)^2 + (-4)^2 + (0)^2 \\ = 25$$

$$r \frac{r \cdot s}{r \cdot r} = \begin{bmatrix} 3 \\ -4 \\ 0 \end{bmatrix} \frac{10}{25} \\ = \begin{bmatrix} 6/5 \\ -8/5 \\ 0 \end{bmatrix}$$

$$= \frac{r}{|r|} \times \frac{r \cdot s}{|r|}, \text{ or } \frac{r}{|r|} \times |s| \cos \theta \\ = r \frac{r \cdot s}{r \cdot r}$$

PROBLEM #5.

Which is larger given $a = [3 \ 0 \ 4]^T$ and $b = [0 \ 5 \ 12]^T$
is $|a+b|$ or $|a|+|b|$?

Options:

(in this case).

I. $|a+b| < |a|+|b|$

II. $|a+b| = |a|+|b|$

III. $|a+b| > |a|+|b|$

$$|a| = \sqrt{(3)^2 + (0)^2 + (4)^2} = 5$$

$$|b| = \sqrt{(0)^2 + (5)^2 + (12)^2} = 13$$

$$a+b = [(3+0) \ (0+5) \ (4+12)]^T$$

$$|a+b| = \sqrt{3^2 + 5^2 + 16^2} = \sqrt{290}$$

Given the two vectors conjoined
have an angle between them
from $[0^\circ - 180^\circ]$, their addition
will either be synergistic or
otherwise.

Therefore $|a+b| \leq |a|+|b|$.

PROBLEM #6.

Which of the following statements about the dot-product are correct?

□ The dot product is not commutative

(i.e. $s \cdot r \neq r \cdot s$) $\Rightarrow \text{dot prod} = \sum_{i=1}^n |s_i| \times |r_i|$. multiplication is always commutative


□ The scalar projection of s -onto- r is always the same as r -onto- s .

$$\begin{aligned} \text{scalar proj of } r\text{-onto-}s &= \frac{s \cdot r}{|s|} = |r| \cos \theta \\ \text{scalar proj of } s\text{-onto-}r &= \frac{r \cdot s}{|r|} = |s| \cos \theta \end{aligned}$$

Both equations share $r \cdot s$, which is commutative. But $|s|$ is not always equal to $|r|$.

Therefore the scalar projections are not always equal.

For ex.



$$|x| \neq |y|$$

✓ The vector proj of s -onto- r is equal to scalar proj of s -onto- r multiplied by vector of unit length with direction of r .

$$\begin{aligned} \text{vector proj} &= \frac{r}{|r|} \cdot \text{scalar proj.} \\ &= \frac{r}{|r|} \times \frac{r \cdot s}{|r|} \\ &= r \frac{r \cdot s}{r \cdot r} \end{aligned}$$

given this formula the statement is true.

- ☑ We can find the angle between two vectors using the dot product.

$$\begin{aligned} \text{dot prod} &= \forall i \in d \quad d_i = s_i \times r_i \\ &= |r||s|\cos\theta, \text{ by equating the dot product} \\ &\quad \text{and cosine rule (using vectors)} \\ &\quad \text{and simplifying.} \end{aligned}$$

to get the angle between two vectors can
dot prod = x , some val

$$\begin{aligned} r \cdot s = x &\iff x = |r||s|\cos\theta \\ &\iff \cos\theta = x \times \frac{1}{|r||s|} \\ &\quad \theta = \cos^{-1}\left(\frac{x}{|r||s|}\right) \end{aligned}$$

Thus, you could find the angle using the dot prod and moduli of participating vectors.

- ☑ The size of the vector is equal to the square root of the dot product of the vector with itself.

Assume the converse,

$$|r| \neq \sqrt{r \cdot r}.$$

$$|r| = \sqrt{\sum_i |r_i|^2}$$

$$\begin{aligned} \sqrt{r \cdot r} &= \sqrt{|r||r|\cos\theta} \\ &= \sqrt{|r||r|\cos(0)} \\ &= \sqrt{|r||r|(1)} \\ &= \sqrt{|r|^2} \\ &= |r|, \end{aligned}$$

Therefore $|r| \neq \sqrt{r \cdot r}$ is false and the hypothesis must be true!