

⇒ Goal to change from standard basis to orthogonal basis.

PROBLEM #1.

What is v in the basis of b_1 and b_2 , given

$$v = [5 \ -1]^T, \quad b_1 = [1 \ 1], \quad b_2 = [1 \ -1]^T.$$

1. Check if basis vectors are orthogonal,

$$b_1 \cdot b_2 = |b_1| |b_2| \cos \theta$$

$$\frac{b_1 \cdot b_2}{|b_1| |b_2|} = \cos \theta$$

$$|b_1| |b_2|$$

If the basis vectors are orthogonal they will be 90° apart from one another.

$$0 = \cos 90$$

$$= \frac{(1)(1) + (1)(-1)}{\sqrt{2}\sqrt{2}}$$

$$= 0/2.$$

Thus b_1 and b_2 are orthogonal.

2. Translate vector's bases,

Project v onto b_1 ,

$$\frac{b_1}{|b_1|} \times \frac{v \cdot b_1}{|b_1|} = v_{b_1}$$

$$= \begin{bmatrix} 1 \\ 1 \end{bmatrix} \times \left(\frac{1}{\sqrt{2}} \right)^2 \times ((5)(1) + (-1)(1))$$

$$= 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

The scalar projection of v onto b_1 is 2.

Project v onto b_2 ,

$$v_{b_2} = \frac{b_2}{|b_2|} \times \frac{v \cdot b_2}{|b_2|}$$

$$= \begin{bmatrix} 1 \\ -1 \end{bmatrix} \times \left(\frac{1}{\sqrt{2}} \right)^2 \times ((5)(1) + (-1)(-1))$$

$$= \begin{bmatrix} 1 \\ -1 \end{bmatrix} \frac{6}{2}$$

The scalar projection of v onto b_2 is 3.

$$v_b = \begin{bmatrix} v_{b_1} \\ v_{b_2} \end{bmatrix} = [2 \ 3]^T.$$

PROBLEM #2

What is v defined in the basis b_1 and b_2 .

Given $v = \begin{bmatrix} 10 \\ -5 \end{bmatrix}$, $b_1 = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$, $b_2 = \begin{bmatrix} 4 \\ -3 \end{bmatrix}$.

Project v onto b_1 .

$$\begin{aligned} v_{b_1} &= \frac{b_1}{|b_1|} \times \frac{b_1 \cdot v}{|b_1|} \\ &= \begin{bmatrix} 3 \\ 4 \end{bmatrix} \times \frac{(10)(3) + (-5)(4)}{(\sqrt{3^2 + 4^2})^2} \\ &= \frac{10}{25} [3 \ 4]^T. \end{aligned}$$

$$\begin{aligned} v &= \begin{bmatrix} v_{b_1} \\ v_{b_2} \end{bmatrix} \\ &= \begin{bmatrix} 2/5 \\ 11/5 \end{bmatrix}. \end{aligned}$$

Project v onto b_2 .

$$\begin{aligned} v_{b_2} &= \frac{b_2}{|b_2|} \times \frac{b_2 \cdot v}{|b_2|} \\ &= \begin{bmatrix} 4 \\ -3 \end{bmatrix} \times \frac{(4)(10) + (-3)(-5)}{(\sqrt{4^2 + (-3)^2})^2} \\ &= \frac{55}{25} \begin{bmatrix} 4 \\ -3 \end{bmatrix}. \end{aligned}$$

PROBLEM #3.

Where $v = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$, $b_1 = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$
 $b_2 = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$.

$$\begin{aligned} v_b &= \begin{bmatrix} \frac{b_1}{|b_1|} \times \frac{b_1 \cdot v}{|b_1|} \\ \frac{b_2}{|b_2|} \times \frac{b_2 \cdot v}{|b_2|} \end{bmatrix} \\ &= \begin{bmatrix} \begin{bmatrix} -3 \\ 1 \end{bmatrix} \times \frac{(2)(-3) + (2)(1)}{(\sqrt{(-3)^2 + 1^2})^2} \\ \begin{bmatrix} 1 \\ 3 \end{bmatrix} \times \frac{(1)(2) + (3)(2)}{(\sqrt{1^2 + 3^2})^2} \end{bmatrix} \end{aligned}$$

$$= \left[-\frac{4}{10} [-3 \ 1]^T \quad \frac{8}{10} [1 \ 3]^T \right] \rightarrow \begin{bmatrix} -2/5 & 4/5 \end{bmatrix}^T$$

PROBLEM #4.

$$v_{b1} = \frac{(1)(2) + (1)(1) + (1)(0)}{(\sqrt{2^2 + 1^2 + 0^2})^2} = \frac{3}{5}$$

$$\begin{aligned} v_{b2} &= \frac{(1)(1) + (1)(-2) + (1)(-1)}{(\sqrt{(1)^2 + (-2)^2 + (-1)^2})^2} \\ &= \frac{-2}{6} \end{aligned}$$

$$\begin{aligned} v_{b3} &= \frac{(1)(-1) + (1)(2) + (1)(-5)}{(\sqrt{(-1)^2 + (2)^2 + (-5)^2})^2} \\ &= \frac{-4}{30} \\ &= \frac{-2}{15} \end{aligned}$$

PROBLEM #5.

$$v_b = [1 \ 0 \ 1 \ 1]^T$$