



A cool thing happens when you differentiate  $e^x$  - also equivalent to Maclaurin series

$$e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$$

$$\frac{d}{dx} e^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$$

The derivative of the infinitely long series of  $e^x$  is the very same series!



$$g(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

$$e^x = 1 + x + \frac{x^2}{2} \dots$$


$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

Similarly expand around  $x=0$ .

"If you know everything about a function at  $x=0$ , you can reconstruct everything about the function everywhere."

"Instead of expanding around  $x=0$ , to expand around  $x=p$ !"

## $e^x$ Taylor Series



The first order derivation at point  $p$ . of a Taylor Series yields the process by which you can produce the other derivation orders.

Approximate the first order - linear component - of  $e^x$ ,

$$y = mx + c$$

$$= f'(x)x + c, \text{ where } f'(x) \text{ is the first order deriv.}$$

In approximating around point  $p$ . the first order derivative shows that the gradient is relative to the point  $(x-p)$ ; that is  $p$ 's distance from  $x$ .

Calculation of  $c$ :

$$y = f'(p)p + c$$

$$= f'(p)p + c, \text{ where } p \text{ is the point we are deriving around}$$

$$\Leftrightarrow f(p) = f'(p)p + c$$

$$\Leftrightarrow c = f(p) - f'(p)p$$

Approximation functions around  $p$ .

$$g_0(x) = f(p)$$

$$g_1(x) = f(p) + f'(p)(x-p)$$

$$g_2(x) = f(p) + f'(p)(x-p) + \frac{f''(p)(x-p)^2}{2}, \text{ where } g_2(x) \text{ is the Maclaurin Series is:}$$

From which generalizes the Taylor Series.

$$g(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(p)}{n!} (x-p)^n$$

Brook Taylor (1685-1721)

Note if you sub in  $p=0$ , Taylor becomes Maclaurin.

$$g_2(x) = f(0) + f'(0)x + \frac{f''(0)x^2}{2}$$