


How to transform from one basis vectors to another.



DETERMINING THE PRE-IMAGE

From the e frame f 's basis vectors are $\begin{bmatrix} 3 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix}$

If the vector u_f in the basis of f , needs to be translated into the basis of e do,

$$u_e = \begin{bmatrix} 3 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3/2 \\ 1/2 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \text{ in basis of } e.$$

DETERMINING THE IMAGE.

To convert from a basis e to another basis f — given the translation matrix $T(x): x_f \rightarrow x_e$, the inverse of that translation will be the applicative matrix required to go from $e \rightarrow f$.

$$T^{-1}(x): x_e \rightarrow x_f$$

In this example,

$$T(x) = \begin{bmatrix} 3 & 1 \\ 1 & 1 \end{bmatrix}$$

$$T^{-1}(x) = \frac{1}{\det(T(x))} \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix}$$

This means that e 's basis will be,

$$T(\hat{e}_1): \hat{e}_1 \rightarrow \frac{1}{2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

$$T(\hat{e}_2): \hat{e}_2 \rightarrow \frac{1}{2} \begin{bmatrix} -1 \\ 3 \end{bmatrix}$$


TESTING, CONVERT u TO f -BASIS

$$u \text{ in } e = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$$

$$V(x): x_e \rightarrow x_f \\ = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix} x_e$$

$$u_f = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 3 \end{bmatrix} \begin{bmatrix} 3 \\ 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 3 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}_f. \quad \square$$

Translating to An Orthonormal Basis Vector Set.



TRANSFORMING FROM f -to- e USING f -basis in e

$$T(x): x_f \rightarrow x_e$$

$$= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

$$T\left(\begin{bmatrix} 2 \\ 1 \end{bmatrix}_f\right) = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \} \xrightarrow{\text{Aug}} u_e = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

e -to- f using inverse of f -basis in e .

$$A = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$$

$$A^{-1} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$$

$$A^{-1} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix} \} \xrightarrow{\text{Aug}} A^{-1} u_e = u_f$$

Since the vectors are orthogonal; can do changes of basis really easily with projections (that simplify to the dot-product).

① Get the basis vectors of f in terms of e .

$$\hat{f}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix} \text{ in } e$$

$$\hat{f}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}_f = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 \\ 1 \end{bmatrix} \text{ in } e$$

② Project the vector to translate u , onto the e -translated basis vectors of f .

$$\text{proj}_{u_e} \hat{f}_1 = \frac{\hat{f}_1}{\|\hat{f}_1\|} \frac{\hat{f}_{1e} \cdot u_e}{\|\hat{f}_{1e}\|} \rightarrow \frac{\begin{bmatrix} 1 \\ 0 \end{bmatrix}}{1} \times \frac{\hat{f}_{1e} \cdot u_e}{1} \rightarrow \begin{bmatrix} \hat{f}_{1e} \cdot u_e \\ 0 \end{bmatrix}$$

$$u_f = \text{proj}_{u_e} \hat{f}_1 + \text{proj}_{u_e} \hat{f}_2$$

Basically a vector is the sum of its components, and to figure out its components in the basis f , project the vector onto equivalent e basis representations then project the scalar proj in the direction of the basis of the f component basis vectors.