

# Class 08: Unsupervised Learning

Nathaniel Lightle (A16669288)

## Outline

Today we will apply the machine learning methods we introduced in the last class on breast cancer biopsy data from fine needle aspiration (FNA).

## Data input

The data is supplied in CSV format

## Exploratory Data analysis

## Preparing the data

```
wisc.df <- read.csv("WisconsinCancer.csv", row.names = 1)
head(wisc.df)
```

	diagnosis	radius_mean	texture_mean	perimeter_mean	area_mean
842302	M	17.99	10.38	122.80	1001.0
842517	M	20.57	17.77	132.90	1326.0
84300903	M	19.69	21.25	130.00	1203.0
84348301	M	11.42	20.38	77.58	386.1
84358402	M	20.29	14.34	135.10	1297.0
843786	M	12.45	15.70	82.57	477.1

	smoothness_mean	compactness_mean	concavity_mean	concave.points_mean
842302	0.11840	0.27760	0.3001	0.14710
842517	0.08474	0.07864	0.0869	0.07017
84300903	0.10960	0.15990	0.1974	0.12790
84348301	0.14250	0.28390	0.2414	0.10520

84358402	0.10030	0.13280	0.1980	0.10430	
843786	0.12780	0.17000	0.1578	0.08089	
	symmetry_mean	fractal_dimension_mean	radius_se	texture_se	perimeter_se
842302	0.2419	0.07871	1.0950	0.9053	8.589
842517	0.1812	0.05667	0.5435	0.7339	3.398
84300903	0.2069	0.05999	0.7456	0.7869	4.585
84348301	0.2597	0.09744	0.4956	1.1560	3.445
84358402	0.1809	0.05883	0.7572	0.7813	5.438
843786	0.2087	0.07613	0.3345	0.8902	2.217
	area_se	smoothness_se	compactness_se	concavity_se	concave.points_se
842302	153.40	0.006399	0.04904	0.05373	0.01587
842517	74.08	0.005225	0.01308	0.01860	0.01340
84300903	94.03	0.006150	0.04006	0.03832	0.02058
84348301	27.23	0.009110	0.07458	0.05661	0.01867
84358402	94.44	0.011490	0.02461	0.05688	0.01885
843786	27.19	0.007510	0.03345	0.03672	0.01137
	symmetry_se	fractal_dimension_se	radius_worst	texture_worst	
842302	0.03003	0.006193	25.38	17.33	
842517	0.01389	0.003532	24.99	23.41	
84300903	0.02250	0.004571	23.57	25.53	
84348301	0.05963	0.009208	14.91	26.50	
84358402	0.01756	0.005115	22.54	16.67	
843786	0.02165	0.005082	15.47	23.75	
	perimeter_worst	area_worst	smoothness_worst	compactness_worst	
842302	184.60	2019.0	0.1622	0.6656	
842517	158.80	1956.0	0.1238	0.1866	
84300903	152.50	1709.0	0.1444	0.4245	
84348301	98.87	567.7	0.2098	0.8663	
84358402	152.20	1575.0	0.1374	0.2050	
843786	103.40	741.6	0.1791	0.5249	
	concavity_worst	concave.points_worst	symmetry_worst		
842302	0.7119	0.2654	0.4601		
842517	0.2416	0.1860	0.2750		
84300903	0.4504	0.2430	0.3613		
84348301	0.6869	0.2575	0.6638		
84358402	0.4000	0.1625	0.2364		
843786	0.5355	0.1741	0.3985		
	fractal_dimension_worst				
842302	0.11890				
842517	0.08902				
84300903	0.08758				
84348301	0.17300				
84358402	0.07678				

843786

0.12440

Removing the diagnosis

```
wisc.data <- wisc.df[,-1]  
head(wisc.data)
```

	radius_mean	texture_mean	perimeter_mean	area_mean	smoothness_mean
842302	17.99	10.38	122.80	1001.0	0.11840
842517	20.57	17.77	132.90	1326.0	0.08474
84300903	19.69	21.25	130.00	1203.0	0.10960
84348301	11.42	20.38	77.58	386.1	0.14250
84358402	20.29	14.34	135.10	1297.0	0.10030
843786	12.45	15.70	82.57	477.1	0.12780
	compactness_mean	concavity_mean	concave.points_mean	symmetry_mean	
842302	0.27760	0.3001	0.14710	0.2419	
842517	0.07864	0.0869	0.07017	0.1812	
84300903	0.15990	0.1974	0.12790	0.2069	
84348301	0.28390	0.2414	0.10520	0.2597	
84358402	0.13280	0.1980	0.10430	0.1809	
843786	0.17000	0.1578	0.08089	0.2087	
	fractal_dimension_mean	radius_se	texture_se	perimeter_se	area_se
842302	0.07871	1.0950	0.9053	8.589	153.40
842517	0.05667	0.5435	0.7339	3.398	74.08
84300903	0.05999	0.7456	0.7869	4.585	94.03
84348301	0.09744	0.4956	1.1560	3.445	27.23
84358402	0.05883	0.7572	0.7813	5.438	94.44
843786	0.07613	0.3345	0.8902	2.217	27.19
	smoothness_se	compactness_se	concavity_se	concave.points_se	
842302	0.006399	0.04904	0.05373	0.01587	
842517	0.005225	0.01308	0.01860	0.01340	
84300903	0.006150	0.04006	0.03832	0.02058	
84348301	0.009110	0.07458	0.05661	0.01867	
84358402	0.011490	0.02461	0.05688	0.01885	
843786	0.007510	0.03345	0.03672	0.01137	
	symmetry_se	fractal_dimension_se	radius_worst	texture_worst	
842302	0.03003	0.006193	25.38	17.33	
842517	0.01389	0.003532	24.99	23.41	
84300903	0.02250	0.004571	23.57	25.53	
84348301	0.05963	0.009208	14.91	26.50	
84358402	0.01756	0.005115	22.54	16.67	
843786	0.02165	0.005082	15.47	23.75	

	perimeter_worst	area_worst	smoothness_worst	compactness_worst
842302	184.60	2019.0	0.1622	0.6656
842517	158.80	1956.0	0.1238	0.1866
84300903	152.50	1709.0	0.1444	0.4245
84348301	98.87	567.7	0.2098	0.8663
84358402	152.20	1575.0	0.1374	0.2050
843786	103.40	741.6	0.1791	0.5249

	concavity_worst	concave.points_worst	symmetry_worst
842302	0.7119	0.2654	0.4601
842517	0.2416	0.1860	0.2750
84300903	0.4504	0.2430	0.3613
84348301	0.6869	0.2575	0.6638
84358402	0.4000	0.1625	0.2364
843786	0.5355	0.1741	0.3985

	fractal_dimension_worst
842302	0.11890
842517	0.08902
84300903	0.08758
84348301	0.17300
84358402	0.07678
843786	0.12440

Making the diagnosis into a factor

```
diagnosis <- as.factor( wisc.df[,1])
head(diagnosis)
```

```
[1] M M M M M M
Levels: B M
```

## Exploratory Data Analysis

Q1. How many observations are in this dataset?

```
dim(wisc.data)
```

```
[1] 569 30
```

There are 569 observations in this dataset.

Q2. How many of the observations have a malignant diagnosis?

```
NROW(diagnosis[diagnosis == "M"])
```

```
[1] 212
```

There are 212 malignant diagnosis

Q3. How many variables/features in the data are suffixed with `_mean`?

```
length(grep("_mean", colnames(wisc.data)))
```

```
[1] 10
```

There are 10 variables/features in the data suffixed with `_mean`

## 2. Principal Component Analysis

### Performing PCA

Checking to see if scaling is necessary

```
colMeans(wisc.data)
```

radius_mean	texture_mean	perimeter_mean
1.412729e+01	1.928965e+01	9.196903e+01
area_mean	smoothness_mean	compactness_mean
6.548891e+02	9.636028e-02	1.043410e-01
concavity_mean	concave.points_mean	symmetry_mean
8.879932e-02	4.891915e-02	1.811619e-01
fractal_dimension_mean	radius_se	texture_se
6.279761e-02	4.051721e-01	1.216853e+00
perimeter_se	area_se	smoothness_se
2.866059e+00	4.033708e+01	7.040979e-03
compactness_se	concavity_se	concave.points_se
2.547814e-02	3.189372e-02	1.179614e-02
symmetry_se	fractal_dimension_se	radius_worst
2.054230e-02	3.794904e-03	1.626919e+01

texture_worst	perimeter_worst	area_worst
2.567722e+01	1.072612e+02	8.805831e+02
smoothness_worst	compactness_worst	concavity_worst
1.323686e-01	2.542650e-01	2.721885e-01
concave.points_worst	symmetry_worst	fractal_dimension_worst
1.146062e-01	2.900756e-01	8.394582e-02

```
apply(wisc.data, 2, sd)
```

radius_mean	texture_mean	perimeter_mean
3.524049e+00	4.301036e+00	2.429898e+01
area_mean	smoothness_mean	compactness_mean
3.519141e+02	1.406413e-02	5.281276e-02
concavity_mean	concave.points_mean	symmetry_mean
7.971981e-02	3.880284e-02	2.741428e-02
fractal_dimension_mean	radius_se	texture_se
7.060363e-03	2.773127e-01	5.516484e-01
perimeter_se	area_se	smoothness_se
2.021855e+00	4.549101e+01	3.002518e-03
compactness_se	concavity_se	concave.points_se
1.790818e-02	3.018606e-02	6.170285e-03
symmetry_se	fractal_dimension_se	radius_worst
8.266372e-03	2.646071e-03	4.833242e+00
texture_worst	perimeter_worst	area_worst
6.146258e+00	3.360254e+01	5.693570e+02
smoothness_worst	compactness_worst	concavity_worst
2.283243e-02	1.573365e-01	2.086243e-01
concave.points_worst	symmetry_worst	fractal_dimension_worst
6.573234e-02	6.186747e-02	1.806127e-02

We need to scale our input data before PCA as one of the columns are measured in terms of very different units with different means and different variances. I'm going to set `scale = TRUE` to fix this.

```
wisc.pr<-prcomp(wisc.data, scale = TRUE)
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172

Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

Q4. From your results, what proportion of the original variance is captured by the first principal components (PC1)?

44.27% of the original variance is captured by PC1.

Q5. How many principal components (PCs) are required to describe at least 70% of the original variance in the data?

3 PCs are required to describe at least 70% of the original variance in the data.

Q6. How many principal components (PCs) are required to describe at least 90% of the original variance in the data?

7 PCs are required to describe at least 90% of the original variance in the data.


## Interpreting PCA Results

Plotting it!

```
biplot(wisc.pr)
```








Q8. Generate a similar plot for principal components 1 and 3. What do you notice about these plots?

```
plot(wisc.pr$x[,1], wisc.pr$x[,3], col=diagnosis)
```




The first plot is better because the two groups are more clearly defined.

ggPlot time!

```
df <- as.data.frame(wisc.pr$x)
df$diagnosis <- diagnosis
library(ggplot2)

ggplot(df) +
  aes (PC1, PC2, col=diagnosis) +
  geom_point()
```




## Variance Explained

```
pr.var <- wisc.pr$sdev^2  
head(pr.var)
```


```
[1] 13.281608  5.691355  2.817949  1.980640  1.648731  1.207357
```

```
pve <- pr.var / sum(pr.var)  
plot(pve, xlab="Principal Component",  
      ylab="Proportion of Variance Explained",  
      ylim=c(0,1), type="o")
```



Optional Stuff


```
barplot(pve, ylab = "Precent of Variance Explained",
        names.arg=paste0("PC",1:length(pve)), las=2, axes = FALSE)
axis(2, at=pve, labels=round(pve,2)*100 )
```



```
#install.packages("factoextra")  
library(factoextra)
```

Welcome! Want to learn more? See two factoextra-related books at <https://goo.gl/ve3WBa>

```
fviz_eig(wisc.pr, addlabels = TRUE)
```



## Communicating PCA Results

Q9. For the first principal component, what is the component of the loading vector (i.e. `wisc.pr$rotation[,1]`) for the feature `concave.points_mean`?

```
wisc.pr$rotation[,1]
```

radius_mean	texture_mean	perimeter_mean
-0.21890244	-0.10372458	-0.22753729
area_mean	smoothness_mean	compactness_mean
-0.22099499	-0.14258969	-0.23928535
concavity_mean	concave.points_mean	symmetry_mean
-0.25840048	-0.26085376	-0.13816696
fractal_dimension_mean	radius_se	texture_se
-0.06436335	-0.20597878	-0.01742803
perimeter_se	area_se	smoothness_se
-0.21132592	-0.20286964	-0.01453145
compactness_se	concavity_se	concave.points_se
-0.17039345	-0.15358979	-0.18341740
symmetry_se	fractal_dimension_se	radius_worst
-0.04249842	-0.10256832	-0.22799663

texture_worst	perimeter_worst	area_worst
-0.10446933	-0.23663968	-0.22487053
smoothness_worst	compactness_worst	concavity_worst
-0.12795256	-0.21009588	-0.22876753
concave.points_worst	symmetry_worst	fractal_dimension_worst
-0.25088597	-0.12290456	-0.13178394

The component of the loading vector for concave.points\_mean is -0.26085376.

Q10. What is the minimum number of principal components required to explain 80% of the variance of the data?

```
summary(wisc.pr)
```

Importance of components:

	PC1	PC2	PC3	PC4	PC5	PC6	PC7
Standard deviation	3.6444	2.3857	1.67867	1.40735	1.28403	1.09880	0.82172
Proportion of Variance	0.4427	0.1897	0.09393	0.06602	0.05496	0.04025	0.02251
Cumulative Proportion	0.4427	0.6324	0.72636	0.79239	0.84734	0.88759	0.91010
	PC8	PC9	PC10	PC11	PC12	PC13	PC14
Standard deviation	0.69037	0.6457	0.59219	0.5421	0.51104	0.49128	0.39624
Proportion of Variance	0.01589	0.0139	0.01169	0.0098	0.00871	0.00805	0.00523
Cumulative Proportion	0.92598	0.9399	0.95157	0.9614	0.97007	0.97812	0.98335
	PC15	PC16	PC17	PC18	PC19	PC20	PC21
Standard deviation	0.30681	0.28260	0.24372	0.22939	0.22244	0.17652	0.1731
Proportion of Variance	0.00314	0.00266	0.00198	0.00175	0.00165	0.00104	0.0010
Cumulative Proportion	0.98649	0.98915	0.99113	0.99288	0.99453	0.99557	0.9966
	PC22	PC23	PC24	PC25	PC26	PC27	PC28
Standard deviation	0.16565	0.15602	0.1344	0.12442	0.09043	0.08307	0.03987
Proportion of Variance	0.00091	0.00081	0.0006	0.00052	0.00027	0.00023	0.00005
Cumulative Proportion	0.99749	0.99830	0.9989	0.99942	0.99969	0.99992	0.99997
	PC29	PC30					
Standard deviation	0.02736	0.01153					
Proportion of Variance	0.00002	0.00000					
Cumulative Proportion	1.00000	1.00000					

5 PCs are required to explain 80% of the variance of the data.

### 3. Hierarchical Clustering

Scaling the data

```
data.scaled <- scale(wisc.data)
```


Finding Euclidian Distance

```
data.dist <- dist(data.scaled)
```

Creating a cluster

```
wisc.hclust <- hclust(data.dist, method = "complete" )  
plot(wisc.hclust)
```

### Cluster Dendrogram




data.dist  
hclust (\*, "complete")

Q11. Using the plot() and abline() functions, what is the height at which the clustering model has 4 clusters?

```
plot(wisc.hclust)  
abline(h=19, col = "red", lty=2)
```



## Cluster Dendrogram



```
data.dist  
hclust (*, "complete")
```

At height 19 there are 4 clusters.

## Selecting number of clusters

```
wisc.hclust.clusters <- cutree(wisc.hclust, k=4)  
  
table(wisc.hclust.clusters, diagnosis)
```

	diagnosis	
wisc.hclust.clusters	B	M
1	12	165
2	2	5
3	343	40
4	0	2

Q12. Can you find a better cluster vs diagnoses match by cutting into a different number of clusters between 2 and 10?

I can't. 4 clusters seems to be the best match.

## Using Different Methods


Q13. Which method gives your favorite results for the same data.dist dataset?  
Explain your reasoning.

ward.D2 gives my favorite results because it's the easiest to look at. It's the best organized.

## 5. Combining Methods

This approach will take not original data, but our PCA results and work with them.

```
d <- dist(wisc.pr$x[,1:3])  
wisc.pr.hclust <- hclust(d, method = "ward.D2")  
plot(wisc.pr.hclust)
```



Generating 2 cluster groups from this hclust object

```
grps <- cutree(wisc.pr.hclust, k=2)  
table(grps)
```


```
grps
  1  2
203 366
```

```
table(grps, diagnosis)
```


```
      diagnosis
grps   B     M
  1   24  179
  2  333   33
```

Plotting it!

```
plot(wisc.pr$x[,1], wisc.pr$x[,2], col=grps)
```



```
plot(wisc.pr$x[,1], wisc.pr$x[,2], col=diagnosis)
```



Making the colors match up!


```
g <- as.factor(grps)
levels(g)
```

```
[1] "1" "2"
```

```
g <- relevel(g,2)
levels(g)
```

```
[1] "2" "1"
```

```
plot(wisc.pr$x[,1:2], col=g)
```



Using more PCs to try and more accurately predict diagnosis!

```
x <- dist(wisc.pr$x[,1:7])
wisc.pr.hclust2 <- hclust(x, method="ward.D2")

wisc.pr.hclust.clusters <- cutree(wisc.pr.hclust2, k=2)

table(wisc.pr.hclust.clusters, diagnosis)
```

	diagnosis	
wisc.pr.hclust.clusters	B	M
1	28	188
2	329	24

Q15. How well does the newly created model with four clusters separate out the two diagnoses?

The newly created model separates out the two diagnosis pretty well. Not quite as well as the other model, but still good.

**END**