

A Finer Calibration Analysis for Adversarial Robustness

Pranjal Awasthi

Google Research and Rutgers University, New York

PRANJAL.AWASTHI@RUTGERS.EDU

Anqi Mao

Courant Institute of Mathematical Sciences, New York

AQMAO@CIMS.NYU.EDU

Mehryar Mohri

Google Research and Courant Institute of Mathematical Sciences, New York

MOHRI@GOOGLE.COM

Yutao Zhong

Courant Institute of Mathematical Sciences, New York

YUTAO@CIMS.NYU.EDU

Abstract

We present a more general analysis of \mathcal{H} -calibration for adversarially robust classification. By adopting a finer definition of calibration, we can cover settings beyond the restricted hypothesis sets studied in previous work. In particular, our results hold for most common hypothesis sets used in machine learning. We both fix some previous calibration results (Bao et al., 2020) and generalize others (Awasthi et al., 2021). Moreover, our calibration results, combined with the previous study of consistency by Awasthi et al. (2021), also lead to more general \mathcal{H} -consistency results covering common hypothesis sets.

Keywords: calibration, consistency, adversarial robustness.

1. Introduction

Rich learning models trained on large datasets often achieve a high accuracy in a variety of applications (Sutskever et al., 2014; Krizhevsky et al., 2012). However, such complex models have been shown to be susceptible to imperceptible perturbations (Szegedy et al., 2013): an unnoticeable perturbation can, for example, result in a dog being classified as an electronics device, which could lead to dramatic consequences in practice in many applications.

This has motivated the introduction and analysis of the notion of *adversarial loss*, which requires a predictor not only to correctly classify an input point \mathbf{x} but also to maintain the same classification for all points at a small ℓ_p distance of \mathbf{x} (Goodfellow et al., 2014; Madry et al., 2017; Tsipras et al., 2018; Carlini and Wagner, 2017).

The problem of designing effective learning algorithms with theoretical guarantees for the *adversarial loss* has been the topic of a number of recent studies (Bao et al., 2020; Awasthi et al., 2021). In particular, these authors have initiated a theoretical analysis of the \mathcal{H} -calibration and \mathcal{H} -consistency of surrogate losses for the *adversarial 0/1 loss*.

Bao et al. (2020) analyzed \mathcal{H} -calibration for adversarially robust classification in the special case where \mathcal{H} is the family of linear models. However, several comments are due regarding that work. First, the definition of calibration adopted by the authors does not coincide with the standard definition (Steinwart, 2007) in the case of the linear models they study, although it does match that definition in the case of the family of all measurable functions (Steinwart, 2007, Section 4.1): the minimal inner risk in the definition should be defined for a fixed \mathbf{x} and the infimum should be over f , instead of an infimum over both f and \mathbf{x} . Second, and this is crucial, \mathcal{H} -calibration, in

general, does not imply \mathcal{H} -consistency, unless a property such as \mathcal{P} -minimizability holds (Steinwart, 2007, Theorem 2.8). \mathcal{P} -minimizability holds for standard binary classification and the family of all measurable functions (Steinwart, 2007, Theorem 3.2). However, it does not hold, in general, for adversarially robust classification and a specific hypothesis set \mathcal{H} . As a result, the claim made by the authors that the calibrated surrogates they propose are \mathcal{H} -consistent is incorrect, as shown by Awasthi et al. (2021). Third, the authors analyze \mathcal{H} -calibration with respect to the loss function $\phi_\gamma: \mathbf{x} \mapsto \mathbb{1}_{yf(\mathbf{x}) \leq \gamma}$ in the case where $\mathcal{H} \supset [-1, 1]$ is the general family of functions. However, ϕ_γ only coincides with the adversarial 0/1 loss ℓ_γ in Equation (10) in the special case where \mathcal{H} is the family of linear models (Bao et al., 2020, Proposition 1).

Awasthi et al. (2021) also recently studied the \mathcal{H} -calibration and \mathcal{H} -consistency of adversarial surrogate losses. They pointed out the issues just mentioned about the study of Bao et al. (2020) and considered more general hypothesis sets, such as generalized linear models, including ReLU-based functions, and one-layer ReLU neural networks. They identified natural conditions under which \mathcal{H} -calibrated losses can be \mathcal{H} -consistent in the adversarial scenario. They also derived calibration results under the correct definition of the minimal inner risk by analyzing the equivalence of two definitions. However, with this method of calibration analysis, the calibration considered by the authors needs to be a uniform calibration (Steinwart, 2007, Definition 2.15) instead of non-uniform calibration (Steinwart, 2007, Definition 2.7). In view of that, their positive result imposes an extra restriction on the parameters of the hypothesis sets, which can be removed through the analysis presented here.

Our Contributions. Building on previous work by Awasthi et al. (2021), we present a more general analysis of \mathcal{H} -calibration for adversarially robust classification for more general hypothesis sets. For example, our Theorem 8, Theorem 10 and Theorem 16 apply to most common hypothesis sets. Furthermore, for the specific hypothesis sets considered in previous work, our results either fix existing calibration results (Bao et al., 2020) or generalize them (Awasthi et al., 2021). More precisely, our Theorem 12 is a correction to the main positive result, Theorem 11 in (Bao et al., 2020), where we prove the theorem under the correct calibration definition. Moreover, our Theorem 13 extends the results for linear models to generalized linear models. Our Corollary 9 and Corollary 11 are stronger versions of the negative calibration results Theorem 10, Corollary 11, Theorem 12 and Corollary 13 in (Awasthi et al., 2021), since the calibration considered in (Awasthi et al., 2021) is uniform calibration (Steinwart, 2007, Definition 2.15), which is stronger than non-uniform calibration (Steinwart, 2007, Definition 2.7) considered in our paper. Our Theorem 15 and Corollary 17 are generalizations of the positive calibration results of Awasthi et al. (2021), since our results hold without the unboundedness assumptions for parameters of the hypothesis sets.

2. Preliminaries

We adopt much of the notation used in (Awasthi et al., 2021). We will denote vectors as lowercase bold letters (e.g. \mathbf{x}). The d -dimensional l_2 -ball with radius r is denoted by $B_2^d(r) := \{\mathbf{z} \in \mathbb{R}^d \mid \|\mathbf{z}\|_2 \leq r\}$. We denote by \mathcal{X} the set of all possible examples. \mathcal{X} is also sometimes referred to as the input space. The set of all possible labels is denoted by \mathcal{Y} . We will limit ourselves to the case of binary classification where $\mathcal{Y} = \{-1, 1\}$. Let \mathcal{H} be a family of functions from \mathbb{R}^d to \mathbb{R} . Given a fixed but unknown distribution \mathcal{P} over $\mathcal{X} \times \mathcal{Y}$, the binary classification learning problem is then formulated as follows. The learner seeks to select a predictor $f \in \mathcal{H}$ with small *generalization error* with respect to the distribution \mathcal{P} . The *generalization error* of a classifier $f \in \mathcal{H}$ is defined

by $\mathcal{R}_{\ell_0}(f) = \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{P}}[\ell_0(f, \mathbf{x}, y)]$, where $\ell_0(f, \mathbf{x}, y) = \mathbb{1}_{yf(\mathbf{x}) \leq 0}$ is the standard 0/1 loss. More generally, the ℓ -risk of a classifier f for a surrogate loss $\ell(f, \mathbf{x}, y)$ is defined by

$$\mathcal{R}_{\ell}(f) = \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{P}}[\ell(f, \mathbf{x}, y)]. \quad (1)$$

Moreover, the *minimal* (ℓ, \mathcal{H}) -risk, which is also called the *Bayes* (ℓ, \mathcal{H}) -risk, is defined by $\mathcal{R}_{\ell, \mathcal{H}}^* = \inf_{f \in \mathcal{H}} \mathcal{R}_{\ell}(f)$. In the standard classification setting, the goal of a consistency analysis is to determine whether the minimization of a surrogate loss ℓ can lead to that of the binary loss generalization error. Similarly, in adversarially robust classification, the goal of a consistency analysis is to determine if the minimization of a surrogate loss ℓ yields that of the *adversarial generalization error* defined by $\mathcal{R}_{\ell_{\gamma}}(f) = \mathbb{E}_{(\mathbf{x}, y) \sim \mathcal{P}}[\ell_{\gamma}(f, \mathbf{x}, y)]$, where

$$\ell_{\gamma}(f, \mathbf{x}, y) := \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \mathbb{1}_{yf(\mathbf{x}') \leq 0} \quad (2)$$

is the *adversarial* 0/1 loss. This motivates the definition of \mathcal{H} -consistency (or simply *consistency*) stated below.

Definition 1 (\mathcal{H} -Consistency) *Given a hypothesis set \mathcal{H} , we say that a loss function ℓ_1 is \mathcal{H} -consistent with respect to loss function ℓ_2 , if the following holds:*

$$\mathcal{R}_{\ell_1}(f_n) - \mathcal{R}_{\ell_1, \mathcal{H}}^* \xrightarrow{n \rightarrow +\infty} 0 \implies \mathcal{R}_{\ell_2}(f_n) - \mathcal{R}_{\ell_2, \mathcal{H}}^* \xrightarrow{n \rightarrow +\infty} 0, \quad (3)$$

for all probability distributions and sequences of $\{f_n\}_{n \in \mathbb{N}} \subset \mathcal{H}$.

For a distribution \mathcal{P} over $\mathcal{X} \times \mathcal{Y}$ with random variables X and Y , let $\eta_{\mathcal{P}}: \mathcal{X} \rightarrow [0, 1]$ be a measurable function such that, for any $\mathbf{x} \in \mathcal{X}$, $\eta_{\mathcal{P}}(\mathbf{x}) = \mathcal{P}(Y = 1 \mid X = \mathbf{x})$. By the property of conditional expectation, we can rewrite (1) as $\mathcal{R}_{\ell}(f) = \mathbb{E}_X[\mathcal{C}_{\ell}(f, \mathbf{x}, \eta_{\mathcal{P}}(\mathbf{x}))]$, where $\mathcal{C}_{\ell}(f, \mathbf{x}, \eta)$ is the *generic conditional ℓ -risk* (or *inner ℓ -risk*) defined as followed:

$$\forall \mathbf{x} \in \mathcal{X}, \forall \eta \in [0, 1], \quad \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) := \eta \ell(f, \mathbf{x}, +1) + (1 - \eta) \ell(f, \mathbf{x}, -1). \quad (4)$$

Moreover, the *minimal inner ℓ -risk* on \mathcal{H} is denoted by $\mathcal{C}_{\ell, \mathcal{H}}^*(\mathbf{x}, \eta) := \inf_{f \in \mathcal{H}} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta)$. The notion of *calibration* for the inner risk is often a powerful tool for the analysis of \mathcal{H} -consistency (Steinwart, 2007).

Definition 2 (\mathcal{H} -Calibration) [Definition 2.7 in (Steinwart, 2007)] *Given a hypothesis set \mathcal{H} , we say that a loss function ℓ_1 is \mathcal{H} -calibrated with respect to a loss function ℓ_2 if, for any $\epsilon > 0$, $\eta \in [0, 1]$, and $\mathbf{x} \in \mathcal{X}$, there exists $\delta > 0$ such that for all $f \in \mathcal{H}$ we have*

$$\mathcal{C}_{\ell_1}(f, \mathbf{x}, \eta) < \mathcal{C}_{\ell_1, \mathcal{H}}^*(\mathbf{x}, \eta) + \delta \implies \mathcal{C}_{\ell_2}(f, \mathbf{x}, \eta) < \mathcal{C}_{\ell_2, \mathcal{H}}^*(\mathbf{x}, \eta) + \epsilon. \quad (5)$$

For comparison with previous work, we also introduce the *uniform \mathcal{H} -calibration* in (Steinwart, 2007), which is stronger than Definition 2.

Definition 3 (Uniform \mathcal{H} -Calibration) [Definition 2.15 in (Steinwart, 2007)] *Given a hypothesis set \mathcal{H} , we say that a loss function ℓ_1 is uniform \mathcal{H} -calibrated with respect to a loss function ℓ_2 if, for any $\epsilon > 0$, there exists $\delta > 0$ such that for all $\eta \in [0, 1]$, $f \in \mathcal{H}$, $\mathbf{x} \in \mathcal{X}$, we have*

$$\mathcal{C}_{\ell_1}(f, \mathbf{x}, \eta) < \mathcal{C}_{\ell_1, \mathcal{H}}^*(\mathbf{x}, \eta) + \delta \implies \mathcal{C}_{\ell_2}(f, \mathbf{x}, \eta) < \mathcal{C}_{\ell_2, \mathcal{H}}^*(\mathbf{x}, \eta) + \epsilon. \quad (6)$$

Note that, in the previous work of [Awasthi et al. \(2021\)](#), Definition 3 is adopted, where δ in (6) is independent of η and \mathbf{x} ; the work of [Bao et al. \(2020\)](#) adopts a similar definition. In this paper, we will focus on the non-uniform case, that is Definition 2, where δ is dependent on η and \mathbf{x} . There are two advantages to considering non-uniform calibration: it makes it possible to provide stronger negative results on calibration properties of convex surrogates and, it helps us prove more general positive results that hold for most common hypothesis sets \mathcal{H} . In contrast, positive results for uniform calibration hold for some restricted hypothesis sets ([Awasthi et al., 2021](#)).

[Steinwart \(2007\)](#) showed that if ℓ_1 is \mathcal{H} -calibrated (it suffices to satisfy non-uniform calibration, that is condition (5)) with respect to ℓ_2 , then \mathcal{H} -consistency, that is condition (3), holds for any probability distribution verifying the additional condition of \mathcal{P} -minimizability ([Steinwart, 2007](#), Definition 2.4). While \mathcal{P} -minimizability does not hold in general for adversarially robust classification, [Awasthi et al. \(2021\)](#) showed that the uniform \mathcal{H} -calibrated losses are \mathcal{H} -consistent under certain conditions. In fact, it also suffices to satisfy non-uniform calibration, that is condition (5) for these results, since their proofs only make use of the weaker non-uniform property.

Next, we introduce the notions of *calibration function* and an important result characterizing \mathcal{H} -calibration from ([Steinwart, 2007](#)).

Definition 4 (Calibration function) *Given a hypothesis set \mathcal{H} , we define the calibration function δ_{\max} for a pair of losses (ℓ_1, ℓ_2) as follows: for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$,*

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}} \left\{ \mathcal{C}_{\ell_1}(f, \mathbf{x}, \eta) - \mathcal{C}_{\ell_1, \mathcal{H}}^*(\mathbf{x}, \eta) \mid \mathcal{C}_{\ell_2}(f, \mathbf{x}, \eta) - \mathcal{C}_{\ell_2, \mathcal{H}}^*(\mathbf{x}, \eta) \geq \epsilon \right\}. \quad (7)$$

Proposition 5 (Lemma 2.9 in ([Steinwart, 2007](#))) *Given a hypothesis set \mathcal{H} , loss ℓ_1 is \mathcal{H} -calibrated with respect to ℓ_2 if and only if its calibration function δ_{\max} satisfies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$.*

For comparison, [Bao et al. \(2020, Definition 2\)](#) and [Awasthi et al. \(2021, Definition 2\)](#) consider the *Uniform Calibration function* $\delta(\epsilon)$ and make use of Lemma 2.16 in ([Steinwart, 2007](#)) to characterize uniform calibration ([Awasthi et al., 2021](#); [Bao et al., 2020, Proposition 5](#)). Note $\delta(\epsilon) > 0$ implies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$, and as a result uniform calibration implies non-uniform calibration. However, the converse does not hold in general.

3. Adversarially Robust Classification

In adversarially robust classification, the loss at (\mathbf{x}, y) is measured in terms of the worst loss incurred over an adversarial perturbation of \mathbf{x} within a ball of a certain radius in a norm. In this work we will consider perturbations in the l_2 norm $\|\cdot\|$. We will denote by γ the maximum magnitude of the allowed perturbations. Given $\gamma > 0$, a data point (\mathbf{x}, y) , a function $f \in \mathcal{H}$, and a margin-based loss $\phi: \mathbb{R} \rightarrow \mathbb{R}_+$, we define the *adversarial loss* of f at (\mathbf{x}, y) as

$$\tilde{\phi}(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}')). \quad (8)$$

The above naturally motivates *supremum-based* surrogate losses that are commonly used to optimize the adversarial 0/1 loss ([Goodfellow et al., 2014](#); [Madry et al., 2017](#); [Shafahi et al., 2019](#); [Wong et al., 2020](#)). We say that a surrogate loss $\tilde{\phi}(f, \mathbf{x}, y)$ is *supremum-based* if it is of the form

defined in (8). We say that the supremum-based surrogate is convex if the function ϕ in (8) is convex. When ϕ is non-increasing, the following equality holds (Yin et al., 2019):

$$\sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}')) = \phi\left(\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} yf(\mathbf{x}')\right). \quad (9)$$

The adversarial 0/1 loss defined in (2) is a special kind of adversarial loss (8), where ϕ is the 0/1 loss, that is, $\phi(yf(\mathbf{x})) = \ell_0(f, \mathbf{x}, y) = \mathbb{1}_{yf(\mathbf{x}) \leq 0}$. Therefore, the adversarial 0/1 loss has the equivalent form

$$\ell_\gamma(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \mathbb{1}_{yf(\mathbf{x}') \leq 0} = \mathbb{1}_{\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} yf(\mathbf{x}') \leq 0}. \quad (10)$$

This alternative equivalent form of adversarial 0/1 loss is more advantageous to analyze than (2) and would be adopted in our proofs. Without loss of generality, let $\mathcal{X} = B_2^d(1)$ and $\gamma \in (0, 1)$. In this paper, we aim to characterize surrogate losses ℓ_1 satisfying \mathcal{H} -calibration (5) with $\ell_2 = \ell_\gamma$ and for the hypothesis sets \mathcal{H} which are *regular for adversarial calibration*.

Definition 6 (Regularity for Adversarial Calibration) We say that a hypothesis set \mathcal{H} is regular for adversarial calibration if there exists a distinguishing \mathbf{x} in \mathcal{X} , that is if there exist $f, g \in \mathcal{H}$ such that $\inf_{\|\mathbf{x}' - \mathbf{x}\| \leq \gamma} f(\mathbf{x}') > 0$ and $\sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \gamma} g(\mathbf{x}') < 0$.

It suffices to study hypothesis sets \mathcal{H} that are regular for adversarial calibration not only because all common hypothesis sets admit that property, but also because the following result holds. We say that a hypothesis set \mathcal{H} is *symmetric*, if for any $f \in \mathcal{H}$, $-f$ is also in \mathcal{H} .

Theorem 7 Let \mathcal{H} be a symmetric hypothesis set. If \mathcal{H} is not regular for adversarial calibration, then any surrogate loss ℓ is \mathcal{H} -calibrated with respect to ℓ_γ .

Proof Since \mathcal{H} is symmetric, for any $\mathbf{x} \in \mathcal{X}$, $f \in \mathcal{H}$, $\inf_{\|\mathbf{x}' - \mathbf{x}\| \leq \gamma} f(\mathbf{x}') \leq 0 \leq \sup_{\|\mathbf{x}' - \mathbf{x}\| \leq \gamma} f(\mathbf{x}')$. Thus by the definition of inner risk (4) and adversarial 0-1 loss ℓ_γ (10), for any $\mathbf{x} \in \mathcal{X}$, $f \in \mathcal{H}$,

$$\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = \eta \mathbb{1}_{\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} f(\mathbf{x}') \leq 0} + (1 - \eta) \mathbb{1}_{\sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} f(\mathbf{x}') \geq 0} = 1 = \mathcal{C}_{\ell_\gamma, \mathcal{H}}^*(\mathbf{x}, \eta),$$

which implies any surrogate loss ℓ is \mathcal{H} -calibrated with respect to ℓ_γ by (5). \blacksquare

Note all the hypothesis sets considered in the previous work (Bao et al., 2020) and (Awasthi et al., 2021) are regular for adversarial calibration. For convenience, we adopt the notation in (Awasthi et al., 2021) to denote these specific hypothesis sets:

- linear models: $\mathcal{H}_{\text{lin}} = \{\mathbf{x} \rightarrow \mathbf{w} \cdot \mathbf{x} \mid \|\mathbf{w}\| = 1\}$, as in (Bao et al., 2020) and (Awasthi et al., 2021).
- generalized linear models: $\mathcal{H}_g = \{\mathbf{x} \rightarrow g(\mathbf{w} \cdot \mathbf{x}) + b \mid \|\mathbf{w}\| = 1, |b| \leq G\}$ where g is a non-decreasing function, as in (Awasthi et al., 2021); and
- one-layer ReLU neural networks: $\mathcal{H}_{\text{NN}} = \{\mathbf{x} \rightarrow \sum_{j=1}^n u_j(\mathbf{w}_j \cdot \mathbf{x})_+ \mid \|\mathbf{u}\|_1 \leq \Lambda, \|\mathbf{w}_j\| \leq W\}$, where $(\cdot)_+ = \max(\cdot, 0)$ as in (Awasthi et al., 2021); and
- all measurable functions: \mathcal{H}_{all} as in (Awasthi et al., 2021).

In the special case of $g = (\cdot)_+$, we denote the corresponding ReLU-based hypothesis set as $\mathcal{H}_{\text{relu}} = \{\mathbf{x} \rightarrow (\mathbf{w} \cdot \mathbf{x})_+ + b \mid \|\mathbf{w}\| = 1, |b| \leq G\}$ as in (Awasthi et al., 2021).

4. \mathcal{H} -Calibration Analysis

4.1. Negative results

In this section, we show that the commonly used convex surrogates and supremum-based convex surrogates are not \mathcal{H} -calibrated with respect to ℓ_γ , even under the weaker notion of non-uniform calibration. These results can be viewed as a generalization of those given by [Awasthi et al. \(2021\)](#).

4.1.1. CONVEX LOSSES

We first study convex losses, which are often used for standard binary classification problems.

Theorem 8 *Let \mathcal{H} be a hypothesis set containing 0 that is regular for adversarial calibration. If a margin-based loss $\phi: \mathbb{R} \rightarrow \mathbb{R}_+$ is convex, then it is not \mathcal{H} -calibrated with respect to ℓ_γ .*

The proof of Theorem 8 is included in Appendix A.1. By Theorem 8, we obtain the following corollary, which fixes the main negative result of [Bao et al. \(2020\)](#) and generalizes negative results of [Awasthi et al. \(2021\)](#). Note \mathcal{H}_{lin} , $\mathcal{H}_{\text{relu}}$ and \mathcal{H}_{all} are all regular for adversarial calibration and contain 0. When $|g(0)| \leq G$, $0 \in \mathcal{H}_g$; when $g(1 - \gamma) + G > 0$ and $g(-1 + \gamma) - G < 0$, \mathcal{H}_g is regular for adversarial calibration.

Corollary 9 *If a margin-based loss $\phi: \mathbb{R} \rightarrow \mathbb{R}_+$ is convex, then,*

1. ϕ is not \mathcal{H}_{lin} -calibrated with respect to ℓ_γ ;
2. Given a non-decreasing and continuous function g such that $|g(0)| \leq G$, $g(1 - \gamma) + G > 0$ and $g(-1 + \gamma) - G < 0$. Then ϕ is not \mathcal{H}_g -calibrated with respect to ℓ_γ ; Specifically, ϕ is not $\mathcal{H}_{\text{relu}}$ -calibrated with respect to ℓ_γ ;
3. ϕ is not \mathcal{H}_{NN} -calibrated with respect to ℓ_γ ;
4. ϕ is not \mathcal{H}_{all} -calibrated with respect to ℓ_γ .

By using the correct calibration Definition 2, 1. of Corollary 9 fixes the main negative result in [\(Bao et al., 2020\)](#).

4.1.2. SUPREMUM-BASED CONVEX LOSSES

While it is natural to consider convex surrogates for the 0/1 loss, convex supremum-based surrogates are widely used in practice for designing algorithms for the adversarial loss ([Madry et al., 2017](#); [Shafahi et al., 2019](#); [Wong et al., 2020](#)). We next present a negative result for convex supremum-based surrogates.

Theorem 10 *Let \mathcal{H} be a hypothesis set containing 0 that is regular for adversarial calibration. If a margin-based loss ϕ is convex and non-increasing, then the surrogate loss defined by $\tilde{\phi}(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}'))$ is not \mathcal{H} -calibrated with respect to ℓ_γ .*

The proof of Theorem 10 is also included in Appendix A.1. Theorem 10 leads to the following corollary.

Corollary 11 *Let ϕ be convex and non-increasing margin-based loss, consider the surrogate loss defined by $\tilde{\phi}(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}'))$. Then*

1. $\tilde{\phi}$ is not \mathcal{H}_{lin} -calibrated with respect to ℓ_γ ;
2. Given a non-decreasing and continuous function g such that $|g(0)| \leq G$, $g(1 - \gamma) + G > 0$ and $g(-1 + \gamma) - G < 0$. Then $\tilde{\phi}$ is not \mathcal{H}_g -calibrated with respect to ℓ_γ ; Specifically, $\tilde{\phi}$ is not $\mathcal{H}_{\text{relu}}$ -calibrated with respect to ℓ_γ ;
3. $\tilde{\phi}$ is not \mathcal{H}_{NN} -calibrated with respect to ℓ_γ ;
4. $\tilde{\phi}$ is not \mathcal{H}_{all} -calibrated with respect to ℓ_γ .

Corollary 9 and Corollary 11 above are stronger versions of the negative calibration results Theorem 10, Corollary 11, Theorem 12 and Corollary 13 in (Awasthi et al., 2021), since the calibration considered in (Awasthi et al., 2021) is uniform calibration (Steinwart, 2007, Definition 2.15), which is stronger than non-uniform calibration (Steinwart, 2007, Definition 2.7) considered in this work.

4.2. Positive results

In this section, we provide alternative surrogate losses that are \mathcal{H} -calibrated with respect to ℓ_γ . These results are similar but more general than their counterparts in (Awasthi et al., 2021),

4.2.1. MARGIN-BASED LOSSES

In light of the negative results of Section 4.1, to find calibrated surrogate losses for adversarially robust classification, we need to consider non-convex ones. One possible candidate is the family of *quasi-concave even* losses introduced by Bao et al. (2020). Theorem 12 below is a correction to the main positive result, Theorem 11 in (Bao et al., 2020), where we prove the theorem under the correct calibration definition.

Theorem 12 *Let a margin-based loss ϕ be bounded, continuous, non-increasing, and quasi-concave even. Assume that $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$. Then ϕ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ if and only if for any $\gamma < t \leq 1$,*

$$\phi(\gamma) + \phi(-\gamma) > \phi(t) + \phi(-t). \quad (11)$$

The proof of Theorem 12 is included in Appendix A.3, where we make use of Lemma 25, which is powerful since it applies to any symmetric hypothesis sets containing 0. Note Bao et al. (2020, Theorem 11) does not hold any more under the correct calibration Definition 2, since their condition $\phi(\gamma) + \phi(-\gamma) > \phi(1) + \phi(-1)$ is much weaker than (11).

We next extend the above to show that under certain conditions, quasi-concave even surrogate losses are \mathcal{H}_g -calibrated for the class of generalized linear models with respect to the adversarial 0/1 loss.

Theorem 13 *Let g be a non-decreasing and continuous function such that $g(1 + \gamma) < G$ and $g(-1 - \gamma) > -G$ for some $G \geq 0$. Let a margin-based loss ϕ be bounded, continuous, non-increasing, and quasi-concave even. Assume that $\phi(g(-t) - G) > \phi(G - g(-t))$ and $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$. Then ϕ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if for any $0 \leq t \leq 1$,*

$$\begin{aligned} & \phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G) \\ \text{and } & \min\{\phi(\overline{A}(t)) + \phi(-\overline{A}(t)), \phi(\underline{A}(t)) + \phi(-\underline{A}(t))\} > \phi(G - g(-t)) + \phi(g(-t) - G), \end{aligned}$$

where $\overline{A}(t) = \max_{s \in [-t, t]} g(s) - g(s - \gamma)$ and $\underline{A}(t) = \min_{s \in [-t, t]} g(s) - g(s + \gamma)$.

The proof of Theorem 13 is included in Appendix A.4. Specifically, when $g = (\cdot)_+$, by Theorem 13, we obtain the following corollary for $\mathcal{H}_{\text{relu}}$ by using that fact that $\phi(t) + \phi(-t) \geq \phi(\gamma) + \phi(-\gamma)$ when $0 \leq t \leq \gamma$ by Part 2 of Lemma 23. Note when $g = (\cdot)_+$,

$$\begin{aligned}\overline{A}(t) &= \max_{s \in [-t, t]} (s)_+ - (s - \gamma)_+ = \begin{cases} t, & 0 \leq t < \gamma, \\ \gamma, & \gamma \leq t \leq 1. \end{cases} \\ \underline{A}(t) &= \min_{s \in [-t, t]} (s)_+ - g(s + \gamma)_+ = -\gamma.\end{aligned}$$

Corollary 14 *Assume that $G > 1 + \gamma$. Let a margin-based loss ϕ be bounded, continuous, non-increasing, and quasi-concave even. Assume that $\phi(-G) > \phi(G)$. Then ϕ is $\mathcal{H}_{\text{relu}}$ -calibrated with respect to ℓ_γ if and only if for any $0 \leq t \leq 1$,*

$$\phi(G) + \phi(-G) = \phi(t + G) + \phi(-t - G) \quad \text{and} \quad \phi(\gamma) + \phi(-\gamma) > \phi(G) + \phi(-G).$$

In order to demonstrate the applicability of Theorem 12, Theorem 13 and Corollary 14, we consider a specific surrogate loss namely the ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$, which is a generalization of the ramp loss (see, for example, Mohri et al. (2018)). Using Theorem 12, Theorem 13 and Corollary 14, we can conclude that the ρ -margin loss is calibrated under reasonable conditions for linear hypothesis sets and non-decreasing g -based hypothesis sets, since $\phi_\rho(t)$ is bounded, non-increasing and quasi-concave even. This is stated formally below.

Theorem 15 *Consider ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$. Then,*

1. ϕ_ρ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ if and only if $\rho > 1$.
2. *Given a non-decreasing and continuous function g such that $g(1 + \gamma) < G$ and $g(-1 - \gamma) > -G$ for some $G \geq 0$. Assume that $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$. Then ϕ_ρ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if for any $0 \leq t \leq 1$,*

$$\phi_\rho(G - g(-t)) = \phi_\rho(g(t) + G) \quad \text{and} \quad \min\{\phi_\rho(\overline{A}(t)), \phi_\rho(-\underline{A}(t))\} > \phi_\rho(G - g(-t)),$$

$$\text{where } \overline{A}(t) = \max_{s \in [-t, t]} g(s) - g(s - \gamma) \text{ and } \underline{A}(t) = \min_{s \in [-t, t]} g(s) - g(s + \gamma).$$

3. *Assume that $G > 1 + \gamma$. Then ϕ_ρ is $\mathcal{H}_{\text{relu}}$ -calibrated with respect to ℓ_γ if and only if $G \geq \rho > \gamma$.*

Theorem 15 is a strict generalization of the positive calibration results in (Awasthi et al., 2021) for \mathcal{H}_g and $\mathcal{H}_{\text{relu}}$ where the authors require G to be unbounded. By working with the weaker notion of non-uniform calibration, we avoid such a restriction on G .

4.2.2. SUPREMUM-BASED MARGIN LOSSES

Recall that in Theorem 10 we ruled out the possibility of finding \mathcal{H} -calibrated supremum-based convex surrogate losses with respect to the adversarial 0/1 loss. However, we show that the supremum-based ρ -margin loss is indeed \mathcal{H} -calibrated. We state the calibration result below and present the proof in Appendix A.3.

Theorem 16 *Consider ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$. Let \mathcal{H} be a symmetric hypothesis set containing 0, then the surrogate loss $\tilde{\phi}_\rho(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi_\rho(yf(\mathbf{x}'))$ is \mathcal{H} -calibrated with respect to ℓ_γ .*

By Theorem 16, we obtain the following corollary, since \mathcal{H}_{lin} , \mathcal{H}_{NN} and \mathcal{H}_{all} are all symmetric and contain 0.

Corollary 17 *Consider ρ -margin loss $\phi_\rho(t) = \min\left\{1, \max\left\{0, 1 - \frac{t}{\rho}\right\}\right\}$, $\rho > 0$. Let $\tilde{\phi}_\rho(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi_\rho(yf(\mathbf{x}'))$ be the surrogate loss. Then,*

1. $\tilde{\phi}_\rho$ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ ;
2. $\tilde{\phi}_\rho$ is \mathcal{H}_{NN} -calibrated with respect to ℓ_γ ;
3. $\tilde{\phi}_\rho$ is \mathcal{H}_{all} -calibrated with respect to ℓ_γ .

2. of Corollary 17 is a strict generalization of the positive calibration result in (Awasthi et al., 2021) for \mathcal{H}_{NN} where the authors require Λ to be unbounded. By working with the weaker notion of non-uniform calibration, we avoid such a restriction on Λ .

5. \mathcal{H} -consistency

Next, we study the implications of our positive results for non-uniform calibration for establishing \mathcal{H} -consistency. As discussed in Section 1, Steinwart (2007) showed that if ℓ_1 is \mathcal{H} -calibrated (it suffices to satisfy non-uniform calibration, that is condition (5)) with respect to ℓ_2 , then \mathcal{H} -consistency, that is condition (3), holds for any probability distribution verifying the additional condition of \mathcal{P} -minimizability (Steinwart, 2007, Definition 2.4). Although the \mathcal{P} -minimizability condition is naturally satisfied and \mathcal{H} -calibration often is a sufficient condition for \mathcal{H} -consistency in the standard classification setting when considering the family of all measurable functions (Steinwart, 2007, Theorem 3.2), Awasthi et al. (2021) points out that the adversarial loss presents new challenges when dealing with \mathcal{P} -minimizability and requires carefully distinguishing among calibration and consistency to avoid drawing false conclusions.

Moreover, Awasthi et al. (2021) shows that the \mathcal{H} -calibrated losses are \mathcal{H} -consistent under certain conditions. Analogously, in this section, we make use of Awasthi et al. (2021, Theorem 25, Theorem 27) to conclude that the \mathcal{H} -calibrated losses studied in previous sections are \mathcal{H} -consistent under the same conditions.

Theorem 18 (Theorem 25 in (Awasthi et al., 2021)) *Let \mathcal{P} be a distribution over $\mathcal{X} \times \mathcal{Y}$ and \mathcal{H} a hypothesis set for which $\mathcal{R}_{\ell_\gamma, \mathcal{H}}^* = 0$. Let ϕ be a margin-based loss. If for $\eta \geq 0$, there exists $f^* \in \mathcal{H} \subset \mathcal{H}_{\text{all}}$ such that $\mathcal{R}_\phi(f^*) \leq \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* + \eta < +\infty$ and ϕ is \mathcal{H} -calibrated with respect to ℓ_γ , then for all $\epsilon > 0$ there exists $\delta > 0$ such that for all $f \in \mathcal{H}$ we have*

$$\mathcal{R}_\phi(f) + \eta < \mathcal{R}_{\phi, \mathcal{H}}^* + \delta \implies \mathcal{R}_{\ell_\gamma}(f) < \mathcal{R}_{\ell_\gamma, \mathcal{H}}^* + \epsilon.$$

Theorem 19 (Theorem 27 in (Awasthi et al., 2021)) *Given a distribution \mathcal{P} over $\mathcal{X} \times \mathcal{Y}$ and a hypothesis set \mathcal{H} such that $\mathcal{R}_{\ell_\gamma, \mathcal{H}}^* = 0$. Let ϕ be a non-increasing margin-based loss. If there exists $f^* \in \mathcal{H} \subset \mathcal{H}_{\text{all}}$ such that $\mathcal{R}_\phi(f^*) = \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* < \infty$ and $\tilde{\phi}(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}'))$ is \mathcal{H} -calibrated with respect to ℓ_γ , then for all $\epsilon > 0$ there exists $\delta > 0$ such that for all $f \in \mathcal{H}$ we have*

$$\mathcal{R}_{\tilde{\phi}}(f) < \mathcal{R}_{\tilde{\phi}, \mathcal{H}}^* + \delta \implies \mathcal{R}_{\ell_\gamma}(f) < \mathcal{R}_{\ell_\gamma, \mathcal{H}}^* + \epsilon.$$

Using Theorem 15 in Section 4.2.1 and Theorem 18 above, we conclude that the calibrated ρ -margin loss in Section 4.2.1 is consistent with respect to ℓ_γ for all distributions that satisfy the realizability assumption, i.e., $\mathcal{R}_{\ell_\gamma, \mathcal{H}}^* = 0$.

Theorem 20 *Consider the ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$. Then,*

1. *If $\rho > 1$, then ϕ_ρ is \mathcal{H}_{lin} -consistent wrt ℓ_γ for all distribution P over $\mathcal{X} \times \mathcal{Y}$ that satisfies $\mathcal{R}_{\ell_\gamma, \mathcal{H}_{\text{lin}}}^* = 0$ and there exists $f^* \in \mathcal{H}_{\text{lin}}$ such that $\mathcal{R}_\phi(f^*) = \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* < \infty$.*
2. *Given a non-decreasing and continuous function g such that $g(1+\gamma) < G$ and $g(-1-\gamma) > -G$ for some $G \geq 0$. Assume that $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$. Let $\bar{A}(t) = \max_{s \in [-t, t]} g(s) - g(s - \gamma)$ and $\underline{A}(t) = \min_{s \in [-t, t]} g(s) - g(s + \gamma)$ for any $0 \leq t \leq 1$. If for any $0 \leq t \leq 1$, $\phi_\rho(G - g(-t)) = \phi_\rho(g(t) + G)$ and $\min\{\phi_\rho(\bar{A}(t)), \phi_\rho(-\underline{A}(t))\} > \phi_\rho(G - g(-t))$, then ϕ_ρ is \mathcal{H}_g -consistent wrt ℓ_γ for all distribution P over $\mathcal{X} \times \mathcal{Y}$ that satisfies $\mathcal{R}_{\ell_\gamma, \mathcal{H}_g}^* = 0$ and there exists $f^* \in \mathcal{H}_g$ such that $\mathcal{R}_\phi(f^*) = \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* < \infty$.*
3. *If $G > 1 + \gamma$ and $G \geq \rho > \gamma$, then ϕ_ρ is $\mathcal{H}_{\text{relu}}$ -consistent wrt ℓ_γ for all distribution P over $\mathcal{X} \times \mathcal{Y}$ that satisfies $\mathcal{R}_{\ell_\gamma, \mathcal{H}_{\text{relu}}}^* = 0$ and there exists $f^* \in \mathcal{H}_{\text{relu}}$ such that $\mathcal{R}_\phi(f^*) = \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* < \infty$.*

Using Theorem 16 in Section 4.2.2 and Theorem 19, we conclude that the calibrated supremum-based ρ -margin loss in Section 4.2.2 is also consistent wrt ℓ_γ for all distributions that satisfy realizability assumptions.

Theorem 21 *Consider ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$. Let \mathcal{H} be a symmetric hypothesis set containing 0, then the surrogate loss $\tilde{\phi}_\rho(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi_\rho(yf(\mathbf{x}'))$ is \mathcal{H} -consistent with respect to ℓ_γ for all distributions P over $\mathcal{X} \times \mathcal{Y}$ that satisfy: $\mathcal{R}_{\ell_\gamma, \mathcal{H}}^* = 0$ and there exists $f^* \in \mathcal{H}$ such that $\mathcal{R}_\phi(f^*) = \mathcal{R}_{\phi, \mathcal{H}_{\text{all}}}^* < \infty$.*

6. Conclusion

We presented a careful analysis of the \mathcal{H} -calibration of surrogate losses, including a series of negative results for surrogate losses commonly used in practice, as well as a number of positive results for surrogate losses that we prove additionally to be \mathcal{H} -consistent, provided that some other natural conditions hold. Our result significantly extend previously known results and provide a solid guidance for the design of algorithms for adversarial robustness with theoretical guarantees. Moreover, several of our proof techniques for calibration and consistency can further be relevant to the analysis of other loss functions.

Acknowledgments

We warmly thank our colleague Natalie Frank for discussions and our previous joint work on this topic.

References

Pranjal Awasthi, Natalie Frank, and Mehryar Mohri. Adversarial learning guarantees for linear hypotheses and neural networks. In *International Conference on Machine Learning*, pages 431–441, 2020.

- Pranjal Awasthi, Natalie Frank, Anqi Mao, Mehryar Mohri, and Yutao Zhong. Calibration and consistency of adversarial surrogate losses. *arXiv preprint arXiv:2104.09658*, 2021.
- Han Bao, Clayton Scott, and Masashi Sugiyama. Calibrated surrogate losses for adversarially robust classification. In *Conference on Learning Theory*, pages 408–451, 2020.
- Stephen P. Boyd and Lieven Vandenbergh. *Convex Optimization*. Cambridge University Press, 2014.
- Nicholas Carlini and David Wagner. Towards evaluating the robustness of neural networks. In *IEEE Symposium on Security and Privacy (SP)*, pages 39–57, 2017.
- Ian J Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. *arXiv preprint arXiv:1412.6572*, 2014.
- Alex Krizhevsky, Ilya Sutskever, and Geoffrey E. Hinton. Imagenet classification with deep convolutional neural networks. In *Advances in Neural Information Processing Systems*, pages 1097–1105, 2012.
- Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. *arXiv preprint arXiv:1706.06083*, 2017.
- Mehryar Mohri, Afshin Rostamizadeh, and Ameet Talwalkar. *Foundations of Machine Learning*. MIT Press, second edition, 2018.
- Ali Shafahi, Mahyar Najibi, Mohammad Amin Ghiasi, Zheng Xu, John Dickerson, Christoph Studer, Larry S Davis, Gavin Taylor, and Tom Goldstein. Adversarial training for free! In *Advances in Neural Information Processing Systems*, pages 3353–3364, 2019.
- Ingo Steinwart. How to compare different loss functions and their risks. *Constructive Approximation*, 26(2):225–287, 2007.
- Ilya Sutskever, Oriol Vinyals, and Quoc V. Le. Sequence to sequence learning with neural networks. In *Advances in Neural Information Processing Systems*, pages 3104–3112, 2014.
- Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. *arXiv preprint arXiv:1312.6199*, 2013.
- Dimitris Tsipras, Shibani Santurkar, Logan Engstrom, Alexander Turner, and Aleksander Madry. Robustness may be at odds with accuracy. *arXiv preprint arXiv:1805.12152*, 2018.
- Eric Wong, Leslie Rice, and J Zico Kolter. Fast is better than free: Revisiting adversarial training. *arXiv preprint arXiv:2001.03994*, 2020.
- Dong Yin, Kannan Ramchandran, and Peter L. Bartlett. Rademacher complexity for adversarially robust generalization. In *International Conference of Machine Learning*, pages 7085–7094, 2019.

Contents of Appendix

A	Deferred Proofs	13
A.1	Proof of Theorem 8 and Theorem 10	13
A.2	Property of $\bar{\mathcal{C}}_\phi(t, \eta)$	16
A.3	Proof of Theorem 12 and Theorem 16	18
A.4	Proof of Theorem 13	26

Appendix A. Deferred Proofs

For convenience, let $\Delta\mathcal{C}_{\ell,\mathcal{H}}(f, \mathbf{x}, \eta) := \mathcal{C}_\ell(f, \mathbf{x}, \eta) - \mathcal{C}_{\ell,\mathcal{H}}^*(\mathbf{x}, \eta)$, $\underline{M}(f, \mathbf{x}, \gamma) := \inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} f(\mathbf{x}')$ and $\overline{M}(f, \mathbf{x}, \gamma) := -\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} -f(\mathbf{x}') = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} f(\mathbf{x}')$.

A.1. Proof of Theorem 8 and Theorem 10

We first characterize the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses (ℓ, ℓ_γ) at $\eta = \frac{1}{2}$, $\epsilon = \frac{1}{2}$ and distinguishing $\mathbf{x}_0 \in \mathcal{X}$ given a hypothesis set \mathcal{H} containing 0 which is regular for adversarial calibration.

Lemma 22 *Let \mathcal{H} be a hypothesis set containing 0 that is regular for adversarial calibration. For distinguishing $\mathbf{x}_0 \in \mathcal{X}$, the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses (ℓ, ℓ_γ) satisfies*

$$\delta_{\max}\left(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}\right) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta\mathcal{C}_{\ell,\mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}).$$

Proof By the definition of inner risk (4) and adversarial 0-1 loss ℓ_γ (10), the inner ℓ_γ -risk is

$$\begin{aligned} \mathcal{C}_{\ell_\gamma}(f, \mathbf{x}, \eta) &= \eta \mathbb{1}_{\{\underline{M}(f, \mathbf{x}, \gamma) \leq 0\}} + (1 - \eta) \mathbb{1}_{\{\overline{M}(f, \mathbf{x}, \gamma) \geq 0\}} \\ &= \begin{cases} 1 & \text{if } \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \\ \eta & \text{if } \overline{M}(f, \mathbf{x}, \gamma) < 0, \\ 1 - \eta & \text{if } \underline{M}(f, \mathbf{x}, \gamma) > 0. \end{cases} \end{aligned}$$

For distinguishing \mathbf{x}_0 and $\eta \in [0, 1]$, $\{f \in \mathcal{H} : \overline{M}(f, \mathbf{x}_0, \gamma) < 0\}$ and $\{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}_0, \gamma) > 0\}$ are not empty sets. Thus

$$\mathcal{C}_{\ell_\gamma, \mathcal{H}}^*(\mathbf{x}_0, \eta) = \inf_{f \in \mathcal{H}} \mathcal{C}_{\ell_\gamma}(f, \mathbf{x}_0, \eta) = \min\{\eta, 1 - \eta\}.$$

Since \mathcal{H} contains 0, $\{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)\}$ is also not empty set. Note for $f \in \{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)\}$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \eta) = \max\{\eta, 1 - \eta\}$; for $f \in \{f \in \mathcal{H} : \overline{M}(f, \mathbf{x}_0, \gamma) < 0\}$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \eta) = \eta - \min\{\eta, 1 - \eta\} = \max\{0, 2\eta - 1\} = |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}_0, \gamma)) \leq 0}$ since $\underline{M}(f, \mathbf{x}_0, \gamma) \leq \overline{M}(f, \mathbf{x}_0, \gamma) < 0$; for $f \in \{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}_0, \gamma) > 0\}$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \eta) = (1 - \eta) - \min\{\eta, 1 - \eta\} = \max\{0, 1 - 2\eta\} = |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}_0, \gamma)) \leq 0}$. Therefore,

$$\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \eta) = \begin{cases} \max\{\eta, 1 - \eta\} & \text{if } \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma), \\ |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}_0, \gamma)) \leq 0} & \text{if } \underline{M}(f, \mathbf{x}_0, \gamma) > 0 \text{ or } \overline{M}(f, \mathbf{x}_0, \gamma) < 0. \end{cases}$$

By (7), for a fixed $\eta \in [0, 1]$ and $\mathbf{x} \in \mathcal{X}$, the calibration function of losses (ℓ, ℓ_γ) is

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}} \{\Delta\mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) \mid \Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) \geq \epsilon\}.$$

Observe that for all $\eta \in [0, 1]$,

$$\max\{\eta, 1 - \eta\} = \frac{1}{2}[(1 - \eta) + \eta + |(1 - \eta) - \eta|] = \frac{1}{2}[1 + |2\eta - 1|] \geq |2\eta - 1|. \quad (12)$$

For distinguishing \mathbf{x}_0 , $\eta = \frac{1}{2}$ and $\epsilon = \frac{1}{2}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}) \geq \frac{1}{2}$ if and only if $\underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)$ since $|2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}$. Therefore,

$$\delta_{\max}\left(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}\right) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}).$$

■

Theorem 8 *Let \mathcal{H} be a hypothesis set containing 0 that is regular for adversarial calibration. If a margin-based loss $\phi: \mathbb{R} \rightarrow \mathbb{R}_+$ is convex, then it is not \mathcal{H} -calibrated with respect to ℓ_γ .*

Proof By Lemma 22, for distinguishing $\mathbf{x}_0 \in \mathcal{X}$, the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses (ϕ, ℓ_γ) satisfies

$$\delta_{\max}\left(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}\right) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta \mathcal{C}_{\phi, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}).$$

Suppose that ϕ is \mathcal{H} -calibrated with respect to ℓ_γ . By Proposition 5, ϕ is \mathcal{H} -calibrated with respect to ℓ_γ if and only if its calibration function δ_{\max} satisfies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$. In particular, the condition requires $\delta_{\max}\left(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}\right) > 0$, that is,

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta \mathcal{C}_{\phi, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}) > 0,$$

which is equivalent to

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) > \inf_{f \in \mathcal{H}} \mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}), \quad (13)$$

By the definition of inner risk (4),

$$\mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) = \frac{1}{2}(\phi(f(\mathbf{x}_0)) + \phi(-f(\mathbf{x}_0))). \quad (14)$$

Since ϕ is convex, by Jensen's inequality, for any $f \in \mathcal{H}$, the following holds:

$$\mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) \geq \phi\left(\frac{1}{2}f(\mathbf{x}_0) - \frac{1}{2}f(\mathbf{x}_0)\right) = \phi(0).$$

For $f = 0$, we have $\underline{M}(f, \mathbf{x}_0, \gamma) = \overline{M}(f, \mathbf{x}_0, \gamma) = 0$ and by (14),

$$\mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) = \frac{1}{2}(\phi(0) + \phi(0)) = \phi(0).$$

Moreover, when $\underline{M}(f, \mathbf{x}_0, \gamma) = \overline{M}(f, \mathbf{x}_0, \gamma) = 0$, $\underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)$ is satisfied. Thus

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) = \inf_{f \in \mathcal{H}} \mathcal{C}_\phi(f, \mathbf{x}_0, \frac{1}{2}) = \phi(0),$$

where the minimum can be achieved by $f = 0$, contradicting (13). Therefore, ϕ is not \mathcal{H} -calibrated with respect to ℓ_γ . ■

Theorem 10 *Let \mathcal{H} be a hypothesis set containing 0 that is regular for adversarial calibration. If a margin-based loss ϕ is convex and non-increasing, then the surrogate loss defined by $\tilde{\phi}(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi(yf(\mathbf{x}'))$ is not \mathcal{H} -calibrated with respect to ℓ_γ .*

Proof By Lemma 22, for distinguishing $\mathbf{x}_0 \in \mathcal{X}$, the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses $(\tilde{\phi}, \ell_\gamma)$ satisfies

$$\delta_{\max}\left(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}\right) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta \mathcal{C}_{\tilde{\phi}, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}).$$

Suppose that $\tilde{\phi}$ is \mathcal{H} -calibrated with respect to ℓ_γ . By Proposition 5, $\tilde{\phi}$ is \mathcal{H} -calibrated with respect to ℓ_γ if and only if its calibration function δ_{\max} satisfies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$. In particular, the condition requires $\delta_{\max}(\frac{1}{2}, \mathbf{x}_0, \frac{1}{2}) > 0$, that is,

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \Delta \mathcal{C}_{\tilde{\phi}, \mathcal{H}}(f, \mathbf{x}_0, \frac{1}{2}) > 0,$$

which is equivalent to

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) > \inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}), \quad (15)$$

As shown by Awasthi et al. (2020), $\tilde{\phi}$ has the equivalent form

$$\tilde{\phi}(f, \mathbf{x}, y) = \phi\left(\inf_{\|\mathbf{x}' - \mathbf{x}\| \leq \gamma} (yf(\mathbf{x}'))\right).$$

By the definition of inner risk (4),

$$\mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) = \frac{1}{2}(\phi(\underline{M}(f, \mathbf{x}_0, \gamma)) + \phi(-\overline{M}(f, \mathbf{x}_0, \gamma))). \quad (16)$$

Since ϕ is convex, by Jensen's inequality, for any $f \in \mathcal{H}$, the following holds:

$$\mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) \geq \phi\left(\frac{1}{2}\underline{M}(f, \mathbf{x}_0, \gamma) - \frac{1}{2}\overline{M}(f, \mathbf{x}_0, \gamma)\right) = \phi\left(\frac{1}{2}(\underline{M}(f, \mathbf{x}_0, \gamma) - \overline{M}(f, \mathbf{x}_0, \gamma))\right) \geq \phi(0),$$

where the last inequality used the fact that

$$\frac{1}{2}(\underline{M}(f, \mathbf{x}_0, \gamma) - \overline{M}(f, \mathbf{x}_0, \gamma)) \leq 0$$

and ϕ is non-increasing. For $f = 0$, we have $\underline{M}(f, \mathbf{x}_0, \gamma) = \overline{M}(f, \mathbf{x}_0, \gamma) = 0$ and by (16),

$$\mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) = \frac{1}{2}(\phi(0) + \phi(0)) = \phi(0).$$

Moreover, when $\underline{M}(f, \mathbf{x}_0, \gamma) = \overline{M}(f, \mathbf{x}_0, \gamma) = 0$, $\underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)$ is satisfied. Thus

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}_0, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}_0, \gamma)} \mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) = \inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}}(f, \mathbf{x}_0, \frac{1}{2}) = \phi(0),$$

where the minimum can be achieved by $f = 0$, contradicting (15). Therefore, $\tilde{\phi}$ is not \mathcal{H} -calibrated with respect to ℓ_γ . ■

A.2. Property of $\bar{\mathcal{C}}_\phi(t, \eta)$

For a margin-based loss ϕ , denote $\bar{\mathcal{C}}_\phi(t, \eta) := \eta\phi(t) + (1-\eta)\phi(-t)$ for any $\eta \in [0, 1]$ and $t \in \mathbb{R}$. In this section, we characterize the property of $\bar{\mathcal{C}}_\phi(t, \eta)$ when ϕ is bounded, continuous, non-increasing and quasi-concave even, which would be useful in the proof of Theorem 12 and Theorem 13. Without loss of generality, assume that g is continuous, non-decreasing and satisfies $g(-1 - \gamma) + G > 0$, $g(1 + \gamma) - G < 0$.

Lemma 23 *Let ϕ be a margin-based loss. If ϕ is bounded, continuous, non-increasing, quasi-concave even, then*

1. $\bar{\mathcal{C}}_\phi(t, \eta)$ is quasi-concave in $t \in \mathbb{R}$ for all $\eta \in [0, 1]$.
2. $\bar{\mathcal{C}}_\phi(t, \frac{1}{2})$ is even and non-increasing in t when $t \geq 0$.
3. For $l, u \in \mathbb{R} (l \leq u)$, $\inf_{t \in [l, u]} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(l, \eta), \bar{\mathcal{C}}_\phi(u, \eta)\}$ for all $\eta \in [0, 1]$.
4. For all $\eta \in (\frac{1}{2}, 1]$, $\bar{\mathcal{C}}_\phi(t, \eta)$ is non-increasing in t when $t \geq 0$.
5. For all $\eta \in [0, \frac{1}{2})$, $\bar{\mathcal{C}}_\phi(t, \eta)$ is non-decreasing in t when $t \leq 0$.
6. If $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$, then, for all $\eta \in (\frac{1}{2}, 1]$ and any $\gamma < t \leq 1$, $\bar{\mathcal{C}}_\phi(-t, \eta) > \bar{\mathcal{C}}_\phi(t, \eta)$.
7. If $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$, then, for all $\eta \in [0, \frac{1}{2})$ and any $\gamma < t \leq 1$, $\bar{\mathcal{C}}_\phi(-t, \eta) < \bar{\mathcal{C}}_\phi(t, \eta)$.
8. If $\phi(g(-t) - G) > \phi(G - g(-t))$, $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$, then, for all $\eta \in (\frac{1}{2}, 1]$ and any $0 \leq t \leq 1$, $\bar{\mathcal{C}}_\phi(g(-t) - G, \eta) > \bar{\mathcal{C}}_\phi(g(t) + G, \eta)$.
9. If $\phi(g(-t) - G) > \phi(G - g(-t))$, $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$, then, for any $0 \leq t \leq 1$, $\bar{\mathcal{C}}_\phi(g(-t) - G, \eta) < \bar{\mathcal{C}}_\phi(g(t) + G, \eta)$ for all $\eta \in [0, \frac{1}{2})$ if and only if $\phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G)$.

Proof Part 1,2,4 of Lemma 23 are stated in (Bao et al., 2020, Lemma 13). Part 3 is a corollary of Part 1 by the characterization of continuous and quasi-convex functions in (Boyd and Vandenberghe, 2014).

Consider Part 5. For $\eta \in [0, \frac{1}{2})$, and $t_1, t_2 \leq 0$. Suppose that $t_1 < t_2$, then

$$\begin{aligned} & \phi(t_1) - \phi(-t_1) - \phi(t_2) + \phi(-t_2) \\ & \geq \phi(t_2) - \phi(-t_2) - \phi(t_2) + \phi(-t_2) \\ & = 0 \end{aligned}$$

since ϕ is non-increasing. By Part 2 of Lemma 23, $\phi(t) + \phi(-t)$ is non-decreasing in t when $t \leq 0$. Therefore, for $\eta \in [0, \frac{1}{2})$,

$$\begin{aligned} & \bar{\mathcal{C}}_\phi(t_1, \eta) - \bar{\mathcal{C}}_\phi(t_2, \eta) \\ &= (\phi(t_1) - \phi(-t_1) - \phi(t_2) + \phi(-t_2))\eta + \phi(-t_1) - \phi(-t_2) \\ &\leq (\phi(t_1) - \phi(-t_1) - \phi(t_2) + \phi(-t_2))\frac{1}{2} + \phi(-t_1) - \phi(-t_2) \\ &= \frac{1}{2}(\phi(t_1) + \phi(-t_1) - \phi(t_2) - \phi(-t_2)) \\ &\leq 0. \end{aligned}$$

Consider Part 6, For $\eta \in (\frac{1}{2}, 1]$ and any $\gamma < t \leq 1$,

$$\begin{aligned} \bar{\mathcal{C}}_\phi(-t, \eta) - \bar{\mathcal{C}}_\phi(t, \eta) &= \eta\phi(-t) + (1-\eta)\phi(t) - \eta\phi(t) - (1-\eta)\phi(-t) \\ &= (2\eta - 1)[\phi(-t) - \phi(t)] > 0 \end{aligned}$$

since $\eta > \frac{1}{2}$ and $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$.

Consider Part 7, For $\eta \in [0, \frac{1}{2})$ and any $\gamma < t \leq 1$,

$$\begin{aligned} \bar{\mathcal{C}}_\phi(t, \eta) - \bar{\mathcal{C}}_\phi(-t, \eta) &= \eta\phi(t) + (1-\eta)\phi(-t) - \eta\phi(-t) - (1-\eta)\phi(t) \\ &= (1-2\eta)[\phi(-t) - \phi(t)] > 0 \end{aligned}$$

since $\eta < \frac{1}{2}$ and $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$.

Consider Part 8. For $\eta \in (\frac{1}{2}, 1]$ and any $0 \leq t \leq 1$,

$$\begin{aligned} & \bar{\mathcal{C}}_\phi(g(-t) - G, \eta) - \bar{\mathcal{C}}_\phi(g(t) + G, \eta) \\ &\geq \bar{\mathcal{C}}_\phi(g(-t) - G, \eta) - \bar{\mathcal{C}}_\phi(G - g(-t), \eta) \quad (g(-t) + g(t) \geq 0, \text{ Part 4 of Lemma 23}) \\ &= (2\eta - 1)[\phi(g(-t) - G) - \phi(G - g(-t))] \\ &> 0 \quad (\phi(g(-t) - G) > \phi(G - g(-t))) \end{aligned}$$

Consider Part 9. Since ϕ is non-increasing, for any $0 \leq t \leq 1$,

$$\begin{aligned} & \phi(g(-t) - G) - \phi(G - g(-t)) + \phi(-g(t) - G) - \phi(g(t) + G) \\ &\geq \phi(g(-t) - G) - \phi(G - g(-t)) + \phi(g(t) + G) - \phi(g(t) + G) \quad (g(t) + G > 0) \\ &= \phi(g(-t) - G) - \phi(G - g(-t)) \\ &> 0 \quad (\phi(g(-t) - G) > \phi(G - g(-t))) \end{aligned}$$

\Leftarrow : Suppose $\phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G)$, then for $\eta \in [0, \frac{1}{2})$,

$$\begin{aligned} & \bar{\mathcal{C}}_\phi(g(-t) - G, \eta) - \bar{\mathcal{C}}_\phi(g(t) + G, \eta) \\ &= (\phi(g(-t) - G) - \phi(G - g(-t)) + \phi(-g(t) - G) - \phi(g(t) + G))\eta \\ &\quad + \phi(G - g(-t)) - \phi(-g(t) - G) \\ &< (\phi(g(-t) - G) - \phi(G - g(-t)) + \phi(-g(t) - G) - \phi(g(t) + G))\frac{1}{2} \\ &\quad + \phi(G - g(-t)) - \phi(-g(t) - G) \\ &= \frac{1}{2}(\phi(G - g(-t)) + \phi(g(-t) - G) - \phi(g(t) + G) - \phi(-g(t) - G)) \\ &= 0. \end{aligned}$$

\implies : Suppose $\bar{\mathcal{C}}_\phi(g(-t) - G, \eta) < \bar{\mathcal{C}}_\phi(g(t) + G, \eta)$ for $\eta \in [0, \frac{1}{2})$, then

$$\begin{aligned} & \bar{\mathcal{C}}_\phi(g(-t) - G, \eta) - \bar{\mathcal{C}}_\phi(g(t) + G, \eta) \\ &= (\phi(g(-t) - G) - \phi(G - g(-t)) + \phi(-g(t) - G) - \phi(g(t) + G))\eta \\ & \quad + \phi(G - g(-t)) - \phi(-g(t) - G) \\ & < 0 \end{aligned}$$

for $\eta \in [0, \frac{1}{2})$. By taking $\eta \rightarrow \frac{1}{2}$, we have

$$\begin{aligned} & \frac{1}{2}(\phi(G - g(-t)) + \phi(g(-t) - G) - \phi(g(t) + G) - \phi(-g(t) - G)) \\ &= (\phi(g(-t) - G) - \phi(G - g(-t)) + \phi(-g(t) - G) - \phi(g(t) + G))\frac{1}{2} \\ & \quad + \phi(G - g(-t)) - \phi(-g(t) - G) \\ & \leq 0. \end{aligned}$$

By Part 2 of Lemma 23, we have

$$\begin{aligned} & \phi(G - g(-t)) + \phi(g(-t) - G) - \phi(g(t) + G) - \phi(-g(t) - G) \\ & \geq \phi(g(t) + G) + \phi(-g(t) - G) - \phi(g(t) + G) - \phi(-g(t) - G) \quad (g(-t) + g(t) \geq 0) \\ & = 0. \end{aligned}$$

Therefore, $\phi(G - g(-t)) + \phi(g(-t) - G) - \phi(g(t) + G) - \phi(-g(t) - G) = 0$, i.e., $\phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G)$. \blacksquare

A.3. Proof of Theorem 12 and Theorem 16

We will make use of general form (10) of the adversarial 0/1 loss:

$$\ell_\gamma(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \mathbb{1}_{yf(\mathbf{x}') \leq 0} = \mathbb{1}_{\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} yf(\mathbf{x}') \leq 0}.$$

Next, we first characterize the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses (ℓ, ℓ_γ) given a symmetric hypothesis set \mathcal{H} containing 0.

Lemma 24 *Let \mathcal{H} be a symmetric hypothesis set containing 0. For a surrogate loss ℓ , the calibration function $\delta_{\max}(\epsilon, \mathbf{x}, \eta)$ of losses (ℓ, ℓ_γ) is*

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \begin{cases} +\infty & \text{if } \mathbf{x} \in \mathcal{X}_1 \text{ or } \mathbf{x} \in \mathcal{X}_2, \epsilon > \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) & \text{if } \mathbf{x} \in \mathcal{X}_2, |2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma) \text{ or } (2\eta - 1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) & \text{if } \mathbf{x} \in \mathcal{X}_2, \epsilon \leq |2\eta - 1|, \end{cases}$$

where $\mathcal{X}_1 = \{\mathbf{x} \in \mathcal{X} : \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \forall f \in \mathcal{H}\}$, $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\}$ and $\mathcal{X} = \mathcal{X}_1 \cup \mathcal{X}_2$, $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$.

Proof By the definition of inner risk (4) and adversarial 0-1 loss ℓ_γ (10), the inner ℓ_γ -risk is

$$\begin{aligned} \mathcal{C}_{\ell_\gamma}(f, \mathbf{x}, \eta) &= \eta \mathbb{1}_{\{\underline{M}(f, \mathbf{x}, \gamma) \leq 0\}} + (1 - \eta) \mathbb{1}_{\{\overline{M}(f, \mathbf{x}, \gamma) \geq 0\}} \\ &= \begin{cases} 1 & \text{if } \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \\ \eta & \text{if } \overline{M}(f, \mathbf{x}, \gamma) < 0, \\ 1 - \eta & \text{if } \underline{M}(f, \mathbf{x}, \gamma) > 0. \end{cases} \end{aligned}$$

Let $\mathcal{X}_1 = \{\mathbf{x} \in \mathcal{X} : \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \forall f \in \mathcal{H}\}$, $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\}$. It is obvious that $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. Since \mathcal{H} is symmetric, for any $\mathbf{x} \in \mathcal{X}$, either there exists $f' \in \mathcal{H}$ such that $\underline{M}(f', \mathbf{x}, \gamma) > 0$ and $\overline{M}(-f', \mathbf{x}, \gamma) < 0$, or $\underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)$ for any $f \in \mathcal{H}$. Thus $\mathcal{X} = \mathcal{X}_1 \cup \mathcal{X}_2$. Note when $\mathbf{x} \in \mathcal{X}_1$, $\{f \in \mathcal{H} : \overline{M}(f, \mathbf{x}, \gamma) < 0\}$ and $\{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}, \gamma) > 0\}$ are both empty sets. Therefore, the minimal inner ℓ_γ -risk is

$$\mathcal{C}_{\ell_\gamma, \mathcal{H}}^*(\mathbf{x}, \eta) = \begin{cases} 1, & \mathbf{x} \in \mathcal{X}_1, \\ \min\{\eta, 1 - \eta\}, & \mathbf{x} \in \mathcal{X}_2. \end{cases}$$

Since $0 \in \mathcal{H}$, for any $\mathbf{x} \in \mathcal{X}$, $\{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)\}$ is not empty set. Note when $\mathbf{x} \in \mathcal{X}_1$, $\mathcal{C}_{\ell_\gamma}(f, \mathbf{x}, \eta) = 1$ for any $f \in \mathcal{H}$, thus $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = 0$. When $\mathbf{x} \in \mathcal{X}_2$, for $f \in \{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)\}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = 1 - \min\{\eta, 1 - \eta\} = \max\{\eta, 1 - \eta\}$; for $f \in \{f \in \mathcal{H} : \overline{M}(f, \mathbf{x}, \gamma) < 0\}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = \eta - \min\{\eta, 1 - \eta\} = \max\{0, 2\eta - 1\} = |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0}$ since $\underline{M}(f, \mathbf{x}, \gamma) \leq \overline{M}(f, \mathbf{x}, \gamma) < 0$; for $f \in \{f \in \mathcal{H} : \underline{M}(f, \mathbf{x}, \gamma) > 0\}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = 1 - \eta - \min\{\eta, 1 - \eta\} = \max\{0, 1 - 2\eta\} = |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0}$ since $\underline{M}(f, \mathbf{x}, \gamma) > 0$. Therefore,

$$\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = \begin{cases} \max\{\eta, 1 - \eta\} & \text{if } \mathbf{x} \in \mathcal{X}_2, \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \\ |2\eta - 1| \mathbb{1}_{(2\eta-1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} & \text{if } \mathbf{x} \in \mathcal{X}_2, \underline{M}(f, \mathbf{x}, \gamma) > 0 \text{ or } \overline{M}(f, \mathbf{x}, \gamma) < 0, \\ 0 & \text{if } \mathbf{x} \in \mathcal{X}_1. \end{cases} \quad (17)$$

By (7), for a fixed $\eta \in [0, 1]$ and $\mathbf{x} \in \mathcal{X}$, the calibration function of losses (ℓ, ℓ_γ) is

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}} \left\{ \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) \mid \Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) \geq \epsilon \right\}$$

If $\mathbf{x} \in \mathcal{X}_1$, then for all $f \in \mathcal{H}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) = 0 < \epsilon$, which implies that $\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \infty$. Next we consider case where $\mathbf{x} \in \mathcal{X}_2$. By the observation (12), if $\epsilon > \max\{\eta, 1 - \eta\}$, then for all $f \in \mathcal{H}$, $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) < \epsilon$, which implies that $\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \infty$; if $|2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}$, then $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) \geq \epsilon$ if and only if $\underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)$, which leads to

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta);$$

if $\epsilon \leq |2\eta - 1|$, then $\Delta \mathcal{C}_{\ell_\gamma, \mathcal{H}}(f, \mathbf{x}, \eta) \geq \epsilon$ if and only if $\underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)$ or $(2\eta - 1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0$, which leads to

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma) \text{ or } (2\eta-1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta).$$

■

We then give the equivalent conditions of calibration based on inner ℓ -risk and \mathcal{H} .

Lemma 25 *Let \mathcal{H} be a symmetric hypothesis set containing 0 and ℓ be a surrogate loss function. If $\mathcal{X}_2 = \emptyset$, any loss ℓ is \mathcal{H} -calibrated with respect to ℓ_γ . If $\mathcal{X}_2 \neq \emptyset$, then ℓ is \mathcal{H} -calibrated with respect to ℓ_γ if and only if for any $\mathbf{x} \in \mathcal{X}_2$,*

$$\begin{aligned} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}) &> \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}), \text{ and} \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned}$$

where $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\}$.

Proof Let δ_{\max} be the calibration function of (ℓ, ℓ_γ) given hypothesis set \mathcal{H} . By Lemma 24,

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \begin{cases} +\infty & \text{if } \mathbf{x} \in \mathcal{X}_1 \text{ or } \mathbf{x} \in \mathcal{X}_2, \epsilon > \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) & \text{if } \mathbf{x} \in \mathcal{X}_2, |2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma) \text{ or } (2\eta - 1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) & \text{if } \mathbf{x} \in \mathcal{X}_2, \epsilon \leq |2\eta - 1|, \end{cases}$$

where $\mathcal{X}_1 = \{\mathbf{x} \in \mathcal{X} : \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma), \forall f \in \mathcal{H}\}$, $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\}$ and $\mathcal{X} = \mathcal{X}_1 \cup \mathcal{X}_2$, $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$. By Proposition 5, ℓ is \mathcal{H} -calibrated with respect to ℓ_γ if and only if its calibration function δ_{\max} satisfies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$. Since $\delta(\epsilon, \mathbf{x}, \eta) = \infty > 0$ when $\mathbf{x} \notin \mathcal{X}_2$, any loss ℓ is \mathcal{H} -calibrated with respect to ℓ_γ when $\mathcal{X}_2 = \emptyset$. Furthermore, when $\mathcal{X}_2 \neq \emptyset$, we only need to analyze $\delta(\epsilon, \mathbf{x}, \eta)$ when $\mathbf{x} \in \mathcal{X}_2$. For $\eta = \frac{1}{2}$, we have for any $\mathbf{x} \in \mathcal{X}_2$,

$$\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0 \text{ for all } \epsilon > 0 \Leftrightarrow \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}). \quad (18)$$

For $1 \geq \eta > \frac{1}{2}$, we have $|2\eta - 1| = 2\eta - 1$, $\max\{\eta, 1 - \eta\} = \eta$, and

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma) \text{ or } (2\eta - 1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta).$$

Therefore, $\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0$ for all $\mathbf{x} \in \mathcal{X}_2, \epsilon > 0$ and $\eta \in (\frac{1}{2}, 1]$ if and only if for all $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{cases} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } 2\eta - 1 < \epsilon \leq \eta, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \epsilon \leq 2\eta - 1, \end{cases}$$

for all $\epsilon > 0$, which is equivalent to for all $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{cases} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \epsilon \leq \eta < \frac{\epsilon+1}{2}, \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \frac{\epsilon+1}{2} \leq \eta, \end{cases} \quad (19)$$

for all $\epsilon > 0$. Observe that

$$\begin{aligned} \left\{ \eta \in \left(\frac{1}{2}, 1 \right] \middle| \epsilon \leq \eta < \frac{\epsilon+1}{2}, \epsilon > 0 \right\} &= \left\{ \frac{1}{2} < \eta \leq 1 \right\}, \text{ and} \\ \left\{ \eta \in \left(\frac{1}{2}, 1 \right] \middle| \frac{\epsilon+1}{2} \leq \eta, \epsilon > 0 \right\} &= \left\{ \frac{1}{2} < \eta \leq 1 \right\}, \text{ and} \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &\geq \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta. \end{aligned}$$

Therefore, we reduce the above condition (19) as for all $\mathbf{x} \in \mathcal{X}_2$,

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in \left(\frac{1}{2}, 1 \right]. \quad (20)$$

For $\frac{1}{2} > \eta \geq 0$, we have $|2\eta - 1| = 1 - 2\eta$, $\max\{\eta, 1 - \eta\} = 1 - \eta$, and

$$\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma) \text{ or } (2\eta - 1)(\underline{M}(f, \mathbf{x}, \gamma)) \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}}(f, \mathbf{x}, \eta).$$

Therefore, $\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0$ for all $\mathbf{x} \in \mathcal{X}_2$, $\epsilon > 0$ and $\eta \in [0, \frac{1}{2})$ if and only if for all $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{cases} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } 1 - 2\eta < \epsilon \leq 1 - \eta, \\ \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \epsilon \leq 1 - 2\eta, \end{cases}$$

for all $\epsilon > 0$, which is equivalent to for all $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{cases} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \frac{1-\epsilon}{2} < \eta \leq 1 - \epsilon, \\ \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \eta \leq \frac{1-\epsilon}{2}, \end{cases} \quad (21)$$

for all $\epsilon > 0$. Observe that

$$\begin{aligned} \left\{ \eta \in [0, \frac{1}{2}) \middle| \frac{1-\epsilon}{2} < \eta \leq 1 - \epsilon, \epsilon > 0 \right\} &= \left\{ 0 \leq \eta < \frac{1}{2} \right\}, \text{ and} \\ \left\{ \eta \in [0, \frac{1}{2}) \middle| \eta \leq \frac{1-\epsilon}{2}, \epsilon > 0 \right\} &= \left\{ 0 \leq \eta < \frac{1}{2} \right\}, \text{ and} \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &\geq \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta. \end{aligned}$$

Therefore, we reduce the above condition (21) as for all $\mathbf{x} \in \mathcal{X}_2$,

$$\inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \quad (22)$$

To sum up, by (18), (20) and (22), we conclude the proof. \blacksquare

Since \mathcal{H}_{lin} is a symmetric hypothesis set containing 0, we could make use of Lemma 24 and Lemma 25 for proving Theorem 12.

Theorem 12 *Let a margin-based loss ϕ be bounded, continuous, non-increasing, and quasi-concave even. Assume that $\phi(-t) > \phi(t)$ for any $\gamma < t \leq 1$. Then ϕ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ if and only if for any $\gamma < t \leq 1$,*

$$\phi(\gamma) + \phi(-\gamma) > \phi(t) + \phi(-t). \quad (11)$$

Proof As shown by [Awasthi et al. \(2020\)](#), for $f \in \mathcal{H}_{\text{lin}} = \{\mathbf{x} \rightarrow \mathbf{w} \cdot \mathbf{x} \mid \|\mathbf{w}\| = 1\}$,

$$\begin{aligned} \underline{M}(f, \mathbf{x}, \gamma) &= \inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} f(\mathbf{x}') = \inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} (\mathbf{w} \cdot \mathbf{x}') = \mathbf{w} \cdot \mathbf{x} - \gamma \|\mathbf{w}\| = f(\mathbf{x}) - \gamma, \\ \overline{M}(f, \mathbf{x}, \gamma) &= - \inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} -f(\mathbf{x}') = - \inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} (-\mathbf{w} \cdot \mathbf{x}') = \mathbf{w} \cdot \mathbf{x} + \gamma \|\mathbf{w}\| = f(\mathbf{x}) + \gamma. \end{aligned}$$

Thus for \mathcal{H}_{lin} , $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H}_{\text{lin}} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\} = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H}_{\text{lin}} \text{ such that } f'(\mathbf{x}) > \gamma\} = \{\mathbf{x} : \gamma < \|\mathbf{x}\| \leq 1\}$ since $f(\mathbf{x}) = \mathbf{w} \cdot \mathbf{x} \in [-\|\mathbf{x}\|, \|\mathbf{x}\|]$ when $f \in \mathcal{H}_{\text{lin}}$. Note \mathcal{H}_{lin} is a symmetric hypothesis set containing 0. Therefore, by Lemma 25, ϕ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ if and only if for any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\begin{aligned} \inf_{f \in \mathcal{H}_{\text{lin}}: |f(\mathbf{x})| \leq \gamma} \mathcal{C}_\phi(f, \mathbf{x}, \frac{1}{2}) &> \inf_{f \in \mathcal{H}_{\text{lin}}} \mathcal{C}_\phi(f, \mathbf{x}, \frac{1}{2}), \text{ and} \\ \inf_{f \in \mathcal{H}_{\text{lin}}: f(\mathbf{x}) \leq \gamma} \mathcal{C}_\phi(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_{\text{lin}}} \mathcal{C}_\phi(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{f \in \mathcal{H}_{\text{lin}}: f(\mathbf{x}) \geq -\gamma} \mathcal{C}_\phi(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_{\text{lin}}} \mathcal{C}_\phi(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned} \quad (23)$$

By the definition of inner risk (4), the inner ϕ -risk is

$$\mathcal{C}_\phi(f, \mathbf{x}, \eta) = \eta \phi(f(\mathbf{x})) + (1 - \eta) \phi(-f(\mathbf{x})).$$

Note $f(\mathbf{x}) = \mathbf{w} \cdot \mathbf{x} \in [-\|\mathbf{x}\|, \|\mathbf{x}\|]$ when $f \in \mathcal{H}_{\text{lin}}$. Therefore, (23) is equivalent to for any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\begin{aligned} \inf_{-\gamma \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) &> \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}), \text{ and} \\ \inf_{-\|\mathbf{x}\| \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \eta) &> \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{-\gamma \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) &> \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned} \quad (24)$$

Suppose that ϕ is \mathcal{H}_{lin} -calibrated with respect to ℓ_γ . Since by Part 2 of Lemma 23,

$$\inf_{-\gamma \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) = \bar{\mathcal{C}}_\phi(\gamma, \frac{1}{2}), \quad \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) = \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \frac{1}{2}),$$

we obtain $\phi(\gamma) + \phi(-\gamma) = \bar{\mathcal{C}}_\phi(\gamma, \frac{1}{2}) > \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \frac{1}{2}) > \phi(t) + \phi(-t)$ for any $\gamma < t \leq 1$.

Now for the other direction, assume that $\phi(\gamma) + \phi(-\gamma) > \phi(t) + \phi(-t)$ for any $\gamma < t \leq 1$. For $\eta = \frac{1}{2}$, by Part 2 of Lemma 23, we obtain for any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\inf_{-\gamma \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) = \bar{\mathcal{C}}_\phi(\gamma, \frac{1}{2}) > \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \frac{1}{2}) = \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}).$$

For $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\begin{aligned} \inf_{-\|\mathbf{x}\| \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\bar{\mathcal{C}}_\phi(\gamma, \eta), \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\ \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta), \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\ &= \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta) \quad (\text{Part 6 of Lemma 23}) \end{aligned}$$

Note for $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$, since ϕ is non-increasing,

$$\phi(\gamma) - \phi(-\gamma) - \phi(\|\mathbf{x}\|) + \phi(-\|\mathbf{x}\|) \geq \phi(\|\mathbf{x}\|) - \phi(-\|\mathbf{x}\|) - \phi(\|\mathbf{x}\|) + \phi(-\|\mathbf{x}\|) = 0.$$

Thus

$$\begin{aligned} \bar{\mathcal{C}}_\phi(\gamma, \eta) - \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta) &= \eta\phi(\gamma) + (1 - \eta)\phi(-\gamma) - \eta\phi(\|\mathbf{x}\|) - (1 - \eta)\phi(-\|\mathbf{x}\|) \\ &= (\phi(\gamma) - \phi(-\gamma) - \phi(\|\mathbf{x}\|) + \phi(-\|\mathbf{x}\|))\eta + \phi(-\gamma) - \phi(-\|\mathbf{x}\|) \\ &\geq (\phi(\gamma) - \phi(-\gamma) - \phi(\|\mathbf{x}\|) + \phi(-\|\mathbf{x}\|))\frac{1}{2} + \phi(-\gamma) - \phi(-\|\mathbf{x}\|) \\ &= \frac{1}{2}[\phi(\gamma) + \phi(-\gamma) - \phi(\|\mathbf{x}\|) - \phi(-\|\mathbf{x}\|)] \\ &> 0. \end{aligned}$$

In addition, we have for $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta) > \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta). \quad (\text{Part 6 of Lemma 23})$$

Therefore for $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\inf_{-\|\mathbf{x}\| \leq t \leq \gamma} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(\gamma, \eta), \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta)\} > \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta) = \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta).$$

For $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\begin{aligned} \inf_{-\gamma \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\bar{\mathcal{C}}_\phi(-\gamma, \eta), \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\ \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta), \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\ &= \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta) \quad (\text{Part 7 of Lemma 23}) \end{aligned}$$

Note for $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$, since ϕ is non-increasing,

$$\phi(-\gamma) - \phi(\gamma) - \phi(-\|\mathbf{x}\|) + \phi(\|\mathbf{x}\|) \leq \phi(-\|\mathbf{x}\|) - \phi(\|\mathbf{x}\|) - \phi(-\|\mathbf{x}\|) + \phi(\|\mathbf{x}\|) = 0.$$

Thus

$$\begin{aligned} \bar{\mathcal{C}}_\phi(-\gamma, \eta) - \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta) &= \eta\phi(-\gamma) + (1 - \eta)\phi(\gamma) - \eta\phi(-\|\mathbf{x}\|) - (1 - \eta)\phi(\|\mathbf{x}\|) \\ &= (\phi(-\gamma) - \phi(\gamma) - \phi(-\|\mathbf{x}\|) + \phi(\|\mathbf{x}\|))\eta + \phi(\gamma) - \phi(\|\mathbf{x}\|) \\ &\geq (\phi(-\gamma) - \phi(\gamma) - \phi(-\|\mathbf{x}\|) + \phi(\|\mathbf{x}\|))\frac{1}{2} + \phi(\gamma) - \phi(\|\mathbf{x}\|) \\ &= \frac{1}{2}[\phi(\gamma) + \phi(-\gamma) - \phi(\|\mathbf{x}\|) - \phi(-\|\mathbf{x}\|)] \\ &> 0. \end{aligned}$$

In addition, we have for $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta) > \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta). \quad (\text{Part 7 of Lemma 23})$$

Therefore for $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$ such that $\gamma < \|\mathbf{x}\| \leq 1$,

$$\inf_{-\gamma \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(-\gamma, \eta), \bar{\mathcal{C}}_\phi(\|\mathbf{x}\|, \eta)\} > \bar{\mathcal{C}}_\phi(-\|\mathbf{x}\|, \eta) = \inf_{-\|\mathbf{x}\| \leq t \leq \|\mathbf{x}\|} \bar{\mathcal{C}}_\phi(t, \eta).$$

■

Theorem 16 Consider ρ -margin loss $\phi_\rho(t) = \min\{1, \max\{0, 1 - \frac{t}{\rho}\}\}$, $\rho > 0$. Let \mathcal{H} be a symmetric hypothesis set containing 0, then the surrogate loss $\tilde{\phi}_\rho(f, \mathbf{x}, y) = \sup_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} \phi_\rho(yf(\mathbf{x}'))$ is \mathcal{H} -calibrated with respect to ℓ_γ .

Proof By Lemma 25, if $\mathcal{X}_2 = \emptyset$, $\tilde{\phi}_\rho$ is \mathcal{H} -calibrated with respect to ℓ_γ . Next consider the case where $\mathcal{X}_2 \neq \emptyset$. By Lemma 25, $\tilde{\phi}_\rho$ is \mathcal{H} -calibrated with respect to ℓ_γ if and only if for all $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{aligned} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \frac{1}{2}) &> \inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \frac{1}{2}), \text{ and} \\ \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}), \end{aligned}$$

where $\mathcal{X}_2 = \{\mathbf{x} \in \mathcal{X} : \text{there exists } f' \in \mathcal{H} \text{ such that } \underline{M}(f', \mathbf{x}, \gamma) > 0\}$. As shown by Awasthi et al. (2020), $\tilde{\phi}_\rho$ has the equivalent form

$$\tilde{\phi}_\rho(f, \mathbf{x}, y) = \phi_\rho\left(\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} (yf(\mathbf{x}'))\right).$$

Thus by the definition of inner risk (4), the inner $\tilde{\phi}_\rho$ -risk is

$$\mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) = \eta \phi_\rho(\underline{M}(f, \mathbf{x}, \gamma)) + (1 - \eta) \phi_\rho(-\overline{M}(f, \mathbf{x}, \gamma)).$$

For any $\mathbf{x} \in \mathcal{X}_2$, let $M_{\mathbf{x}} = \sup_{f \in \mathcal{H}} \underline{M}(f, \mathbf{x}, \gamma) > 0$. Since \mathcal{H} is symmetric, we have $-M_{\mathbf{x}} = \inf_{f \in \mathcal{H}} \overline{M}(f, \mathbf{x}, \gamma) < 0$. Since ϕ_ρ is continuous, for any $\mathbf{x} \in \mathcal{X}_2$ and $\epsilon > 0$, there exists $f_{\mathbf{x}}^\epsilon \in \mathcal{H}$ such that $\phi_\rho(\underline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) < \phi_\rho(M_{\mathbf{x}}) + \epsilon$ and $\overline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma) \geq \underline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma) > 0$, $\underline{M}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma) \leq \overline{M}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma) = -\underline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma) < 0$. Next we analyze three cases:

- When $\eta = \frac{1}{2}$, since ϕ_ρ is non-increasing,

$$\begin{aligned} &\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \frac{1}{2}) \\ &= \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \frac{1}{2} \phi_\rho(\underline{M}(f, \mathbf{x}, \gamma)) + \frac{1}{2} \phi_\rho(-\overline{M}(f, \mathbf{x}, \gamma)) \\ &\geq \frac{1}{2} \phi_\rho(0) + \frac{1}{2} \phi_\rho(0) = \phi_\rho(0) = 1. \end{aligned}$$

For any $\mathbf{x} \in \mathcal{X}_2$, there exists $f' \in \mathcal{H}$ such that $\underline{M}(f', \mathbf{x}, \gamma) > 0$ and $-\overline{M}(f', \mathbf{x}, \gamma) \leq -\underline{M}(f', \mathbf{x}, \gamma) < 0$, we obtain

$$\mathcal{C}_{\tilde{\phi}_\rho}(f', \mathbf{x}, \frac{1}{2}) = \frac{1}{2}\phi_\rho(\underline{M}(f', \mathbf{x}, \gamma)) + \frac{1}{2}\phi_\rho(-\overline{M}(f', \mathbf{x}, \gamma)) = \frac{1}{2}\phi_\rho(\underline{M}(f', \mathbf{x}, \gamma)) + \frac{1}{2} < 1.$$

Therefore for any $\mathbf{x} \in \mathcal{X}_2$,

$$\inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \frac{1}{2}) \leq \mathcal{C}_{\tilde{\phi}_\rho}(f', \mathbf{x}, \frac{1}{2}) < 1 \leq \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0 \leq \overline{M}(f, \mathbf{x}, \gamma)} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \frac{1}{2}). \quad (25)$$

- When $\eta \in (\frac{1}{2}, 1]$, since ϕ_ρ is non-increasing, for any $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{aligned} \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) &= \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \eta\phi_\rho(\underline{M}(f, \mathbf{x}, \gamma)) + (1 - \eta)\phi_\rho(-\overline{M}(f, \mathbf{x}, \gamma)) \\ &= \eta + \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} (1 - \eta)\phi_\rho(-\overline{M}(f, \mathbf{x}, \gamma)) \\ &\geq \eta + (1 - \eta)\phi_\rho(M_{\mathbf{x}}). \end{aligned}$$

On the other hand, for any $\mathbf{x} \in \mathcal{X}_2$ and $\epsilon > 0$,

$$\begin{aligned} \mathcal{C}_{\tilde{\phi}_\rho}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) &= \eta\phi_\rho(\underline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) + (1 - \eta)\phi_\rho(-\overline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) \\ &< \eta\phi_\rho(M_{\mathbf{x}}) + \epsilon + (1 - \eta). \end{aligned}$$

Since $\eta > \frac{1}{2}$ and $M_{\mathbf{x}} > 0$, we have

$$\begin{aligned} &\inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) - \mathcal{C}_{\tilde{\phi}_\rho}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) \\ &> [\eta + (1 - \eta)\phi_\rho(M_{\mathbf{x}})] - [\eta\phi_\rho(M_{\mathbf{x}}) + \epsilon + (1 - \eta)] \\ &= (2\eta - 1)(1 - \phi_\rho(M_{\mathbf{x}})) - \epsilon \\ &> 0, \end{aligned}$$

where we take $0 < \epsilon < (2\eta - 1)(1 - \phi_\rho(M_{\mathbf{x}}))$.

Therefore for any $\eta \in (\frac{1}{2}, 1]$ and $\mathbf{x} \in \mathcal{X}_2$, there exists $0 < \epsilon < (2\eta - 1)(1 - \phi_\rho(M_{\mathbf{x}}))$ such that

$$\inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) \leq \mathcal{C}_{\tilde{\phi}_\rho}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) < \inf_{f \in \mathcal{H}: \underline{M}(f, \mathbf{x}, \gamma) \leq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta). \quad (26)$$

- When $\eta \in [0, \frac{1}{2})$, since ϕ_ρ is non-increasing, for any $\mathbf{x} \in \mathcal{X}_2$,

$$\begin{aligned} \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) &= \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \eta\phi_\rho(\underline{M}(f, \mathbf{x}, \gamma)) + (1 - \eta)\phi_\rho(-\overline{M}(f, \mathbf{x}, \gamma)) \\ &= 1 - \eta + \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \eta\phi_\rho(\underline{M}(f, \mathbf{x}, \gamma)) \\ &\geq 1 - \eta + \eta\phi_\rho(M_{\mathbf{x}}) \end{aligned}$$

On the other hand, for any $\mathbf{x} \in \mathcal{X}_2$ and $\epsilon > 0$,

$$\begin{aligned} \mathcal{C}_{\tilde{\phi}_\rho}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) &= \eta\phi_\rho(\underline{M}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) + (1 - \eta)\phi_\rho(-\overline{M}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) \\ &= \eta + (1 - \eta)\phi_\rho(\underline{M}(f_{\mathbf{x}}^\epsilon, \mathbf{x}, \gamma)) \\ &< \eta + (1 - \eta)\phi_\rho(M_{\mathbf{x}}) + \epsilon \end{aligned}$$

Since $\eta < \frac{1}{2}$ and $M_{\mathbf{x}} > 0$, we have

$$\begin{aligned} & \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) - \mathcal{C}_{\tilde{\phi}_\rho}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) \\ & > [1 - \eta + \eta\phi_\rho(M_{\mathbf{x}})] - [\eta + (1 - \eta)\phi_\rho(M_{\mathbf{x}}) + \epsilon] \\ & = (1 - 2\eta)(1 - \phi_\rho(M_{\mathbf{x}})) - \epsilon \\ & > 0 \end{aligned}$$

where we take $0 < \epsilon < (1 - 2\eta)(1 - \phi_\rho(M_{\mathbf{x}}))$.

Therefore for any $\eta \in [0, \frac{1}{2})$ and $\mathbf{x} \in \mathcal{X}_2$, there exists $0 < \epsilon < (1 - 2\eta)(1 - \phi_\rho(M_{\mathbf{x}}))$ such that

$$\inf_{f \in \mathcal{H}} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta) \leq \mathcal{C}_{\tilde{\phi}_\rho}(-f_{\mathbf{x}}^\epsilon, \mathbf{x}, \eta) < \inf_{f \in \mathcal{H}: \overline{M}(f, \mathbf{x}, \gamma) \geq 0} \mathcal{C}_{\tilde{\phi}_\rho}(f, \mathbf{x}, \eta). \quad (27)$$

To sum up, by (25), (26) and (27), we conclude that $\tilde{\phi}_\rho$ is \mathcal{H} -calibrated with respect to ℓ_γ . \blacksquare

A.4. Proof of Theorem 13

As shown by Awasthi et al. (2020), for $f \in \mathcal{H}_g$, the adversarial 0/1 loss has the equivalent form

$$\ell_\gamma(f, \mathbf{x}, y) = \mathbb{1}_{\inf_{\mathbf{x}': \|\mathbf{x} - \mathbf{x}'\| \leq \gamma} (yg(\mathbf{w} \cdot \mathbf{x}') + by) \leq 0} = \mathbb{1}_{yg(\mathbf{w} \cdot \mathbf{x} - \gamma y \|\mathbf{w}\|) + by \leq 0} = \mathbb{1}_{yg(\mathbf{w} \cdot \mathbf{x} - \gamma y) + by \leq 0}. \quad (28)$$

The proofs of \mathcal{H} -consistency 13 will closely follow the proofs of Theorem 12 and Theorem 16. We will first prove Lemma 26 and Lemma 27 analogous to Lemma 24 and Lemma 25 respectively. Without loss of generality, assume that g is continuous and satisfies $g(-1 - \gamma) + G > 0$, $g(1 + \gamma) - G < 0$. Then observe that $g(-\gamma) + G > 0$, $g(\gamma) - G < 0$ and $0 \in \mathcal{H}_g$ since g is non-decreasing.

Lemma 26 *For a surrogate loss ℓ and hypothesis set \mathcal{H}_g , the calibration function of losses (ℓ, ℓ_γ) is*

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \begin{cases} +\infty & \text{if } \epsilon > \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) & \text{if } |2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \text{ or } (2\eta - 1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) & \text{if } \epsilon \leq |2\eta - 1|. \end{cases}$$

Proof Similarly to the proof of Lemma 24, we first characterize the inner ℓ -risk and minimal inner ℓ_γ -risk for \mathcal{H}_g . By the definition of inner risk (4) and equivalent form of adversarial 0-1 loss ℓ_γ for \mathcal{H}_g (28), the inner ℓ_γ -risk is

$$\begin{aligned} \mathcal{C}_{\ell_\gamma}(f, \mathbf{x}, \eta) &= \eta \mathbb{1}_{g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} + (1 - \eta) \mathbb{1}_{g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \\ &= \begin{cases} 1 & \text{if } g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b, \\ \eta & \text{if } g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b < 0, \\ 1 - \eta & \text{if } g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b > 0. \end{cases} \end{aligned}$$

where we used the fact that g is non-decreasing and $g(\mathbf{w} \cdot \mathbf{x} - \gamma) \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma)$. Note for any $\mathbf{x} \in \mathcal{X}$, $\mathbf{w} \cdot \mathbf{x} \in [-\|\mathbf{x}\|, \|\mathbf{x}\|]$. Thus we have $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \in [g(-\|\mathbf{x}\| - \gamma) - G, g(\|\mathbf{x}\| - \gamma) + G]$ and

$g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \in [g(-\|\mathbf{x}\| + \gamma) - G, g(\|\mathbf{x}\| + \gamma) + G]$ since g is non-decreasing. By the fact that $g(-\gamma) + G > 0$ and $g(\gamma) - G < 0$, we obtain the minimal inner ℓ_γ -risk, which is for any $\mathbf{x} \in \mathcal{X}$,

$$\mathcal{C}_{\ell_\gamma, \mathcal{H}_{\text{lin}}}^*(\mathbf{x}, \eta) = \min\{\eta, 1 - \eta\}.$$

Similarly to the derivation of $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_{\text{lin}}}(f, \mathbf{x}, \eta)$ (17), we derive $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta)$ as follows. By the observation (12), for any $\mathbf{x} \in \mathcal{X}$, for $f \in \mathcal{H}_g$ such that $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) = 1 - \min\{\eta, 1 - \eta\} = \max\{\eta, 1 - \eta\}$; for $f \in \mathcal{H}_g$ such that $g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b < 0$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) = \eta - \min\{\eta, 1 - \eta\} = \max\{0, 2\eta - 1\} = |2\eta - 1|\mathbb{1}_{(2\eta-1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0}$ since $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b < 0$; for $f \in \mathcal{H}_g$ such that $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b > 0$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) = 1 - \eta - \min\{\eta, 1 - \eta\} = \max\{0, 1 - 2\eta\} = |2\eta - 1|\mathbb{1}_{(2\eta-1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0}$ since $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b > 0$. Therefore,

$$\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) = \begin{cases} \max\{\eta, 1 - \eta\} & \text{if } g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b, \\ |2\eta - 1|\mathbb{1}_{(2\eta-1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} & \text{if } g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b < 0 \text{ or } g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b > 0. \end{cases}$$

By (7), for a fixed $\eta \in [0, 1]$ and $\mathbf{x} \in \mathcal{X}$, the calibration function of losses (ℓ, ℓ_γ) given \mathcal{H}_g is

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}_g} \{ \Delta\mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) \mid \Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) \geq \epsilon \}.$$

Similarly to the proof of Lemma 24, we then make use of the observation (12) for deriving the the calibration function. By the observation (12), if $\epsilon > \max\{\eta, 1 - \eta\}$, then for all $f \in \mathcal{H}_g$, $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) < \epsilon$, which implies that $\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \infty$; ff $|2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}$, then $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) \geq \epsilon$ if and only if $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b$, which leads to

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \Delta\mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta);$$

if $\epsilon \leq |2\eta - 1|$, then $\Delta\mathcal{C}_{\ell_\gamma, \mathcal{H}_g}(f, \mathbf{x}, \eta) \geq \epsilon$ if and only if $g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b$ or $(2\eta - 1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0$, which leads to

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \text{ or } (2\eta-1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} \Delta\mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta).$$

■

Lemma 27 *Let ℓ be a surrogate loss function. Then ℓ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if for any $\mathbf{x} \in \mathcal{X}$,*

$$\begin{aligned} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}), \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned}$$

Proof As the proof of Lemma 25 first make use of Lemma 24 and Proposition 5, we also first make use of Lemma 26 and Proposition 5 in the following proof. Let δ_{\max} be the calibration function of (ℓ, ℓ_γ) for hypothesis set \mathcal{H}_g . By Lemma 26,

$$\delta_{\max}(\epsilon, \mathbf{x}, \eta) = \begin{cases} +\infty & \text{if } \epsilon > \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) & \text{if } |2\eta - 1| < \epsilon \leq \max\{\eta, 1 - \eta\}, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \text{ or } (2\eta - 1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) & \text{if } \epsilon \leq |2\eta - 1|. \end{cases}$$

By Proposition 5, ℓ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if its calibration function δ_{\max} satisfies $\delta_{\max}(\epsilon, \mathbf{x}, \eta) > 0$ for all $\mathbf{x} \in \mathcal{X}$, $\eta \in [0, 1]$ and $\epsilon > 0$. The following steps are similar to the steps in the proof of Lemma 25, where we analyze by considering three cases.

For $\eta = \frac{1}{2}$, we have for any $\mathbf{x} \in \mathcal{X}$,

$$\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0 \text{ for all } \epsilon > 0 \Leftrightarrow \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \frac{1}{2}). \quad (29)$$

For $1 \geq \eta > \frac{1}{2}$, we have $|2\eta - 1| = 2\eta - 1$, $\max\{\eta, 1 - \eta\} = \eta$, and

$$\inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \text{ or } (2\eta - 1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta).$$

Therefore, $\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0$ for any $\mathbf{x} \in \mathcal{X}$, $\epsilon > 0$ and $\eta \in (\frac{1}{2}, 1]$ if and only if for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{cases} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } 2\eta - 1 < \epsilon \leq \eta, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \epsilon \leq 2\eta - 1, \end{cases}$$

for all $\epsilon > 0$, which is equivalent to for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{cases} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \epsilon \leq \eta < \frac{\epsilon+1}{2}, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) & \text{for all } \eta \in (\frac{1}{2}, 1] \text{ such that } \frac{\epsilon+1}{2} \leq \eta, \end{cases} \quad (30)$$

for all $\epsilon > 0$. Observe that

$$\begin{aligned} \left\{ \eta \in (\frac{1}{2}, 1] \mid \epsilon \leq \eta < \frac{\epsilon+1}{2}, \epsilon > 0 \right\} &= \left\{ \frac{1}{2} < \eta \leq 1 \right\}, \text{ and} \\ \left\{ \eta \in (\frac{1}{2}, 1] \mid \frac{\epsilon+1}{2} \leq \eta, \epsilon > 0 \right\} &= \left\{ \frac{1}{2} < \eta \leq 1 \right\}, \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_\ell(f, \mathbf{x}, \eta) &\geq \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta. \end{aligned}$$

Therefore, we reduce the above condition (30) as for any $\mathbf{x} \in \mathcal{X}$,

$$\inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_\ell(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_\ell(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1]. \quad (31)$$

For $\frac{1}{2} > \eta \geq 0$, we have $|2\eta - 1| = 1 - 2\eta$, $\max\{\eta, 1 - \eta\} = 1 - \eta$, and

$$\inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \text{ or } (2\eta - 1)[g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b] \leq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta) = \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \Delta \mathcal{C}_{\ell, \mathcal{H}_g}(f, \mathbf{x}, \eta).$$

Therefore, $\delta_{\max}(\epsilon, \mathbf{x}, \frac{1}{2}) > 0$ for any $\mathbf{x} \in \mathcal{X}$, $\epsilon > 0$ and $\eta \in [0, \frac{1}{2})$ if and only if for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{cases} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } 1 - 2\eta < \epsilon \leq 1 - \eta, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \epsilon \leq 1 - 2\eta, \end{cases}$$

for all $\epsilon > 0$, which is equivalent to for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{cases} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \frac{1-\epsilon}{2} < \eta \leq 1 - \epsilon, \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) & \text{for all } \eta \in [0, \frac{1}{2}) \text{ such that } \eta \leq \frac{1-\epsilon}{2}, \end{cases} \quad (32)$$

for all $\epsilon > 0$. Observe that

$$\begin{aligned} \left\{ \eta \in [0, \frac{1}{2}) \left| \frac{1-\epsilon}{2} < \eta \leq 1 - \epsilon, \epsilon > 0 \right. \right\} &= \left\{ 0 \leq \eta < \frac{1}{2} \right\}, \text{ and} \\ \left\{ \eta \in [0, \frac{1}{2}) \left| \eta \leq \frac{1-\epsilon}{2}, \epsilon > 0 \right. \right\} &= \left\{ 0 \leq \eta < \frac{1}{2} \right\}, \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) &\geq \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) \text{ for all } \eta. \end{aligned}$$

Therefore we reduce the above condition (32) as for any $\mathbf{x} \in \mathcal{X}$,

$$\inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) > \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\ell}(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \quad (33)$$

To sum up, by (29), (31) and (33), we conclude the proof. \blacksquare

Theorem 13 Let g be a non-decreasing and continuous function such that $g(1 + \gamma) < G$ and $g(-1 - \gamma) > -G$ for some $G \geq 0$. Let a margin-based loss ϕ be bounded, continuous, non-increasing, and quasi-concave even. Assume that $\phi(g(-t) - G) > \phi(G - g(-t))$ and $g(-t) + g(t) \geq 0$ for any $0 \leq t \leq 1$. Then ϕ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if for any $0 \leq t \leq 1$,

$$\begin{aligned} \phi(G - g(-t)) + \phi(g(-t) - G) &= \phi(g(t) + G) + \phi(-g(t) - G) \\ \text{and } \min\{\phi(\overline{A}(t)) + \phi(-\overline{A}(t)), \phi(\underline{A}(t)) + \phi(-\underline{A}(t))\} &> \phi(G - g(-t)) + \phi(g(-t) - G), \end{aligned}$$

where $\overline{A}(t) = \max_{s \in [-t, t]} g(s) - g(s - \gamma)$ and $\underline{A}(t) = \min_{s \in [-t, t]} g(s) - g(s + \gamma)$.

Proof By Lemma 27, ϕ is \mathcal{H}_g -calibrated with respect to ℓ_γ if and only if for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned} \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b} \mathcal{C}_{\phi}(f, \mathbf{x}, \frac{1}{2}) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\phi}(f, \mathbf{x}, \frac{1}{2}), \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0} \mathcal{C}_{\phi}(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\phi}(f, \mathbf{x}, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \inf_{f \in \mathcal{H}_g: g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0} \mathcal{C}_{\phi}(f, \mathbf{x}, \eta) &> \inf_{f \in \mathcal{H}_g} \mathcal{C}_{\phi}(f, \mathbf{x}, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned} \quad (34)$$

By the definition of inner risk (4), the inner ϕ -risk is

$$\mathcal{C}_\phi(f, \mathbf{x}, \eta) = \eta\phi(f(\mathbf{x})) + (1 - \eta)\phi(-f(\mathbf{x})).$$

and $f(\mathbf{x}) = g(\mathbf{w} \cdot \mathbf{x}) + b \in [g(-\|\mathbf{x}\|) - G, g(\|\mathbf{x}\|) + G]$ when $f \in \mathcal{H}_g$ since g is continuous and non-decreasing. Specifically, by the assumption that $g(-1 - \gamma) + G > 0$, $g(1 + \gamma) - G < 0$, when $f \in \{f \in \mathcal{H}_g : g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0 \leq g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b\}$, $f(\mathbf{x}) = g(\mathbf{w} \cdot \mathbf{x}) + b \in [\min_{-\|\mathbf{x}\| \leq s \leq \|\mathbf{x}\|} g(s) - g(s + \gamma), \max_{-\|\mathbf{x}\| \leq s \leq \|\mathbf{x}\|} g(s) - g(s - \gamma)]$; when $f \in \{f \in \mathcal{H}_g : g(\mathbf{w} \cdot \mathbf{x} - \gamma) + b \leq 0\}$, $f(\mathbf{x}) = g(\mathbf{w} \cdot \mathbf{x}) + b \in [g(-\|\mathbf{x}\|) - G, \max_{-\|\mathbf{x}\| \leq s \leq \|\mathbf{x}\|} g(s) - g(s - \gamma)]$; when $f \in \{f \in \mathcal{H}_g : g(\mathbf{w} \cdot \mathbf{x} + \gamma) + b \geq 0\}$, $f(\mathbf{x}) = g(\mathbf{w} \cdot \mathbf{x}) + b \in [\min_{-\|\mathbf{x}\| \leq s \leq \|\mathbf{x}\|} g(s) - g(s + \gamma), g(\|\mathbf{x}\|) + G]$. For convenience, we denote $\bar{A}(t) = \max_{-t \leq s \leq t} g(s) - g(s - \gamma) \geq 0$ and $\underline{A}(t) = \min_{-t \leq s \leq t} g(s) - g(s + \gamma) \leq 0$ for any $0 \leq t \leq 1$. Therefore, (34) is equivalent to for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned} \underline{A}(\|\mathbf{x}\|) \leq t \leq \bar{A}(\|\mathbf{x}\|) \quad \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) &> \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}), \text{ and} \\ \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq \bar{A}(\|\mathbf{x}\|)} \bar{\mathcal{C}}_\phi(t, \eta) &> \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) \text{ for all } \eta \in (\frac{1}{2}, 1], \text{ and} \\ \underline{A}(\|\mathbf{x}\|) \leq t \leq g(\|\mathbf{x}\|) + G \quad \bar{\mathcal{C}}_\phi(t, \eta) &> \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) \text{ for all } \eta \in [0, \frac{1}{2}). \end{aligned} \quad (35)$$

Suppose that ϕ is \mathcal{H}_g -calibrated with respect to ℓ_γ . Since for $\eta \in [0, \frac{1}{2})$,

$$\begin{aligned} \inf_{\underline{A}(\|\mathbf{x}\|) \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\bar{\mathcal{C}}_\phi(\underline{A}(\|\mathbf{x}\|), \eta), \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\ \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) &= \min\{\mathcal{C}_\phi(g(-\|\mathbf{x}\|) - G, \eta), \mathcal{C}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \quad (\text{Part 3 of Lemma 23}) \end{aligned}$$

we have $\mathcal{C}_\phi(g(-\|\mathbf{x}\|) - G, \eta) < \mathcal{C}_\phi(g(\|\mathbf{x}\|) + G, \eta)$ for any $\mathbf{x} \in \mathcal{X}$, otherwise

$$\inf_{\underline{A}(\|\mathbf{x}\|) \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) \leq \mathcal{C}_\phi(g(\|\mathbf{x}\|) + G, \eta) = \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta).$$

By Part 9 of Lemma 23, $\phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G)$ for all $0 \leq t \leq 1$. Also, for any $0 \leq t \leq 1$,

$$\begin{aligned} &\frac{1}{2} \min\{\phi(\bar{A}(t)) + \phi(-\bar{A}(t)), \phi(\underline{A}(t)) + \phi(-\underline{A}(t))\} \\ &= \inf_{\underline{A}(t) \leq t \leq \bar{A}(t)} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) \quad (\text{Part 3 of Lemma 23}) \\ &> \inf_{g(-t) - G \leq t \leq g(t) + G} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) \quad (35) \\ &= \frac{1}{2} \min\{\phi(G - g(-t)) + \phi(g(-t) - G), \phi(g(t) + G) + \phi(-g(t) - G)\} \quad (\text{Part 3 of Lemma 23}) \\ &= \frac{1}{2} (\phi(G - g(-t)) + \phi(g(-t) - G)) \end{aligned}$$

Now for the other direction, assume that for any $0 \leq t \leq 1$,

$$\begin{aligned} &\phi(G - g(-t)) + \phi(g(-t) - G) = \phi(g(t) + G) + \phi(-g(t) - G) \\ \text{and } \min\{\phi(\bar{A}(t)) + \phi(-\bar{A}(t)), \phi(\underline{A}(t)) + \phi(-\underline{A}(t))\} &> \phi(G - g(-t)) + \phi(g(-t) - G). \end{aligned}$$

Then for $\eta = \frac{1}{2}$ and any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned}
 & \inf_{\underline{A}(\|\mathbf{x}\|) \leq t \leq \bar{A}(\|\mathbf{x}\|)} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}) \\
 &= \frac{1}{2} \min\{\phi(\bar{A}(\|\mathbf{x}\|)) + \phi(-\bar{A}(\|\mathbf{x}\|)), \phi(\underline{A}(\|\mathbf{x}\|)) + \phi(-\underline{A}(\|\mathbf{x}\|))\} \quad (\text{Part 3 of Lemma 23}) \\
 &> \frac{1}{2}(\phi(G - g(-\|\mathbf{x}\|)) + \phi(g(-\|\mathbf{x}\|) - G)) \quad (\text{by Assumption}) \\
 &= \frac{1}{2} \min\{\phi(G - g(-\|\mathbf{x}\|)) + \phi(g(-\|\mathbf{x}\|) - G), \phi(g(\|\mathbf{x}\|) + G) + \phi(-g(\|\mathbf{x}\|) - G)\} \quad (\text{by Assumption}) \\
 &= \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \frac{1}{2}). \quad (\text{Part 3 of Lemma 23})
 \end{aligned}$$

For $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned}
 & \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq \bar{A}(\|\mathbf{x}\|)} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta), \bar{\mathcal{C}}_\phi(\bar{A}(\|\mathbf{x}\|), \eta)\} \quad (\text{Part 3 of Lemma 23}) \\
 & \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta), \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \quad (\text{Part 3 of Lemma 23}) \\
 &= \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta) \quad (\text{Part 8 of Lemma 23})
 \end{aligned}$$

Since ϕ is non-increasing, we have for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned}
 & \phi(-g(\|\mathbf{x}\|) - G) - \phi(g(\|\mathbf{x}\|) + G) + \phi(\bar{A}(\|\mathbf{x}\|)) - \phi(-\bar{A}(\|\mathbf{x}\|)) \\
 & \geq \phi(-g(\|\mathbf{x}\|) - G) - \phi(g(\|\mathbf{x}\|) + G) + \phi(g(\|\mathbf{x}\|) + G) - \phi(-g(\|\mathbf{x}\|) - G) \\
 & = 0.
 \end{aligned}$$

Then for $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned}
 & \bar{\mathcal{C}}_\phi(\bar{A}(\|\mathbf{x}\|), \eta) - \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta) \\
 &= (\phi(\bar{A}(\|\mathbf{x}\|)) - \phi(-\bar{A}(\|\mathbf{x}\|)) + \phi(-g(\|\mathbf{x}\|) - G) - \phi(g(\|\mathbf{x}\|) + G))\eta + \phi(-\bar{A}(\|\mathbf{x}\|)) - \phi(-g(\|\mathbf{x}\|) - G) \\
 &\geq (\phi(\bar{A}(\|\mathbf{x}\|)) - \phi(-\bar{A}(\|\mathbf{x}\|)) + \phi(-g(\|\mathbf{x}\|) - G) - \phi(g(\|\mathbf{x}\|) + G))\frac{1}{2} + \phi(-\bar{A}(\|\mathbf{x}\|)) - \phi(-g(\|\mathbf{x}\|) - G) \\
 &= \frac{1}{2}(\phi(\bar{A}(\|\mathbf{x}\|)) - \phi(-\bar{A}(\|\mathbf{x}\|)) - \phi(-g(\|\mathbf{x}\|) - G) - \phi(g(\|\mathbf{x}\|) + G)) \\
 &> 0.
 \end{aligned}$$

In addition, by Part 8 of Lemma 23, for all $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$, $\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) - \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta) > 0$. As a result, for $\eta \in (\frac{1}{2}, 1]$ and any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned}
 & \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq \bar{A}(\|\mathbf{x}\|)} \bar{\mathcal{C}}_\phi(t, \eta) - \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) \\
 &= \min\{\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) - \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta), \bar{\mathcal{C}}_\phi(\bar{A}(\|\mathbf{x}\|), \eta) - \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \\
 &> 0.
 \end{aligned}$$

Finally, for $\eta \in [0, \frac{1}{2})$, by Part 9 of Lemma 23, we have $\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) < \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)$ and

$$\inf_{\underline{A}(\|\mathbf{x}\|) \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(\underline{A}(\|\mathbf{x}\|), \eta), \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \quad (\text{Part 3 of Lemma 23})$$

$$\inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) = \min\{\bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta), \bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta)\} \quad (\text{Part 3 of Lemma 23})$$

$$= \bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) \quad (\text{Part 9 of Lemma 23})$$

Since $\phi(\underline{A}(\|\mathbf{x}\|)) + \phi(-\underline{A}(\|\mathbf{x}\|)) > \phi(G - g(-\|\mathbf{x}\|)) + \phi(g(-\|\mathbf{x}\|) - G)$ and ϕ is non-increasing, we have for any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned} & \phi(G - g(-\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) + \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(-\underline{A}(\|\mathbf{x}\|)) \\ &= \phi(G - g(-\|\mathbf{x}\|)) - \phi(-\underline{A}(\|\mathbf{x}\|)) + \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) \\ &< \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) + \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) \\ &= 2[\phi(\underline{A}(\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G)] \\ &\leq 0. \end{aligned}$$

Then for $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$.

$$\begin{aligned} & \bar{\mathcal{C}}_\phi(\underline{A}(\|\mathbf{x}\|), \eta) - \bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) \\ &= [\phi(G - g(-\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) + \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(-\underline{A}(\|\mathbf{x}\|))] \eta + \phi(-\underline{A}(\|\mathbf{x}\|)) - \phi(G - g(-\|\mathbf{x}\|)) \\ &\geq [\phi(G - g(-\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) + \phi(\underline{A}(\|\mathbf{x}\|)) - \phi(-\underline{A}(\|\mathbf{x}\|))] \frac{1}{2} + \phi(-\underline{A}(\|\mathbf{x}\|)) - \phi(G - g(-\|\mathbf{x}\|)) \\ &= \frac{1}{2} [\phi(\underline{A}(\|\mathbf{x}\|)) + \phi(-\underline{A}(\|\mathbf{x}\|)) - \phi(g(-\|\mathbf{x}\|) - G) - \phi(G - g(-\|\mathbf{x}\|))] \\ &> 0. \end{aligned}$$

In addition, by Part 9 of Lemma 23, for all $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$, $\bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta) - \bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta) > 0$. As a result, for $\eta \in [0, \frac{1}{2})$ and any $\mathbf{x} \in \mathcal{X}$,

$$\begin{aligned} & \inf_{\underline{A}(\|\mathbf{x}\|) \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) - \inf_{g(-\|\mathbf{x}\|) - G \leq t \leq g(\|\mathbf{x}\|) + G} \bar{\mathcal{C}}_\phi(t, \eta) \\ &= \min\{\bar{\mathcal{C}}_\phi(g(\|\mathbf{x}\|) + G, \eta) - \bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta), \bar{\mathcal{C}}_\phi(\underline{A}(\|\mathbf{x}\|), \eta) - \bar{\mathcal{C}}_\phi(g(-\|\mathbf{x}\|) - G, \eta)\} \\ &> 0. \end{aligned}$$

■