
A All 12 hypotheses from runs 0001-0004
(bar colour = source run)
1.H1 GI-lineage TFs (HNF1A/HNF4A/FOXA2) define the Metabo
1.H2 Natural-EBV proteome module; TMK1 a cryptic EB
[carrier rejected]
1.H3 In-vitro EBV infection (eAGS) lowers MHC-I antigen p
2.H1 Metabolism subtype runs a glutathione/antioxidant de
2.H2 MYC-target load is coupled to the proteostasis/UPR m
2.H3 No Warburg glycolysis<->OXPH
[weak]
3.H1 Metabolism subtype = convergent RTK (MET/FGFR/RON/Ep
3.H2 EMT subtype = PI3K-AKT-mTOR with a co-active SGK arm
3.H3 The robust subtype signalling difference is tyrosine
4.H1 Candidate novel ER/secretory-r
[candidate]
4.H2 Subtype-coordinated complex abundance: OXPHOS comple
4.H3 Co-abundance here is a reliable known-complex map bu
run 1
run 2
run 3
run 4
0
1
2
3
4
5
6
evidence strength (author rating 1-5)
B Prioritised integrated hypotheses
(convergence = same data, not independent validation)
P1: The 'Metabolism' subtype is a well-differentia
P2: Metabolism subtype depends on RTK->RAS-
P3: The Metabolism subtype carries a glutathione/a
P4: A MYC-target / proteostasis (UPR) proliferatio
P5: In-vitro EBV infection imposes a cell-intrinsi
0.0
0.5
1.0
1.5
2.0
2.5
3.0
3.5
4.0
supporting analyses (of 4 runs)