
A 26 most variable hallmark pathways across lines
MYC_TARGETS_V2
OXIDATIVE_PHOSPHORYLATION
TGF_BETA_SIGNALING
E2F_TARGETS
G2M_CHECKPOINT
FATTY_ACID_METABOLISM
INTERFERON_ALPHA_RESPONSE
COAGULATION
ADIPOGENESIS
CHOLESTEROL_HOMEOSTASIS
ESTROGEN_RESPONSE_LATE
BILE_ACID_METABOLISM
EPITHELIAL_MESENCHYMAL_TRANSITION
XENOBIOTIC_METABOLISM
UV_RESPONSE_DN
HYPOXIA
MYC_TARGETS_V1
INTERFERON_GAMMA_RESPONSE
PEROXISOME
MYOGENESIS
ESTROGEN_RESPONSE_EARLY
APICAL_JUNCTION
UNFOLDED_PROTEIN_RESPONSE
P53_PATHWAY
MITOTIC_SPINDLE
PI3K_AKT_MTOR_SIGNALING
AGS
eAGS
AZ521
SNU1750
MKN28
N87
TMK1
NCC24
SNU719
YCCEL1
KatoIII
Ocum1
SNU601
YCC3
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
pathway activity (mean protein z)


B Proliferation (mod 1) and metabolism (mod 4) vary independently
(module r = -0.12, n.s.); the metabolism module separates
EMT (low) from Metabolism (high) lines, Welch p = 0.008
module 4: OXIDATIVE_PHOSPHORYLATION / FATTY_ACID_METABOLISM / ADIPOGENESIS...
module 1: MYC_TARGETS_V2 / E2F_TARGETS / G2M_CHECKPOINT...
NCC24
YCCEL1
SNU719
TMK1
YCC3
KatoIII
Ocum1
N87
AGS
SNU601
MKN28
eAGS
SNU1750
AZ521
Hu et al.
EMT
Metabolism
0.4
0.2
0.0
−0.2
−0.4
−0.4
−0.2
0.0
0.2
0.4