


Introduction to Logic

Lecture 5: Sets, Relations, Functions

Davide Grossi

4 December 2023



Overview

Sets

Relations

Functions


Overview

Sets

Relations

Functions

The beginning of Set Theory



Georg Cantor

Beiträge zur Begründung der transfiniten Mengenlehre (1869)

Contributions to the Founding of the Theory of Transfinite Numbers (1915)

<https://archive.org/details/contributionstof00cant>

Set (classical definition)



By an “aggregate” (*Menge*) we are to understand any collection into a whole (*Zusammenfassung zu einem Ganzen*) M of definite and separate objects m of our intuition or our thought. These objects are called the “elements” of M .

Set theory (LPL Section 1.6 and Syllabus, Section 8.1)

Examples of **finite sets**:

- ▶ $\{1, 2, 3\}$
- ▶ $\{\textit{aristoteles}, \textit{plato}, \textit{socrates}\}$
- ▶ $\{\textit{mars}, \textit{merkel}, \textit{mounteverest}, 42\}$
- ▶ $\{1, \{2, 3\}, \{4, \{5, 6\}\}\}$

Set theory (LPL Section 1.6 and Syllabus, Section 8.1)

Examples of **finite sets**:

- ▶ $\{1, 2, 3\}$
- ▶ $\{\text{aristoteles}, \text{plato}, \text{socrates}\}$
- ▶ $\{\text{mars}, \text{merkel}, \text{mounteverest}, 42\}$
- ▶ $\{1, \{2, 3\}, \{4, \{5, 6\}\}\}$

Examples of **infinite sets**:

- ▶ \mathbb{N} , the set of natural numbers $0, 1, 2, \dots$
- ▶ \mathbb{Z} , the set of integers $\dots, -2, -1, 0, 1, 2, \dots$

Set theory (LPL Section 1.6 and Syllabus, Section 8.1)

Examples of **finite sets**:

- ▶ $\{1, 2, 3\}$
- ▶ $\{\text{aristoteles}, \text{plato}, \text{socrates}\}$
- ▶ $\{\text{mars}, \text{merkel}, \text{mounteverest}, 42\}$
- ▶ $\{1, \{2, 3\}, \{4, \{5, 6\}\}\}$


Examples of **infinite sets**:

- ▶ \mathbb{N} , the set of natural numbers $0, 1, 2, \dots$
- ▶ \mathbb{Z} , the set of integers $\dots, -2, -1, 0, 1, 2, \dots$

Examples which are **not sets**:


- ▶ $\langle 1, 2, 3 \rangle$
- ▶ $\langle 2, 1, 3 \rangle$
- ▶ $P \wedge Q$
- ▶ $\text{Large}()$

Membership in a set



- ▶ Notation: $a \in b$.
- ▶ “ a is in b ” or “ a is an element of b ” or “ a is a member of b ”
- ▶ For “ a is not in b ” we write $a \notin b$ (or $\neg(a \in b)$).

Membership in a set



- ▶ Notation: $a \in b$.
- ▶ “ a is in b ” or “ a is an element of b ” or “ a is a member of b ”
- ▶ For “ a is not in b ” we write $a \notin b$ (or $\neg(a \in b)$).



Giuseppe Peano

Arithmetices Principia Nova Methodo Exposita (1889)

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : $\{2, 4, 6\}$

c : $\{2, \{2, 4, 6\}\}$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$ FALSE

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$ FALSE

$a \in c$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$ FALSE

$a \in c$ TRUE

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$ FALSE

$a \in c$ TRUE

$b \in c$

Membership in a set

When is the sentence $a \in b$ true?

A sentence of the form $a \in b$ is true if and only if b is a set, and a is an element of that set.

Example

Given:

a : 2

b : {2, 4, 6}

c : {2, {2, 4, 6}}

Then we have:

$a \in a$ FALSE Note: 2 is not the same as {2}.

$a \in b$ TRUE

$b \in a$ FALSE

$b \in b$ FALSE

$a \in c$ TRUE

$b \in c$ TRUE

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} \quad \{2, 1\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\begin{array}{l} \{1, 2\} = \{2, 1\} \\ \{1, 2\} \neq \{1, 1, 2\} \end{array}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

$$\{1, 2\} = \{1, 1, 2\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

$$\{1, 2\} = \{1, 1, 2\}$$

$$\{1, 2\} \neq \{1, 2, 3\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

$$\{1, 2\} = \{1, 1, 2\}$$

$$\{1, 2\} \neq \{1, 2, 3\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

$$\{1, 2\} = \{1, 1, 2\}$$

$$\{1, 2\} \neq \{1, 2, 3\}$$

$$\{1, 2\} \neq \{\{1, 2\}\}$$

Relations between sets: Equality

Definition

$A = B$ iff A and B contain exactly the same elements.

That is, for all x it holds that: $x \in A$ if and only if $x \in B$.

Example

$$\{1, 2\} = \{2, 1\}$$

$$\{1, 2\} = \{1, 1, 2\}$$

$$\{1, 2\} \neq \{1, 2, 3\}$$

$$\{1, 2\} \neq \{\{1, 2\}\}$$

Relations between sets: Empty set & Inclusion

Definition

\emptyset The **empty set** \emptyset or \varnothing is the set that contains no elements. That is, *there is no x such that $x \in \emptyset$.*

Relations between sets: Empty set & Inclusion

Definition

- \emptyset The **empty set** \emptyset or \varnothing is the set that contains no elements.
That is, *there is no x such that $x \in \emptyset$.*
- \subseteq Set A is a **subset** of B , written $A \subseteq B$, iff all elements of A are also elements of B .
That is, *for all x it holds that (if $x \in A$ then $x \in B$).*

Relations between sets: Empty set & Inclusion

Definition

- \emptyset The **empty set** \emptyset or \varnothing is the set that contains no elements.
That is, *there is no x such that $x \in \emptyset$.*
- \subseteq Set A is a **subset** of B , written $A \subseteq B$, iff all elements of A are also elements of B .
That is, *for all x it holds that (if $x \in A$ then $x \in B$).*
- \subsetneq Set A is a **proper subset**, written $A \subsetneq B$, iff A is a subset of B , but B has at least one additional element.
That is, *$A \subseteq B$ and there is an x s.t. ($x \in B$ and $x \notin A$).*
Note: LPL and old exams use ' \subset ' for proper subsets.

Relations between sets: Empty set & Inclusion

Definition

- ∅ The **empty set** \emptyset or \varnothing is the set that contains no elements.
That is, *there is no x such that $x \in \emptyset$.*
- \subseteq Set A is a **subset** of B , written $A \subseteq B$, iff all elements of A are also elements of B .
That is, *for all x it holds that (if $x \in A$ then $x \in B$).*
- \subsetneq Set A is a **proper subset**, written $A \subsetneq B$, iff A is a subset of B , but B has at least one additional element.
That is, *$A \subseteq B$ and there is an x s.t. ($x \in B$ and $x \notin A$).*
Note: LPL and old exams use ' \subset ' for proper subsets.

We have for all sets A and B :

- ▶ $\emptyset \subseteq A$ and $A \subseteq A$

Relations between sets: Empty set & Inclusion

Definition

- ∅ The **empty set** \emptyset or \varnothing is the set that contains no elements.
That is, *there is no x such that $x \in \emptyset$.*
- \subseteq Set A is a **subset** of B , written $A \subseteq B$, iff all elements of A are also elements of B .
That is, *for all x it holds that (if $x \in A$ then $x \in B$).*
- \subsetneq Set A is a **proper subset**, written $A \subsetneq B$, iff A is a subset of B , but B has at least one additional element.
That is, *$A \subseteq B$ and there is an x s.t. ($x \in B$ and $x \notin A$).*
Note: LPL and old exams use ' \subset ' for proper subsets.

We have for all sets A and B :

- ▶ $\emptyset \subseteq A$ and $A \subseteq A$
- ▶ if $A \subseteq B$ and $B \subseteq A$ then $A = B$
- ▶ if $A \subseteq \emptyset$ then $A = \emptyset$

Relations between sets: Empty set & Inclusion

Definition

- ∅ The **empty set** \emptyset or \varnothing is the set that contains no elements.
That is, *there is no x such that $x \in \emptyset$.*
- \subseteq Set A is a **subset** of B , written $A \subseteq B$, iff all elements of A are also elements of B .
That is, *for all x it holds that (if $x \in A$ then $x \in B$).*
- \subsetneq Set A is a **proper subset**, written $A \subsetneq B$, iff A is a subset of B , but B has at least one additional element.
That is, *$A \subseteq B$ and there is an x s.t. ($x \in B$ and $x \notin A$).*
Note: LPL and old exams use ' \subset ' for proper subsets.

We have for all sets A and B :

- ▶ $\emptyset \subseteq A$ and $A \subseteq A$
- ▶ if $A \subseteq B$ and $B \subseteq A$ then $A = B$
- ▶ if $A \subseteq \emptyset$ then $A = \emptyset$
- ▶ $A \subsetneq A$ never holds
- ▶ $A \subsetneq \emptyset$ never holds

Operations on sets

Intersection $A \cap B$ is $\{x \mid x \in A \text{ and } x \in B\}$

Union $A \cup B$ is $\{x \mid x \in A \text{ or } x \in B\}$

Operations on sets

Intersection $A \cap B$ is $\{x \mid x \in A \text{ and } x \in B\}$

Union $A \cup B$ is $\{x \mid x \in A \text{ or } x \in B\}$

Difference $A \setminus B$ is $\{x \mid x \in A \text{ and } x \notin B\}$

Operations on sets

Intersection $A \cap B$ is $\{x \mid x \in A \text{ and } x \in B\}$

Union $A \cup B$ is $\{x \mid x \in A \text{ or } x \in B\}$

Difference $A \setminus B$ is $\{x \mid x \in A \text{ and } x \notin B\}$
(alternative notation: $A - B$)

Operations on sets

Intersection $A \cap B$ is $\{x \mid x \in A \text{ and } x \in B\}$

Union $A \cup B$ is $\{x \mid x \in A \text{ or } x \in B\}$

Difference $A \setminus B$ is $\{x \mid x \in A \text{ and } x \notin B\}$
(alternative notation: $A - B$)

We often work in a large set U called *universe* of discourse.
If U is given, then we can define:

Complement $A^c := U \setminus A = \{x \in U \mid x \notin A\}$

Overview of Symbols

You should know by now:

▶ \in

▶ \notin

▶ \subseteq

▶ \subsetneq

▶ \emptyset

▶ \cap

▶ \cup

▶ \setminus

Overview of Symbols

You should know by now:

- ▶ \in `\in`
- ▶ \notin `\notin`
- ▶ \subseteq `\subseteq`
- ▶ \subsetneq `\subsetneq`

- ▶ \emptyset `\emptyset`

- ▶ \cap `\cap`
- ▶ \cup `\cup`
- ▶ \setminus `\setminus`

Next to the symbols you can see their \LaTeX commands.

A very useful tool/website if you are looking for symbol commands: <https://detexify.kirelabs.org/>.

Building sets: Abstraction

Let E be any property, for example “being an even number”, “being a mathematical entity”, “being nice”, ...

General abstraction says:

*There exists a set A such that
for any x , $x \in A$ if and only if $E(x)$*

We write this new set as:

$$\{x \mid E(x)\}$$

That is, the set of all objects with property E .

Building sets: Abstraction

Let E be any property, for example “being an even number”, “being a mathematical entity”, “being nice”, ...

General abstraction says:

*There exists a set A such that
for any x , $x \in A$ if and only if $E(x)$*

We write this new set as:

$$\{x \mid E(x)\}$$

That is, the set of all objects with property E .

Abstraction can also be **relative** to a set, for example:

$$\{x \in \mathbb{N} \mid \text{Even}(x)\}$$

Building sets: Abstraction examples

Examples of **general** abstraction:

- ▶ $\{x \mid Loud(x) \wedge InGroningen(x)\} = \{martiniToren, forum, \dots\}$
- ▶ $\{x \mid x = bob \vee x = alice\} = \{alice, bob\}$

Building sets: Abstraction examples

Examples of **general** abstraction:

- ▶ $\{x \mid Loud(x) \wedge InGroningen(x)\} = \{martiniToren, forum, \dots\}$
- ▶ $\{x \mid x = bob \vee x = alice\} = \{alice, bob\}$

Examples of **relative** abstraction:

- ▶ $\{x \in \mathbb{N} \mid Odd(x)\} = \{1, 3, 5, 7, \dots\}$
- ▶ $\{x \in \{1, 23, 42\} \mid Even(x)\} = \{42\}$

Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

What about R itself? Does R belong to R ? Do we have $R \in R$?

Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

What about R itself? Does R belong to R ? Do we have $R \in R$?



Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

What about R itself? Does R belong to R ? Do we have $R \in R$?



$$R \in R \text{ iff } R \notin R$$

Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

What about R itself? Does R belong to R ? Do we have $R \in R$?



$R \in R$ iff $R \notin R$ Contradiction! Such a set R cannot exist!

Russell's paradox

Using general abstraction, let R be the set defined by the property “not being a member of itself”

$$R := \{x \mid x \notin x\}$$

Then we have for any x : $x \in R$ iff $x \notin x$

What about R itself? Does R belong to R ? Do we have $R \in R$?



$R \in R$ iff $R \notin R$ Contradiction! Such a set R cannot exist!

Moral: General abstraction $\{x \mid \dots\}$ is “dangerous”!

Only *relative* abstraction is safe: $\{x \in A \mid \dots\}$.


Bertrand Russell (1872–1970)

Either a thing is true or it isn't.

If it is true, you should believe it.

And if it isn't you shouldn't.

— Bertrand Russell



Logicomix: An Epic Search for Truth by Papadimitriou et al.

<https://www.logicomix.com/>

Overview

Sets

Relations

Functions

Ordered pairs

Ordered pair of a and b is written as $\langle a, b \rangle$.

Main property: $\langle a, b \rangle = \langle c, d \rangle$ if and only if $a = c$ and $b = d$

Ordered pairs

Ordered pair of a and b is written as $\langle a, b \rangle$.

Main property: $\langle a, b \rangle = \langle c, d \rangle$ if and only if $a = c$ and $b = d$

Order matters:

$$\langle 1, 2 \rangle \neq \langle 2, 1 \rangle$$

Recall that order does not matter for sets:

$$\{1, 2\} = \{2, 1\}$$

Relations

Definition

A (binary) relation is a set of ordered pairs.


Relations

Definition

A (binary) relation is a set of ordered pairs.

Example:

$\{\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle\}$



Relations

Definition


A (binary) relation is a set of ordered pairs.

The domain of R is the set $\{x \mid \langle x, y \rangle \in R, \text{ for some } y\}$.

The codomain (range) of R is the set $\{y \mid \langle x, y \rangle \in R, \text{ for some } x\}$.

Example:

$\{\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle\}$



Relations

Definition

A (binary) relation is a set of ordered pairs.

The domain of R is the set
 $\{x \mid \langle x, y \rangle \in R, \text{ for some } y\}$.

The codomain (range) of R is the set
 $\{y \mid \langle x, y \rangle \in R, \text{ for some } x\}$.


The Cartesian product of A and B is
 $A \times B := \{\langle x, y \rangle \mid x \in A \text{ and } y \in B\}$.
This is the largest relation with domain A
and codomain B .

A relation on A is a subset of $A \times A$.

Notation: $R \subseteq A \times A$.

Example:

$\{\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle\}$



Relations

Definition

A **(binary) relation** is a set of ordered pairs.

The **domain** of R is the set $\{x \mid \langle x, y \rangle \in R, \text{ for some } y\}$.


The **codomain (range)** of R is the set $\{y \mid \langle x, y \rangle \in R, \text{ for some } x\}$.

The **Cartesian product** of A and B is $A \times B := \{\langle x, y \rangle \mid x \in A \text{ and } y \in B\}$.
This is the largest relation with domain A and codomain B .

A **relation on A** is a subset of $A \times A$.
Notation: $R \subseteq A \times A$.

Example:

$\{\langle 1, 2 \rangle, \langle 1, 3 \rangle, \langle 2, 3 \rangle\}$



Some relations on \mathbb{N} are $=$, \leq , \geq , $<$ and $>$.

Properties of relations

Let R be a relation on some set A , i.e. $R \subseteq A \times A$.

We define the following properties:

reflexivity for all $x \in A$: $\langle x, x \rangle \in R$

symmetry for all $x, y \in A$: if $\langle x, y \rangle \in R$, then $\langle y, x \rangle \in R$.

transitivity for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle y, z \rangle \in R$, then $\langle x, z \rangle \in R$.

density for all $x, y \in A$: if $\langle x, y \rangle \in R$,
then there is a $z \in A$
such that $\langle x, z \rangle \in R$ and $\langle z, y \rangle \in R$.

Properties of relations

Let R be a relation on some set A , i.e. $R \subseteq A \times A$.

We define the following properties:

reflexivity for all $x \in A$: $\langle x, x \rangle \in R$

symmetry for all $x, y \in A$: if $\langle x, y \rangle \in R$, then $\langle y, x \rangle \in R$.

transitivity for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle y, z \rangle \in R$, then $\langle x, z \rangle \in R$.

density for all $x, y \in A$: if $\langle x, y \rangle \in R$,
then there is a $z \in A$
such that $\langle x, z \rangle \in R$ and $\langle z, y \rangle \in R$.

functionality for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.

Overview

Sets

Relations

Functions

Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



- If a relation R on A is functional, it is a **partial function** on A .

Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



- ▶ If a relation R on A is functional, it is a **partial function** on A .
- ▶ If a relation R on A is functional and its domain is A , then R is a **(total) function** on A .

Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



- ▶ If a relation R on A is functional, it is a **partial function** on A .
- ▶ If a relation R on A is functional and its domain is A , then R is a **(total) function** on A .
- ▶ To denote functions (functional relations), we often use f, g, h, \dots (rather than R).

Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



- ▶ If a relation R on A is functional, it is a **partial function** on A .
- ▶ If a relation R on A is functional and its domain is A , then R is a **(total) function** on A .
- ▶ To denote functions (functional relations), we often use f, g, h, \dots (rather than R).
- ▶ Instead of $\langle x, y \rangle \in f$, we often write **$f(x) = y$ or $y = f(x)$** .

Functions

functionality For all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle x, z \rangle \in R$, then $y = z$.



- ▶ If a relation R on A is functional, it is a **partial function** on A .
- ▶ If a relation R on A is functional and its domain is A , then R is a **(total) function** on A .
- ▶ To denote functions (functional relations), we often use f, g, h, \dots (rather than R).
- ▶ Instead of $\langle x, y \rangle \in f$, we often write **$f(x) = y$ or $y = f(x)$** .
- ▶ $f: A \rightarrow B$ means: f is a (total) function with domain A and range $\subseteq B$.

Properties of functions

Let $f : A \rightarrow B$ be a (total) function with domain A and range $\subseteq B$. Such a function f may have some additional properties:

- ▶ If $f : A \rightarrow B$ is such that the range of f is B , then f is called a **surjection**.

Properties of functions

Let $f : A \rightarrow B$ be a (total) function with domain A and range $\subseteq B$. Such a function f may have some additional properties:

- ▶ If $f : A \rightarrow B$ is such that the range of f is B , then f is called a **surjection**.
- ▶ If $f : A \rightarrow B$ is such that:
 for all $x, y, z \in A$:
 if $\langle x, y \rangle \in R$ and $\langle z, y \rangle \in R$, then $x = z$,
then f is called an **injection**.

Properties of functions

Let $f : A \rightarrow B$ be a (total) function with domain A and range $\subseteq B$. Such a function f may have some additional properties:

- ▶ If $f : A \rightarrow B$ is such that the range of f is B , then f is called a **surjection**.
- ▶ If $f : A \rightarrow B$ is such that:
for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle z, y \rangle \in R$, then $x = z$,
then f is called an **injection**.
- ▶ If f is both an injection and a surjection, then f is a **bijection**.
If there is such a bijection from A to B , then A and B have equal size.

Properties of functions

Let $f : A \rightarrow B$ be a (total) function with domain A and range $\subseteq B$. Such a function f may have some additional properties:

- ▶ If $f : A \rightarrow B$ is such that the range of f is B , then f is called a **surjection**.
- ▶ If $f : A \rightarrow B$ is such that:
for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle z, y \rangle \in R$, then $x = z$,
then f is called an **injection**.
- ▶ If f is both an injection and a surjection, then f is a **bijection**.
If there is such a bijection from A to B , then A and B have equal size.
- ▶ There are also binary functions such as $+$ and \times , as well as n -ary functions $f : (A_1 \times \cdots \times A_n) \rightarrow B$.

Properties of functions

Let $f : A \rightarrow B$ be a (total) function with domain A and range $\subseteq B$. Such a function f may have some additional properties:

- ▶ If $f : A \rightarrow B$ is such that the range of f is B , then f is called a **surjection**.
- ▶ If $f : A \rightarrow B$ is such that:
for all $x, y, z \in A$:
if $\langle x, y \rangle \in R$ and $\langle z, y \rangle \in R$, then $x = z$,
then f is called an **injection**.
- ▶ If f is both an injection and a surjection, then f is a **bijection**.
If there is such a bijection from A to B , then A and B have equal size.
- ▶ There are also binary functions such as $+$ and \times , as well as n -ary functions $f : (A_1 \times \cdots \times A_n) \rightarrow B$.

Note: This was not in past exams, but may be used now!

Functions in First-order logic

We can add function symbols to the language of first-order logic.

Examples:

- ▶ $\text{father}(x)$: the father of x
- ▶ $\text{mother}(x)$: the mother of x
- ▶ $x + y$: the sum of x and y

Note: function symbols, just like constants, start with a lower-case letter, except special functions in infix notation such as $+$.

Functions in First-order logic

We can add function symbols to the language of first-order logic.

Examples:

- ▶ $\text{father}(x)$: the father of x
- ▶ $\text{mother}(x)$: the mother of x
- ▶ $x + y$: the sum of x and y

Note: function symbols, just like constants, start with a lower-case letter, except special functions in infix notation such as $+$.

(Constants, for example 5 or mary, are 0-place function symbols.)

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Definition

Terms in FOL are defined inductively:

BASIS 1 constants $a, b, c, d, e, n_1, n_2, \dots$ are terms;

BASIS 2 variables t, u, v, w, x, y, z are terms;

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Definition

Terms in FOL are defined inductively:

BASIS 1 constants $a, b, c, d, e, n_1, n_2, \dots$ are terms;

BASIS 2 variables t, u, v, w, x, y, z are terms;

CONSTRUCTION If t_1, \dots, t_n are terms and f is an n -ary function symbol, then $f(t_1, \dots, t_n)$ is also a term;

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Definition

Terms in FOL are defined inductively:

BASIS 1 constants $a, b, c, d, e, n_1, n_2, \dots$ are terms;

BASIS 2 variables t, u, v, w, x, y, z are terms;

CONSTRUCTION If t_1, \dots, t_n are terms and f is an n -ary function symbol, then $f(t_1, \dots, t_n)$ is also a term;

CLOSURE Nothing else is a term that is not constructed in finitely many steps from rules BASIS 1, BASIS 2, CONSTRUCTION.

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Definition

Terms in FOL are defined inductively:

BASIS 1 constants $a, b, c, d, e, n_1, n_2, \dots$ are terms;

BASIS 2 variables t, u, v, w, x, y, z are terms;

CONSTRUCTION If t_1, \dots, t_n are terms and f is an n -ary function symbol, then $f(t_1, \dots, t_n)$ is also a term;

CLOSURE Nothing else is a term that is not constructed in finitely many steps from rules BASIS 1, BASIS 2, CONSTRUCTION.

Example: $a(\text{OlderThan}(b))$ is not a term.

Inductive definition of FOL terms (Syllabus Section 10.1)

Functions can be nested: $\text{father}(\text{mother}(\text{max}))$ is a term.

Definition

Terms in FOL are defined inductively:

BASIS 1 constants $a, b, c, d, e, n_1, n_2, \dots$ are terms;

BASIS 2 variables t, u, v, w, x, y, z are terms;

CONSTRUCTION If t_1, \dots, t_n are terms and f is an n -ary function symbol, then $f(t_1, \dots, t_n)$ is also a term;

CLOSURE Nothing else is a term that is not constructed in finitely many steps from rules BASIS 1, BASIS 2, CONSTRUCTION.

Example: $a(\text{OlderThan}(b))$ is not a term.

Complex terms are used just like names/constants, to refer to objects in the domain of discourse.

Function symbols give more expressivity to FOL

Note that function symbols give extra expressivity to FOL:

- ▶ The set of atomic formulas is extended to include expressions with function symbols.

Example: $\text{Older}(\text{father}(\text{max}), \text{max})$ is an atomic sentence.

Function symbols give more expressivity to FOL

Note that function symbols give extra expressivity to FOL:

- ▶ The set of atomic formulas is extended to include expressions with function symbols.

Example: $\text{Older}(\text{father}(\text{max}), \text{max})$ is an atomic sentence.

- ▶ Functions can be nested. Note: predicates cannot be nested.

Example

$\text{father}(\text{mother}(\text{max}))$ is a term.

$\text{Loves}(\text{father}(\text{mother}(\text{max})), \text{mary})$ is a sentence.

$\text{Older}(\text{Loves}(\text{juliet}, \text{romeo}))$ is not well-formed.

Function symbols give more expressivity to FOL

Note that function symbols give extra expressivity to FOL:

- ▶ The set of atomic formulas is extended to include expressions with function symbols.

Example: $\text{Older}(\text{father}(\text{max}), \text{max})$ is an atomic sentence.

- ▶ Functions can be nested. Note: predicates cannot be nested.

Example

$\text{father}(\text{mother}(\text{max}))$ is a term.

$\text{Loves}(\text{father}(\text{mother}(\text{max})), \text{mary})$ is a sentence.

$\text{Older}(\text{Loves}(\text{juliet}, \text{romeo}))$ is not well-formed.

- ▶ You can express more properties using function symbols.

Example: “Everyone’s paternal grandmother is older and nicer than one’s father”

$$\forall x (\text{OlderThan}(\text{mother}(\text{father}(x)), \text{father}(x)) \wedge \\ \text{NicerThan}(\text{mother}(\text{father}(x)), \text{father}(x)))$$

Example exercise from the 2019 exam

Given are the following five sets:

$$A = \{1, 2\}, \quad B = \{\{1\}, \{2\}\}, \quad C = \{1, \{1\}\}, \quad R = \{\langle 1, 2 \rangle, \langle 2, 1 \rangle\} \\ \text{and } S = \{1, \langle 2, 2 \rangle\}.$$

For each of the following statements, determine whether it is true or false. You are not required to explain the answer.

1. $C \subseteq A$
2. $\emptyset \in B$
3. $R \subseteq A$
4. $A \cap C \subseteq S$
5. $(A \cap C) \in (B \cap C)$
6. $C \subseteq A \cup B$
7. $A \cap S = \emptyset$
8. R is transitive

Next lecture

Wednesday 6 December 15:00 to 17:00

Coming soon

Quantification: meaning, translation, proof rules