

Question :-

Bigram -

what is the probability of word  $w$  given  
before word  $v$

$$P(w_1 | v) \rightarrow P(w_2 | w_1)$$

→ sometimes bigram won't even leading  
to 0 counts

→ for this we use interpolation

$$\text{unigram} + \text{bigram}$$

$$\rightarrow P(w_1 | v) = \lambda f(w_1 | v) + (1 - \lambda) f(w)$$

→ Directly modify counts in training data

3 cases

$$i) \quad c^*(v, w) = c(v, w) + \theta$$

$$ii) \quad c^*(v, w) = c(v, w) + \theta(v)$$

$$iii) \quad c^*(v, w) = c(v, w) + \theta(v) f(w)$$

so

smoothed bigram prob

$$p^*(w|v) = \frac{c^*(v, w)}{\sum_{w' \in V} c^*(v, w')}$$

$N(v, w)$  - held out data

i)  $\theta$  that maximizes the log-probability

$$p(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \log p^*(w|v)$$

Case 1 :-

$$p(w_i | w_{i-1}) = \frac{\text{count}(w_{i-1}, w_i)}{\text{count}(w_{i-1})}$$

$$p^*(w|v) = \frac{c^*(v, w)}{c^*(v)}$$

$$p(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \log \frac{c^*(v, w)}{c^*(v)}$$

$$\frac{d}{d\theta} p(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \frac{d}{d\theta} \log \frac{c^*(v, w) + \theta}{c^*(w) + \theta}$$

$$P(\theta) = \sum_{v \in V} \sum_{w \in V} N(v, w) \frac{1}{c^*(v, w) + \theta} - \frac{1}{c^*(v) + \theta}$$

set the derivative equal to 0 & solve for  $\theta$  to find critical points

$$\sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{c^*(v) + \cancel{\theta} - c^*(v, w) - \cancel{\theta}}{c^*(v, w) \cdot c^*(v) + c^*(v, w) \cdot \theta + \theta c^*(v) + \theta^2} \right) = 0$$

$$\sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{c^*(v) - c^*(v, w)}{c^*(v, w) \cdot c^*(v) + \theta (c^*(v, w) + c^*(v)) + \theta^2} \right) = 0$$

Now compute the second derivative of  $P(\theta)$  at the critical point  
if negative then critical point corresponds to local maximum

$$\frac{d}{d\theta} \sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{c^*(v) - c^*(v, w)}{c^*(v, w) \cdot c^*(v) + \theta (c^*(v, w) + c^*(v)) + \theta^2} \right)$$

$$= - \sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{c^*(v) - c^*(v, w)}{c^*(v, w) \cdot c^*(v) + \theta (c^*(v, w) + c^*(v)) + \theta^2} \right) \times \left[ 2\theta + (c^*(v, w) + c^*(v)) \right]$$

In the numerator we have 2 values which will definitely be  $\geq 0$

all the values are a positive number

so it is Maximum

Case 2:-

$$c^*(v, w) = c(v, w) + \theta c(w)$$

$$\sum_{w \in V} c^*(v, w)$$

$$= \sum_{w' \in V} c^*(v, w') + \theta c(w)$$

$$= c^*(v) + \theta \sum_{w' \in V} c(w')$$

$$= c^*(v) + N\theta$$

$\xrightarrow{\quad}$   $N$  is the length  
of vocabulary

$$P(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \log \left[ \frac{c^*(v, w) + \theta c(w)}{c^*(v) + N\theta} \right]$$

differentiate & equate to 0

$$\frac{dP(H)}{d\theta} = \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c(w)}{c^*(v, w) + \theta c(w)} - \frac{N}{c^*(v) + N\theta} \right] = 0$$

$$= \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c(w) c^*(v) + \cancel{\theta N c^*(w)} - N c^*(v, w) - \cancel{N\theta c^*(w)}}{c^*(v, w) \cdot c^*(v) + N c^*(v, w) \theta + N\theta^2 c(w) + \theta c(w) c(v)} \right]$$

$$= \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(w) c^*(v) - N c^*(v, w)}{c^*(v, w) \cdot c^*(v) + N c^*(v, w) \theta + N\theta^2 c(w) + \theta c(w) c(v)} \right]$$

Second differentiation

$$= \frac{d}{d\theta} \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(w) c^*(v) - N c^*(v, w)}{c^*(v, w) \cdot c^*(v) + N c^*(v, w) \theta + N\theta^2 c(w) + \theta c(w) c(v)} \right]$$

$$= - \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(w) c^*(v) - N c^*(v, w)}{N\theta^2 c(w) + (c^*(v, w) \cdot c^*(v))^2 + \theta (c^*(v, w) N + c^*(w) c(v))} \right]$$

$$\left[ N\theta c(w) + c^*(v, w) N + c^*(w) c(v) \right]$$

= it is - (number) so maximum

case 3:

$$c^*(v, w) = c^*(v, w) + \theta c^*(v) f(w)$$

$$p^*(w|v) = \frac{c^*(v, w)}{\sum_{w' \in V} c^*(v, w')}$$

$$\sum_{w' \in V} c^*(v, w') = \sum_{w' \in V} c(v, w') + \theta c^*(v) \frac{c^*(w)}{N}$$

$$= c^*(v) \left[ 1 + \theta \sum_{w' \in V} \frac{c^*(w)}{N} \right]$$

$$= c^*(v) \left[ 1 + \theta \frac{N}{N} \right]$$

$$= c^*(v) [1 + \theta]$$

Now substitute

$$P(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \log P^*(w|v)$$

$$= \sum_{v \in V} \sum_{w \in V} N(v, w) \log \frac{c(v, w) + \theta c(v) f(w)}{c(v) [1 + \theta]}$$

differentiate and equate to 0

$$\frac{d}{d\theta} P(H) = \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(v) f(w)}{c^*(v, w) + \theta c^*(v) f(w)} - \frac{c^*(v)}{c^*(v) [1 + \theta]} \right]$$

$$= \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(v)^2 f(w) [1 + \theta] - c^*(v) (c^*(v, w) + c^*(v)^2 f(w) \theta)}{c^*(v) c^*(v, w) + c^*(v) c^*(v, w) \theta + \theta c^*(v)^2 f(w) + \theta^2 c^*(v)^2 f(w)} \right]$$

$$= \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(v)^2 f(w) - c^*(v) c^*(v, w)}{\theta^2 c^*(v)^2 f(w) + \theta c^*(v) [c^*(v, w) + c^*(v) f(w)] + c^*(v) c^*(v, w)} \right]$$

differentiate again

differentiate again

$$= - \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{c^*(v)^2 f(w) - c^*(v) c^*(v, w)}{\theta^2 c^*(v)^2 f(w) + \theta c^*(v) [c^*(v, w) + c^*(v) f(w)] + c^*(v) c^*(v, w)} \right] \cdot [2\theta c^*(v)^2 f(w) + c^*(v) [c^*(v, w) + c^*(v) f(w)]]$$

its -ve and the value inside is +ve

so maximum

2) Case 1:-

$$c^*(v, w) = c^*(v, w) + \theta$$

here  $N(v, w) = c(v, w)$

so substitute here

$$p^*(w|v) = \frac{c(v, w) + \theta}{\sum_{w' \in V} [c(v, w') + \theta]}$$

$$= \frac{c(v, w) + \theta}{\sum_{w' \in V} c(v, w') + \theta |V|}$$

differentiate

$$\sum_{w' \in V} \frac{c(v, w') - c(v, w) |V|}{\left( \sum_{w' \in V} c(v, w') + \theta |V| \right)^2}$$

its constant

its +ve for any  
+ve  $\theta$



the  $\theta$  does not change the probabilities

Therefore  $\theta$  doesn't affect  $P(H)$  & optimal value is  $\theta = 0$

$$- \sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{(c^*(v) - c^*(v, w))}{(c^*(v, w) \cdot c^*(v) + \theta (c^*(v, w) + c^*(v)) + \theta^2)} \right) \times \left[ 2\theta + (c^*(v, w) + c^*(v)) \right]$$

from the above equation in question 4, after double differentiating wrt to  $\theta$  evaluated at  $\theta$  is negative, it means that  $\theta = 0$  is indeed a local maximum.

function is concave at that point

Given  $\theta = 0$  emerges as a local maxima, it reinforces the conclusion that  $\theta = 0$  is good when  $N(v, w) = c(v, w)$

Case 2:-

$$C^*(v, w) = C(v, w) + \theta C(w)$$

$$P^*(w|v) = \frac{C(v, w) + \theta C(w)}{\sum_{w' \in V} C(v, w') + \theta C(w)}$$

Based on the equation derived above

$$= - \sum_{v \in V} \sum_{w \in V} N(v, w) \left( \frac{(C^*(v) - C^*(v, w))}{C^*(v, w) \cdot C^*(v) + \theta (C^*(v, w) + C^*(v)) + \theta^2} \right) \times [2\theta + (C^*(v, w) + C^*(v))]$$

we can see that

the  $\theta$  does not change the probabilities

Therefore  $\theta$  doesn't affect  $P(H)$  & optimal value is  $\theta = 0$

Case 3:-

$$C^*(v, w) = C(v, w) + \theta C(v) f(w)$$

Based on above equation

$$= - \sum_{v \in V} \sum_{w \in V} N(v, w) \left[ \frac{C^*(v)^2 f(w) - C^*(v) C^*(v, w)}{\theta^2 C^*(v)^2 f(w) + \theta C^*(v) [C^*(v, w) + C^*(v) f(w)] + C^*(v) C^*(v)^2} \right] \cdot [2\theta C^*(v)^2 f(w) + C^*(v) [C^*(v, w) + C^*(v) f(w)]]$$

we can see that

the  $\theta$  does not change the probabilities

Therefore  $\theta$  doesn't affect  $P(H)$  & optimal value is  $\theta = 0$

3) case 1 :-

$$c^*(v, w) = c(v, w) + \theta$$

$$p^*(w|v) = \frac{c(v, w) + \theta}{\sum_{w' \in V} [c(v, w') + \theta]}$$

$$= \frac{c(v, w) + \theta}{c(v) + \theta}$$

$$= \frac{c(v, w)}{c(v) + \theta} + \frac{\theta}{c(v) + \theta}$$

$$f_2(w|v) = \frac{c(w, v)}{c(v)}$$

$$f_1(w) = \frac{c(w)}{N}$$

$$\gamma f(w|v) + (1-\gamma) f(w) = \frac{c(v,w)}{c(v) + \theta} + \frac{\theta}{c(v) + \theta}$$

$$= f(w|v) \left[ \frac{c(v)}{c(v) + \theta} \right] + \frac{\theta}{c(v) + \theta}$$

$$f(w|v) = \frac{c(w,v)}{c(v)} \quad \gamma = \frac{c(v)}{c(v) + \theta}$$

$$f(w) = 1$$

case 2:-

$$p^*(w|v) = \frac{c(v,w) + \theta c(w)}{c(v) + N\theta}$$

$$p^*(w|v) = \delta f_2(w|v) + (1-\delta) f_1(w)$$

$$\frac{c(v,w) + \theta c(w)}{c(v) + N\theta} = \delta f_2(w|v) + (1-\delta) f_1(w)$$

$$f(w|v) = \frac{c(w,v)}{c(v)} + f(w) \cdot \frac{c(w)}{N}$$

$$\frac{c(v,w)}{c(v) + N\theta} + \frac{\theta c(w)}{c(v) + N\theta}$$

$$= \delta f(w|v) + (1-\delta) f(w)$$

$$= c(w,v) = f(w|v) \cdot c(v)$$

$$= c(w) = f(w) \cdot n$$

$$= f(w|v) \left[ \frac{c(v)}{c(v) + N\theta} \right] + \frac{\theta n f(w)}{c(v) + N\theta}$$

$$= \lambda f(w|v) + (1-\lambda) f(w)$$

$$\lambda = \frac{c(v)}{c(v) + N\theta}$$

$$f(w|v) = \frac{c(w,v)}{c(v)}$$

$$f(w) = \frac{c(w)}{n}$$

Case 3:-

$$p^*(w|v) = \frac{c(v,w) + \theta c(v) f(w)}{c(v) [1 + \theta]}$$

$$= \frac{c(v,w)}{c(v) [1 + \theta]} + \frac{\theta c(v) f(w)}{c(v) [1 + \theta]}$$

$$= \frac{c(v) f(w/v)}{c(v) [1+\theta]} + \frac{\theta c(v) f(w)}{c(v) [1+\theta]}$$

$$\delta = \frac{1}{1+\theta}$$

$$f(w) = f(w)$$

$$f(w/v) = \frac{c(v, w)}{c(v)}$$