

Lecture 4c. Training logistic regression with the IRLS algorithm


COMP90051 Statistical Machine Learning

Semester 2, 2020
Lecturer: Ben Rubinstein



THE UNIVERSITY OF
MELBOURNE

This lecture

- Iterative optimisation for extremum estimators
 - * First-order method: Gradient descent
 - * Second-order: Newton-Raphson method
 -  Later: Lagrangian duality
- Logistic regression: workhorse linear classifier
 - * Possibly familiar derivation: frequentist
 - * Decision-theoretic derivation
 - * **Training with Newton-Raphson** looks like repeated, weighted linear regression

Training Logistic Regression: the IRLS Algorithm

Analytical? Newton-Raphson!


Iterative optimisation

- Training logistic regression: \mathbf{w} maximising log-likelihood $L(\mathbf{w})$ or cross-entropy loss
- **Bad news:** No closed form solution
- **Good news:** Problem is strictly convex, if no irrelevant features \rightarrow convergence!

 Look ahead: regularisation for irrelevant features

How does gradient descent work?

- $\mu(z) = \frac{1}{1+\exp(-z)}$ then $\frac{d\mu}{dz} = \mu(z)(1 - \mu(z))$
- Then $\nabla L(\mathbf{w}) = \sum_{i=1}^n (y_n - \mu(\mathbf{x}_n)) \mathbf{x}_n = \mathbf{X}'(\mathbf{y} - \boldsymbol{\mu})$, stacking instances in \mathbf{X} , labels in \mathbf{y} , $\mu(\mathbf{x}_n)$ in $\boldsymbol{\mu}$



Murphy, Fig 8.3, p247

Note I'm abusing notation:
 $\mu(\mathbf{x}_n) = \mu(z)$ where $z = \mathbf{w}'\mathbf{x}_n$
 Meaning by input type

Iteratively-Reweighted Least Squares

- Instead of GD, let's apply Newton-Raphson → **IRLS algorithm**
- Recall: $\nabla L(\mathbf{w}) = \mathbf{X}'(\mathbf{y} - \boldsymbol{\mu})$. Differentiate again for **Hessian**:

$$\nabla_2 L(\mathbf{w}) = -\sum_i \frac{d\mu}{dz_i} \mathbf{x}_n \mathbf{x}_n' = -\sum_i \mu(\mathbf{x}_i)(1 - \mu(\mathbf{x}_i)) \mathbf{x}_n \mathbf{x}_n'$$

$$= -\mathbf{X}'\mathbf{M}\mathbf{X}, \text{ where } M_{ii} = \mu_i(1 - \mu_i) \text{ otherwise } 0$$
- **Newton-Raphson** then says (now with $\mathbf{M}_t, \boldsymbol{\mu}_t$ dependence on \mathbf{w}_t)

$$\mathbf{w}_{t+1} = \mathbf{w}_t - (\nabla_2 L)^{-1} \nabla L = \mathbf{w}_t + (\mathbf{X}'\mathbf{M}_t\mathbf{X})^{-1} \mathbf{X}'(\mathbf{y} - \boldsymbol{\mu}_t)$$

$$= (\mathbf{X}'\mathbf{M}_t\mathbf{X})^{-1} [\mathbf{X}'\mathbf{M}_t\mathbf{X}\mathbf{w}_t + \mathbf{X}'(\mathbf{y} - \boldsymbol{\mu}_t)]$$

$$= (\mathbf{X}'\mathbf{M}_t\mathbf{X})^{-1} \mathbf{X}'\mathbf{M}_t \mathbf{b}_t, \text{ where } \mathbf{b}_t = \mathbf{X}\mathbf{w}_t + \mathbf{M}_t^{-1}(\mathbf{y} - \boldsymbol{\mu}_t)$$

Compare to
normal
equations
- Each IRLS iteration solves a least squares problem weighted by \mathbf{M}_t , which are reweighted iteratively!

IRLS intuition: Putting labels on linear scale

$$\begin{aligned} \text{IRLS: } \mathbf{w}_{t+1} &= (\mathbf{X}'\mathbf{M}_t\mathbf{X})^{-1}\mathbf{X}'\mathbf{M}_t\mathbf{b}_t \\ \text{where } \mathbf{b}_t &= \mathbf{X}\mathbf{w}_t + \mathbf{M}_t^{-1}(\mathbf{y} - \boldsymbol{\mu}_t) \\ \text{and } M_{ii} &= \mu_t(\mathbf{x}_i)[1 - \mu_t(\mathbf{x}_i)] \text{ otherwise } 0 \\ \text{and } \mu_t(\mathbf{x}) &= [1 + \exp(-\mathbf{w}_t'\mathbf{x})]^{-1} \end{aligned}$$

- The \mathbf{y} are not on linear scale. Invert logistic function?
- The \mathbf{b}_t are a “linearised” approximation to these: the \mathbf{b}_t equation matches a linear approx. to $\mu_t^{-1}(\mathbf{y})$.
- Linear regression on new labels!

IRLS intuition: Equalising label variance

IRLS: $\mathbf{w}_{t+1} = (\mathbf{X}'\mathbf{M}_t\mathbf{X})^{-1}\mathbf{X}'\mathbf{M}_t\mathbf{b}_t$
where $\mathbf{b}_t = \mathbf{X}\mathbf{w}_t + \mathbf{M}_t^{-1}(\mathbf{y} - \boldsymbol{\mu}_t)$
and $M_{ii} = \mu_t(\mathbf{x}_i)[1 - \mu_t(\mathbf{x}_i)]$ otherwise 0
and $\mu_t(\mathbf{x}) = [1 + \exp(-\mathbf{w}_t'\mathbf{x})]^{-1}$

- In linear regression, each y_i has equal variance σ^2
- Our y_i are Bernoulli, variance: $\mu_t(\mathbf{x}_i)[1 - \mu_t(\mathbf{x}_i)]$
- Our reweighting standardises, dividing by variances!!

Fun exercise: Show that Newton-Raphson for linear regression gives you the normal equations!

Summary

- Training logistic regression
 - * No analytical solution
 - * Gradient descent possible, but convergence rate not ideal
 - * Newton-Raphson: iteratively reweighted least squares

Next time: Regularised linear regression for avoiding overfitting and ill-posed optimisation