



Congratulations! You passed!
 Grade received 100%
 Latest Submission Grade 100%
 To pass 70% or higher

[Go to next item](#)

1. Gradient descent is an algorithm for finding values of parameters w and b that minimize the cost function J .

1 / 1 point

repeat until convergence {

$$w = w - \alpha \frac{\partial}{\partial w} J(w, b)$$

$$b = b - \alpha \frac{\partial}{\partial b} J(w, b)$$

When $\frac{\partial J(w, b)}{\partial w}$ is a negative number (less than zero), what happens to w after one update step?



w increases.



w stays the same



w decreases



It is not possible to tell if w will increase or decrease.



Correct

The learning rate is always a positive number, so if you take w minus a negative number, you end up with a new value for w that is larger (more positive).

2. For linear regression, what is the update step for parameter b ?

1 / 1 point



$$b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)}) x^{(i)}$$



$$b = b - \alpha \frac{1}{m} \sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$$



Correct

The update step is $b = b - \alpha \frac{\partial J(w, b)}{\partial b}$ where $\frac{\partial J(w, b)}{\partial b}$ can be computed with this expression: $\sum_{i=1}^m (f_{w,b}(x^{(i)}) - y^{(i)})$