

Solutions Sheet 3

Nina Fischer and Yannick Zelle

October 30, 2021

Exercise 1

Task: Describe the maximum likelihood estimator for the following distributions:

(a)

$$\mathcal{N}(x \mid \mu, \sigma^2)$$

We denote with f_μ the PDF of the normal distribution. By using the Lemma 6.2.3 we can write the log-likelihood function as:

$$\begin{aligned} l(\mu) &= \ln\left(\prod_{i=1}^n f_\mu\right) \\ &= \ln\left(\prod_{i=1}^n e^{a+\eta x_i - \frac{1}{2}\lambda^2 x_i^2}\right) \\ &= \sum_{i=1}^n \ln(e^{a+\eta x_i - \frac{1}{2}\lambda^2 x_i^2}) \\ &= \sum_{i=1}^n -\frac{1}{2}(\log(2\pi) - \log(\lambda^2) + \frac{\mu^2}{\sigma^2}) + \frac{\mu}{\sigma^2}x_i - \frac{1}{2}\lambda^2 x_i^2 \end{aligned}$$

To find μ which maximizes $l(\mu)$ we calculate:

$$\begin{aligned} \frac{\delta l(\mu)}{\delta \mu} &= \sum_{i=1}^n -\frac{\mu}{\sigma^2} + \frac{1}{\sigma^2}x_i \\ &= -n\frac{\mu}{\sigma^2} + \frac{1}{\sigma^2}\sum_{i=1}^n x_i \end{aligned}$$

$l(\mu)$ is maximal if :

$$\begin{aligned}
l(\mu) = 0 &\leftrightarrow 0 = -n \frac{\mu}{\sigma^2} + \frac{1}{\sigma^2} \sum_{i=1}^n x_i \\
&\leftrightarrow \mu = \frac{1}{n} \sum_{i=1}^n x_i
\end{aligned}$$

Since $\frac{\delta^2 l(\mu)}{\delta^2 \mu} = -\frac{1}{\sigma^2} < 0$, with $\mu_o := \frac{1}{n} \sum_{i=1}^n x_i$ we obtain a global maximum for $l(\mu)$ and we chose μ_0 therefore as our estimator.

(b)

Exponential distribution with the density function $f(x|\lambda) = \lambda e^{-\lambda x}$ with $\lambda > 0$ and observations $x_i \geq 0$. We have :

$$\begin{aligned}
f(x|\lambda) &= \lambda e^{-\lambda x} \\
&= e^{-\lambda x + \ln(\lambda)}
\end{aligned}$$

and the likelihood function:

$$L(\lambda) = \prod_{i=1}^n f(x_i|\lambda)$$

Therefore the log-likelihood is giveb by:

$$\begin{aligned}
l(\lambda) &= \ln\left(\prod_{i=1}^n f(x_i|\lambda)\right) \\
&= \sum_{i=1}^n \ln(f(x_i|\lambda)) \\
&= \sum_{i=1}^n \ln(e^{-\lambda x_i + \ln(\lambda)}) \\
&= \sum_{i=1}^n (-\lambda x_i + \ln(\lambda)) \\
&= -\lambda \sum_{i=1}^n x_i + n \ln(\lambda)
\end{aligned}$$

With the same argument as in (a) we derive:

$$\begin{aligned}
\frac{\delta l(\lambda)}{\delta \lambda} &= \sum_{i=1}^n (-x_i) + \frac{n}{\lambda} \\
&= -\sum_{i=1}^n x_i + \frac{n}{\lambda}
\end{aligned}$$

Then we have :

$$0 = \frac{\delta l(\lambda)}{\delta \lambda} \leftrightarrow \lambda = \frac{n}{\sum_{i=1}^n x_i}$$

We also have

$$\frac{\delta^2 l(\lambda)}{\delta^2 \lambda} = -\frac{1}{\lambda^2} < 0$$

therefore we chose $\lambda_0 = \frac{n}{\sum_{i=1}^n x_i}$ as our estimator.

Exercise 2

Exercise 2