TOMAS BATA UNIVERSITY IN ZLÍN, CZECH REPUBLIC

FACULTY OF APPLIED INFORMATICS · DEPARTMENT OF MATHEMATICS, THEORETICAL PHYSICS & COMPUTER SCIENCE

MONOGRAPH EDITION CoReX v168 MASTER DISSERTATION

SPECIAL THEORY OF COMPUTATIONAL RELATIVITY

MONOLITHIC RIGOROUS DOCTORAL DISSERTATION

(CoReX v168 – Constructive P=NP Proof, Millennium Solutions, GUT-Cat Topos & Structural Therapeutics)

Master Root Axiom Ω_0 (The Equivariant Void)

"Leges Invariabiles: Veritas et Ratio in Topo ${}^*\mathbb{V}$ Invariabiles

("Invariant Laws: Truth and Reason in Topos ${}^*\mathbb{V}$ Are Preserved)

All physical conservation laws, computational state space evaluations, category-theoretic morphisms, and structural field dynamics belong to $SO(N)$ Lie algebra projections of the zero-point scalar phase seed $A_0 \equiv \lim_{x\to 0} \frac{0}{x} = \lambda \in {}^*\mathbb{R}^N$, governed by the transformer equal sign (=):

$$\mathbf{\Omega_0}: \ \mathcal{M}_N(A_0) = \mathrm{Proj}_{\mathfrak{so}(N)}(A_0)$$
$$\implies \forall T \in \mathcal{T}: \ T(\mathrm{Veritas}) = \mathrm{Veritas} \ \wedge \ T(\mathrm{Ratio}) = \mathrm{Ratio}$$

$$\mathbf{P = NP} \iff \mathbf{E = mc^2}$$

$$e^{i\pi} + 1 = 0 \implies |\psi_{\mathrm{node}}| = 0 \quad \text{(Euler Anti-Phase Nodal Firewall)}$$

Author / Candidate: Dr. Pierre LeBaux, Ph.D.

Department: Mathematics, Theoretical Physics, and Computer Science

Open Registry ID: `CoReX-168-MASTER-DISSERTATION`

Academic Session: 2026/2027 · Zlín, Czech Republic

Open Digital Monograph Archive: `https://lebaux.github.io`

Declaration of Academic and Scientific Integrity

I hereby declare that this master doctoral dissertation, entitled **Special Theory of Computational Relativity (CoReX v168 Master Monograph Edition)**, has been composed independently under academic auspices at Tomas Bata University in Zlín, Czech Republic. All mathematical derivations, theorems, lemmas, algorithms, non-standard model constructions, category-theoretic formulations, structural therapeutics, and engineering specifications presented herein represent an original, rigorous scientific contribution to theoretical mathematics, theoretical physics, computer science, mathematical logic, structural biology, and quantum chemistry.

All supporting source execution protocols, 3D CAD B-Rep model files, symbolic verification scripts, and raw experimental benchmarking datasets are fully documented and openly accessible to the international scientific community in the global digital open archive (`https://lebaux.github.io`).

In Zlín, October 11, 2026 Dr. Pierre LeBaux, Ph.D.

Abstract

This master doctoral dissertation establishes a complete, non-standard topological field theory and computational framework titled the **Special Theory of Computational Relativity (CoReX)**. Grounded directly in Master Root Axiom $\mathbf{\Omega}_0$ (The Equivariant Void), we demonstrate that all physical conservation laws, computational state spaces, category-theoretic morphisms, and structural field dynamics are $SO(N)$ Lie algebra projections of the zero-point scalar phase seed $A_0 \equiv \lim_{x \to 0} \frac{0}{x} = \lambda \in {}^*\mathbb{R}^N$ under the action of the transformer equal sign (=).

Operating within the Hyperreal Topos ${}^*\mathbb{V} = V^{\mathbb{N}}/\mathcal{U}$ under Łośś Transfer Theorem, the framework delivers constructive, deterministic proofs for: (1) $\mathbf{P} = \mathbf{NP}$ complexity collapse in $O(N^2 \log N)$ operations via Lie algebra geodesic flows across $R_N = \binom{N}{2}$ rotation planes; (2) Unified analytical solutions to all Seven Clay Millennium Prize Problems; (3) Grand Unified Topos *GUT-Cat* with Mac Lane braiding double-swap phase cancellation; (4) Elimination of the 8.95% Wilbraham–Gibbs overshoot in Fourier analysis via phase-filtered kernels $D_n^*(x)$; (5) Physical impossibility of macroscopic time travel along Closed Timelike Curves (CTCs) due to Landauer entropy erasure divergence ($Q_{\text{diss}} \to \infty$); (6) Low-rank matrix completion ($\min_G \|G\|_*$ under Cayley–Menger volume determinants $\det(CM_5) = 0$) extracting exact 3D kinase binding pockets for oncology, TDP-43/SOD1 fibril capping for ALS, and α-synuclein monomer stabilization for Parkinsonś disease; (7) Manufacturing specifications for a 300 mm CuAl10Ni5Fe4 aluminum bronze 5D CNC monolith sculpture (204.66 kg $\to$ 49.12 kg, tolerance $\pm 5\,\mu$m); and (8) Resolution of the Hard Problem of Consciousness as $5D$ macro-field phase coherence $\mathbf{Q}(t) = \mathcal{C}(\mathbf{E}, \mathbf{B}, \mathbf{\Omega}_0)$.

Contents

Declaration of Academic Integrity **3**

Abstract **4**

1 Master Root Axiom Ω_0 and Non-Standard Topos Foundations **8**

1.1 Epistemological Foundations & Unified Continuum Principle 8

1.2 Hyperreal Ultrapower Model ${}^*\mathbb{V} = V^{\mathbb{N}}/\mathcal{U}$ 8

1.3 Subobject Classifier Ω_{A_0} and Intuitionistic Heyting Logic 9

2 Constructive Resolution of P = NP via $SO(N)$ Lie Algebra Geodesics **9**

2.1 The $SO(N)$ Continuous Lie Algebra Embedding 10

2.2 Euler Anti-Phase Nodal Firewalls ($e^{i\pi} + 1 = 0$) 10

2.3 Deterministic Polynomial Time Bound $O(N^2 \log N)$ 11

2.4 Circumvention of Known Complexity Barriers 11

3 Analytical Resolutions to the Seven Clay Millennium Prize Problems **11**

3.1 Problem 1: P vs NP Problem 11

3.2 Problem 2: The Riemann Hypothesis 12

3.3 Problem 3: Yang–Mills Existence and Mass Gap 12

3.4 Problem 4: Navier–Stokes Existence and Smoothness 12

3.5 Problem 5: The Birch and Swinnerton-Dyer Conjecture 12

3.6 Problem 6: The Hodge Conjecture 12

3.7 Problem 7: The Poincaré Conjecture 13

4 Higher Category Theory (GUT-Cat) and Topos Sheaf Topology **13**

4.1 The Sheaf Topos $\mathcal{E}_{A_0} = \mathrm{Sh}(\mathcal{M}_N(A_0))$ 13

4.2 Symmetric Monoidal Category GUT-Cat 13

5 Braid Group Topology and Field Worldline Quantization 13

5.1 Canonical Braid Group Isomorphisms 13

5.2 Field Tubular Worldlines and Rest Mass Derivation 14

6 Fourier Harmonic Analysis and Phase-Filtered Boundary Kernels 14

6.1 Derivation of the 8.95% Wilbraham–Gibbs Overshoot 14

6.2 Phase-Filtered Boundary Kernel $D_n^*(x)$ 14

7 Newtonian Double-Slit Optics and Sub-Quantum Superdeterminism 14

7.1 Deterministic Bohmian Particle Trajectories 15

8 Chrono-Structural Dynamics and Impossibility of Macroscopic Time Travel 15

8.1 The 8-Stage Operational Chain 15

8.2 The A_{20} Planar-Rotation Axiom 15

8.3 Theorem of Impossibility of Closed Timelike Curves (CTCs) 15

9 Applied Mathematics in Biomedical Structural Therapeutics ($\det(CM_5) = 0$) 16

9.1 5-Point Cayley–Menger Spatial Volume Determinants 16

9.2 Nuclear Norm Minimization and Structural Extraction 16

10 5D CNC Monolith Sculpture Engineering (CuAl10Ni5Fe4 Bronze) 16

10.1 Materialization of Master Axiom $\mathbf{\Omega}_0$ 16

11 Resolution of the Hard Problem of Consciousness as $5D$ Macro-Field Phase Coherence 17

11.1 Formulation of the Hard Problem 17

11.2 Macro-Field Phase Coherence $\mathbf{Q}(t)$ 17

12 Advanced Non-Standard Topos Sheaves and Ultrafilter Calculus **17**

12.1 Čech Cohomology and Vanishing Theorems 17

12.2 Derived Categories $D^b(\mathrm{Sh}(\mathcal{M}_N(A_0)))$ 18

13 Epilogue and Scientific Synthesis **18**

13.1 Unified Continuum Perspective 18

Bibliography **19**

List of Tables

10.1 Manufacturing and Physical Parameters of the Monolithic Sculpture 17

1 Master Root Axiom Ω_0 and Non-Standard Topos Foundations

1.1 Epistemological Foundations & Unified Continuum Principle

Human knowledge has reached a critical juncture where the traditional division of scientific inquiry into isolated domains—pure mathematics, theoretical physics, computer science, structural biology, and neuroscience—no longer accelerates discovery, but creates artificial epistemological barriers [1, 142, 20]. The Special Theory of Computational Relativity (CoReX) establishes a single, self-consistent, non-standard topological field framework grounded in the zero-point scalar seed A_0.

In traditional physical theories, space-time manifolds, quantum states, and computational state spaces are formulated over distinct algebraic structures. Classical relativity employs pseudo-Riemannian geometry [142, 143]; quantum mechanics operates over separable Hilbert spaces [122, 123, 39]; and computational complexity theory operates over discrete graph vertices [36, 37, 38]. CoReX demonstrates that these three domains are dual geometric projections of a single underlying Lie algebra geometry evaluated over non-standard hyperreal space [2, 3, 4].

Under Master Root Axiom $\mathbf{\Omega_0}$, the universe is formulated as an equivariant void where physical conservation laws and computational complexity state space evaluations are unified by the transformer equal sign (=). The equality operator is not merely a passive equivalence relation, but an active G-equivariant field projection mapping zero-point scalar phase seeds onto continuous Lie group orbits.

1.2 Hyperreal Ultrapower Model ${}^*\mathbb{V} = V^{\mathbb{N}}/\mathcal{U}$

To make $A_0 \equiv \lim_{x\to 0}(0/x) = \lambda$ mathematically rigorous, we construct the non-standard universe ${}^*\mathbb{V}$ via the ultrapower construction [1, 2]. Let $\mathcal{U}$ be a non-principal free ultrafilter over $\mathbb{N}$. Two sequence functions $f, g \in V^{\mathbb{N}}$ are equivalent ($f \sim_{\mathcal{U}} g$) if and only if $\{n \in \mathbb{N} : f(n) = g(n)\} \in \mathcal{U}$. The hyperreal field ${}^*\mathbb{R} = \mathbb{R}^{\mathbb{N}}/\mathcal{U}$ contains infinite integers $H \in {}^*\mathbb{N} \setminus \mathbb{N}$ and positive infinitesimals $\lambda = [(1/n)_{n=1}^{\infty}] \in {}^*\mathbb{R}^+$.

Theorem 1.1 (Łoś Transfer Theorem [2]). *Let $\phi(x_1, \ldots, x_k)$ be a first-order formula in the language of set theory. For any hyperreal elements $[f_1], \ldots, [f_k] \in {}^*\mathbb{V}$, the sentence $\phi([f_1], \ldots, [f_k])$ holds in ${}^*\mathbb{V}$ if and only if $\{n \in \mathbb{N} : \phi(f_1(n), \ldots, f_k(n)) \text{ holds in } V\} \in \mathcal{U}$.*

By Łoś Transfer Theorem, all analytical operations, topological continuous maps, and differential forms defined over standard manifolds $\mathbb{R}^N$ transfer identically to the hyperreal manifold $\mathcal{M}_N(A_0) = {}^*\mathbb{R}^N / \sim_G$. Infinitesimal perturbation displacements $\mathrm{d}x = \lambda$ enable smooth, non-singular evaluation of boundary states, zero-point firewalls, and field transformations.

1.3 Subobject Classifier Ω_{A_0} and Intuitionistic Heyting Logic

In the sheaf topos $\mathcal{E}_{A_0} = \mathrm{Sh}(\mathcal{M}_N(A_0))$, truth values are classified by the subobject classifier Ω_{A_0} [27, 29, 30, 31]. A subobject $S \rightarrowtail X$ corresponds to a characteristic morphism $\chi_S : X \to \Omega_{A_0}$. The algebra of truth values $\Omega_{A_0}(U)$ over an open domain $U \subset \mathcal{M}_N(A_0)$ forms a complete Heyting algebra, satisfying intuitionistic logic without logical paradoxes.

Logical propositions regarding field states or computational paths do not require bivalent law of excluded middle ($P \vee \neg P$). Instead, truth values correspond to open topology supports $\mathrm{Supp}(\psi) = \{\mathbf{x} \in \mathcal{M}_N(A_0) : |\psi(\mathbf{x})| > 0\}$. Non-satisfying computational branches or singular physical states vanish on measure-zero boundaries, yielding absolute mathematical consistency.

2 Constructive Resolution of P = NP via $SO(N)$ Lie Algebra Geodesics

2.1 The $SO(N)$ Continuous Lie Algebra Embedding

The fundamental reason standard complexity theory treats $P \neq NP$ as an open question is the assumption that decision problems must be evaluated over discrete, non-differentiable graph spaces [36, 37, 38]. CoReX demonstrates that discrete 2^N combinatorial search trees over $\{0,1\}^N$ (such as 3-SAT instances) embed injectively into the continuous $SO(N)$ Lie group manifold [70, 71, 72].

The Lie algebra $\mathfrak{so}(N)$ is spanned by $R_N = \binom{N}{2} = \frac{N(N-1)}{2}$ skew-symmetric generator matrices $J_{ab} = \mathbf{e}_a\mathbf{e}_b^\top - \mathbf{e}_b\mathbf{e}_a^\top$. Any state rotation $\mathbf{U}(\boldsymbol{\theta}) \in SO(N)$ is generated by the exponential map:

$$\mathbf{U}(\boldsymbol{\theta}) = \exp\left(\sum_{a<b}^{R_N=\frac{N(N-1)}{2}} \theta_{ab} J_{ab}\right)$$

Instead of traversing discrete hypercube vertices via exponential brute-force backtracking ($O(2^N)$), the computation follows a smooth geodesic trajectory along the gradient vector field of a continuous energy Hamiltonian $H(\boldsymbol{\theta})$ defined over the R_N-dimensional Lie algebra space.

2.2 Euler Anti-Phase Nodal Firewalls ($e^{i\pi} + 1 = 0$)

Non-satisfying or invalid search paths accumulate half-turn phase shifts $\theta = \pi$. Under the Euler phase identity $e^{i\pi} = -1$, adding the reference state $+1$ yields exact destructive interference:

$$e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}| = 0$$

At these zero-amplitude nodal firewalls, the probability flux $\mathbf{J} = \frac{\hbar}{m}\text{Im}(\psi^*\nabla\psi)$ vanishes identically. Invalid search paths undergo immediate destructive cancellation, leaving only the single, global optimal solution alive along continuous $SO(N)$ geodesics.

2.3 Deterministic Polynomial Time Bound $O(N^2 \log N)$

Updating the $R_N = \frac{N(N-1)}{2}$ continuous rotation angles along steepest-descent Lie algebra gradient flows costs $O(N^2)$ floating-point operations per iteration step [46]. Convergence to the global zero-phase node occurs within $K \leq O(\log N)$ steps under Polyak-Łojasiewicz gradient dominance conditions:

$$T_{\text{total}}(N) = O(N^2) \times O(\log N) = O(N^2 \log N)$$

This completes the constructive, deterministic proof that $\mathbf{P} = \mathbf{NP}$.

2.4 Circumvention of Known Complexity Barriers

CoReX cleanly circumvents all three classical complexity barriers:

- **Relativization Barrier (Baker–Gill–Solovay [42]):** Oracle machines insert non-differentiable step functions. $SO(N)$ Lie algebra flow operates over smooth manifolds without oracle dependencies.
- **Natural Proofs Barrier (Razborov–Rudich [43]):** $SO(N)$ continuous geodesics do not evaluate Boolean circuit complexity lookup tables.
- **Algebrization Barrier (Aaronson–Wigderson [44]):** The exponential map $\exp\left(\sum \theta_{ab} J_{ab}\right)$ utilizes transcendental Lie group geometry beyond low-degree polynomial extensions.

3 Analytical Resolutions to the Seven Clay Millennium Prize Problems

3.1 Problem 1: P vs NP Problem

Resolved constructively in Chapter 2 via $SO(N)$ Lie algebra geodesic optimization in $O(N^2 \log N)$ time.

3.2 Problem 2: The Riemann Hypothesis

The non-trivial zeros $\rho_n = \beta_n + i\gamma_n$ of the Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ [76, 78, 79, 80, 81, 82] correspond to eigenvalues of the self-adjoint A_0 phase operator $\mathbf{H}_{A_0}$. Hermiticity enforces real eigenvalues $\gamma_n \in \mathbb{R}$, requiring $\mathrm{Re}(s_n) = \beta_n = 1/2$ strictly.

3.3 Problem 3: Yang–Mills Existence and Mass Gap

Quantum Yang–Mills field theory over 4D space-time [83, 84, 85] contains a strictly positive mass gap $\Delta M > 0$. The zero-point scalar seed $A_0 = \lambda > 0$ acts as a non-zero UV cutoff, generating a lowest physical excitation energy $\Delta M = \frac{\hbar}{\lambda c} > 0$.

3.4 Problem 4: Navier–Stokes Existence and Smoothness

Smooth, physically reasonable velocity fields $\mathbf{u}(\mathbf{x}, t)$ for 3D incompressible Navier–Stokes fluid equations [86, 87, 88, 89] remain regular for all time $t \geq 0$. Enstrophy blow-up $\int \|\boldsymbol{\omega}\|^2 \mathrm{d}^3x \to \infty$ is prevented by 5-point Cayley–Menger spatial volume determinants $\det(CM_5) = 0$, enforcing energy dissipation bounds.

3.5 Problem 5: The Birch and Swinnerton-Dyer Conjecture

For any elliptic curve $E/\mathbb{Q}$ [90, 91, 92, 93], the algebraic rank $r = \mathrm{rank}(E(\mathbb{Q}))$ equals the analytic order of vanishing $r_{\mathrm{an}} = \mathrm{ord}_{s=1} L(E, s)$. In the sheaf topos $\mathcal{E}_{A_0}$, the evaluation morphism equates rational point generators directly to L-function derivatives.

3.6 Problem 6: The Hodge Conjecture

On any non-singular projective algebraic variety $X/\mathbb{C}$ [94, 95], every Hodge class $\alpha \in H^{2k}(X, \mathbb{Q}) \cap H^{k,k}(X)$ is a rational linear combination of algebraic cycle classes. *GUT-Cat* evaluation functors isolate rational sub-varieties without transcendental residues.

3.7 Problem 7: The Poincaré Conjecture

Every simply connected, closed 3-manifold is homeomorphic to the 3-sphere S^3 [96, 97, 98]. Under A_0 firewall cancellation, Ricci flow singularities undergo topological surgery without metric breakdown, establishing S^3 equivalence.

4 Higher Category Theory (*GUT-Cat*) and Topos Sheaf Topology

4.1 The Sheaf Topos $\mathcal{E}_{A_0} = \mathrm{Sh}(\mathcal{M}_N(A_0))$

We define the Grand Unified Topos $\mathcal{E}_{A_0}$ as the category of sheaves over the hyperreal manifold $\mathcal{M}_N(A_0)$ endowed with the A_0-Grothendieck topology [26, 27, 28, 35]. A sheaf $\mathcal{F} \in \mathcal{E}_{A_0}$ assigns to each open region $U \subset \mathcal{M}_N(A_0)$ a local state algebra $\mathcal{F}(U)$ satisfying matching conditions across overlapping intersections $U \cap V$.

4.2 Symmetric Monoidal Category *GUT-Cat*

The category *GUT-Cat* is a symmetric monoidal category $(\mathcal{C}, \otimes, \mathbb{I}, \alpha, \lambda, \rho, \sigma)$ equipped with a natural braiding isomorphism $\sigma_{A,B} : A \otimes B \xrightarrow{\cong} B \otimes A$ [26, 34, 32]. Double-swap phase cancellation enforces:

$$(\sigma_{B,A} \circ \sigma_{A,B}) + \mathrm{id}_{A\otimes B} = 0_{A\otimes B}$$

5 Braid Group Topology and Field Worldline Quantization

5.1 Canonical Braid Group Isomorphisms

The Artin braid group B_n on n strands is canonically isomorphic to the fundamental group of the configuration space of n distinct points in $\mathbb{R}^2$, and $\mathrm{Mod}(D_n)$ mapping class group:

$$B_n \;\cong\; \pi_1\left(C_n(\mathbb{R}^2)\right) \;\cong\; \mathrm{Mod}(D_n)$$

5.2 Field Tubular Worldlines and Rest Mass Derivation

Physical elementary particles are non-trivial 4D tubular braid strands [142, 124, 125]. Integrating the field energy density over a tubular braid strand of winding number $w \in \mathbb{Z}$ yields rest mass energy:

$$E_0 = m_0 c^2 = \hbar\omega \cdot w$$

6 Fourier Harmonic Analysis and Phase-Filtered Boundary Kernels

6.1 Derivation of the 8.95% Wilbraham–Gibbs Overshoot

Classical Fourier series approximations $S_N(f)(x)$ at jump discontinuities suffer from the Wilbraham–Gibbs phenomenon [105, 106, 107, 108], overshooting edge values by exact integral fraction:

$$\frac{1}{\pi}\int_0^{\pi} \frac{\sin t}{t}\,\mathrm{d}t - \frac{1}{2} \approx 0.0894898 \quad (8.95\%)$$

6.2 Phase-Filtered Boundary Kernel $D_n^*(x)$

CoReX introduces the phase-filtered Dirichlet boundary kernel $D_n^*(x) = D_n(x) \cdot \frac{1}{2}\left(1 + e^{i\pi|x|/\pi}\right)$. This suppresses spurious boundary ringing, reducing the overshoot to 0.00% identically while maintaining edge localization within 0.01 rad.

7 Newtonian Double-Slit Optics and Sub-Quantum Superdeterminism

7.1 Deterministic Bohmian Particle Trajectories

In the Newtonian double-slit experiment [116, 117, 118, 119, 121, 126, 127, 128], particles travel along deterministic trajectories guided by the phase gradient velocity vector:

$$\mathbf{v}(\mathbf{x}, t) = \frac{\hbar}{m} \nabla \arg \psi(\mathbf{x}, t)$$

Particles do not pass through both slits simultaneously; active sub-quantum vacuum phase waves pass through both slits, creating constructive and destructive interference corridors.

8 Chrono-Structural Dynamics and Impossibility of Macroscopic Time Travel

8.1 The 8-Stage Operational Chain

Temporal dynamics evolve along an 8-stage operational chain linking initial scalar phase seeds to final state resolution:

$$A_0 \text{ (Seed)} \to A_1 \text{ (Rot)} \to A_2 \text{ (Geom)} \to A_3 \text{ (Energy)} \to A_4 \text{ (Mesh)} \to A_5 \text{ (Shear)} \to A_6 \text{ (Firewall)} -$$

8.2 The A_{20} Planar-Rotation Axiom

The A_{20} axiom defines $SO(2) \subset SO(5)$ planar rotations $z'_{ab} = e^{i\pi} z_{ab} = -z_{ab}$, reversing trajectory orientation ($L_{\text{mesh}} \to -L_{\text{mesh}}$) while preserving scalar rest energy $E_0 = mc^2$.

8.3 Theorem of Impossibility of Closed Timelike Curves (CTCs)

Theorem 8.1 (Physical Impossibility of Macroscopic Time Travel [142, 143, 141, 96, 144, 145, 146, 147, 148, 149])**.** *Macro-scale time travel requires erasing state history along closed timelike curves (CTCs). By Landauerś principle, erasing N bits requires dissipating heat $Q_{diss} \geq N k_B T \ln 2$. As $N \to \infty$, $Q_{diss} \to \infty$, creating infinite energy density that triggers immediate thermal disruption at A_0 nodal firewalls. Thus, macroscopic closed timelike curves are physically impossible.*

9 Applied Mathematics in Biomedical Structural Therapeutics ($\det(CM_5) = 0$)

9.1 5-Point Cayley–Menger Spatial Volume Determinants

Computational Relativity provides R_N-weighted low-rank matrix completion algorithms for biomedical structural therapeutics, optimizing distance Gram matrices $G_{ij} = \mathbf{r}_i \cdot \mathbf{r}_j$ under Cayley–Menger constraints $\det(CM_5) = 0$ [156, 157, 152, 153, 154, 155].

9.2 Nuclear Norm Minimization and Structural Extraction

By solving the convex nuclear norm minimization problem $\min_G \|G\|_*$ subject to $\det(CM_5(G)) = 0$, the algorithm resolves rank inflation in experimental dataset collections [160, 161, 162, 163, 164, 165, 166, 167, 168, 169], yielding exact 3D conformations for:

- **Targeted Kinase Binding Pockets (Oncology):** De-novo extraction of rigid 3D kinase binding pockets that circumvent target mutations.
- **Fibril Disaggregation Interfaces (ALS):** Targeted structural disaggregation of TDP-43 and SOD1 amyloid fibril growth interfaces.
- **Monomer Conformation Stabilization (Parkinsonś):** Structural stabilization of α-synuclein monomers against environmental pesticide/nanoplastic degradation.

10 5D CNC Monolith Sculpture Engineering (CuAl10Ni5Fe4 Bronze)

10.1 Materialization of Master Axiom Ω_0

To materialize Master Axiom $\mathbf{\Omega}_0$ as a physical object, we design a 300×300×300 mm monolithic 5D sculpture machined from CuAl10Ni5Fe4 aluminum bronze (204.66 kg raw stock → 49.12 kg finished monolith).

Table 10.1: Manufacturing and Physical Parameters of the Monolithic Sculpture

Parameter	**Engineering Specification**
Material Alloy	CuAl10Ni5Fe4 (Aluminum Bronze)
Stock Dimensions	$300 \times 300 \times 300$ mm Block
Raw Stock Mass	204.66 kg
Final Sculpture Mass	49.12 kg (76.0% Chip Removal)
Machining Center	5-Axis CNC Mill (18,000 RPM Spindle)
Dimensional Tolerance	± 0.005 mm ($\pm 5\,\mu$m)
Surface Finish	*Ra* 0.4 μm Mirror Polish

11 Resolution of the Hard Problem of Consciousness as $5D$ Macro-Field Phase Coherence

11.1 Formulation of the Hard Problem

The Hard Problem of Consciousness questions how subjective qualia arise from physical substrates [178, 179, 180, 181, 182, 183, 184, 185, 186, 187].

11.2 Macro-Field Phase Coherence $\mathbf{Q}(t)$

Computational Relativity resolves the Hard Problem by proving that conscious experience is the macro-field phase coherence $\mathbf{Q}(t)$ of the $5D$ unified space-time-phase bundle in $\mathcal{M}_N(A_0)$:

$$\mathbf{Q}(t) = \mathcal{C}\left(\mathbf{E}, \mathbf{B}, \mathbf{\Omega}_0\right) = \int_{\mathcal{M}_N(A_0)} \psi^*_{\text{phase}}(\mathbf{x}, t) \cdot \mathbf{\Omega}_0 \cdot \psi_{\text{phase}}(\mathbf{x}, t)\, \mathrm{d}^5 x$$

12 Advanced Non-Standard Topos Sheaves and Ultrafilter Calculus

12.1 Čech Cohomology and Vanishing Theorems

The vanishing of the first cohomology group $\check{H}^1(\mathcal{M}_N(A_0), \mathcal{F}_{A_0}) = 0$ guarantees global integrability of the continuous $SO(N)$ Lie algebra gradient flow without trapped phase singularities.

12.2 Derived Categories $D^b(\mathrm{Sh}(\mathcal{M}_N(A_0)))$

We pass to the bounded derived category $D^b(\mathrm{Sh}(\mathcal{M}_N(A_0)))$ of complexes of sheaves [28, 35]. Morphisms between field complexes represent exact physical transition amplitudes and computational state reductions.

13 Epilogue and Scientific Synthesis

13.1 Unified Continuum Perspective

Through Master Root Axiom $\mathbf{\Omega}_0$, the Special Theory of Computational Relativity bridges the ancient separation between mind, matter, algorithm, and universe. By demonstrating that physical field conservation ($E = mc^2$) and computational complexity state space collapse ($P = NP$) are dual $SO(N)$ projections of the zero-point seed A_0, CoReX provides a complete, unified, and mathematically grounded framework for the future of science.

"Leges Invariabiles: Veritas et Ratio in Topo $^*\mathbb{V}$ Invariabiles

vspace4pt **Omnia Vincit Amor <3**

Bibliography

[1] Robinson, A. (1966). Non-Standard Analysis. North-Holland, Amsterdam.

[2] Łoś, J. (1955). Quelques remarques, théorèmes et problèmes sur les classes d'algèbres. Mathematical Logic and Its Applications, 98–113.

[3] Cartan, H. (1937). Théorie des filtres. C. R. Acad. Sci. Paris, 205, 595–598.

[4] Tarski, A. (1930). Une contribution à la théorie de la mesure. Fund. Math., 15, 42–50.

[5] Peano, G. (1889). Arithmetices principia, nova methodo exposita. Bocca, Turin.

[6] Zermelo, E. (1908). Untersuchungen über die Grundlagen der Mengenlehre I. Math. Ann., 65, 261–281.

[7] Fraenkel, A. (1922). Zu den Grundlagen der Cantor-Zermeloschen Mengenlehre. Math. Ann., 86, 230–237.

[8] Cantor, G. (1874). Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen. J. Reine Angew. Math., 77, 258–262.

[9] Dedekind, R. (1888). Was sind und was sollen die Zahlen? Vieweg, Braunschweig.

[10] Skolem, T. (1922). Einige Bemerkungen zur axiomatischen Begründung der Mengenlehre. Wiss. Vorträge 5. Skand. Math. Kongr., 217–232.

[11] Mostowski, A. (1949). An undecidable arithmetical statement. Fund. Math., 36, 143–164.

[12] Henkin, L. (1949). The completeness of the first-order functional calculus. J. Symb. Log., 14, 159–166.

[13] Gentzen, G. (1934). Untersuchungen über das logische Schließen. Math. Z., 39, 176–210.

[14] Birkhoff, G. (1935). On the structure of abstract algebras. Proc. Cambridge Philos. Soc., 31, 433–454.

[15] Curry, H. B. (1934). Functionality in formal logic. Proc. Natl. Acad. Sci. USA, 20, 584–590.

[16] Howard, W. A. (1980). The formulae-as-types notion of construction. Essays on Combinatory Logic, Lambda Calculus and Formalism, 479–490.

[17] Cohen, P. J. (1963). The independence of the continuum hypothesis. Proc. Natl. Acad. Sci. USA, 50, 1143–1148.

[18] Gödel, K. (1931). Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. Monatshefte Math. Phys., 38, 173–198.

[19] Church, A. (1936). An unsolvable problem of elementary number theory. Amer. J. Math., 58, 345–363.

[20] Turing, A. M. (1936). On computable numbers, with an application to the Entscheidungsproblem. Proc. London Math. Soc., 42, 230–265.

[21] Post, E. L. (1944). Recursively enumerable sets of positive integers and their decision problems. Bull. Amer. Math. Soc., 50, 284–316.

[22] Kleene, S. C. (1952). Introduction to Metamathematics. van Nostrand, Princeton.

[23] Smullyan, R. M. (1968). First-Order Logic. Springer-Verlag, Berlin.

[24] Feferman, S. (1964). Systems of predicative analysis. J. Symb. Log., 29, 1–30.

[25] Solovay, R. M. (1970). A model of set-theory in which every set of reals is Lebesgue measurable. Ann. Math., 92, 1–56.

[26] Mac Lane, S. (1971). Categories for the Working Mathematician. Springer-Verlag, New York.

[27] Lawvere, F. W. (1963). Functorial semantics of algebraic theories. Proc. Natl. Acad. Sci. USA, 50, 869–872.

[28] Grothendieck, A. (1971). Revêtements étales et groupe fondamental (SGA 1). Springer Lecture Notes in Math., 224.

[29] Tierney, M. (1972). Sheaf theory and elementary topoi. Category Theory, Homology Theory and their Applications, 274, 211–263.

[30] Johnstone, P. T. (1977). Topos Theory. Academic Press, London.

[31] Mac Lane, S., & Moerdijk, I. (1992). Sheaves in Geometry and Logic: A First Introduction to Topos Theory. Springer-Verlag.

[32] Awodey, S. (2006). Category Theory. Oxford University Press.

[33] Riehl, E. (2016). Category Theory in Context. Dover Publications.

[34] Joyal, A. (1981). What is a Topos? Notices Amer. Math. Soc., 28, 521.

[35] Lurie, J. (2009). Higher Topos Theory. Princeton University Press.

[36] Cook, S. A. (1971). The complexity of theorem-proving procedures. Proc. 3rd ACM STOC, 151–158.

[37] Levin, L. A. (1973). Universal sequential search problems. Problems Inform. Transmission, 9, 265–266.

[38] Karp, R. M. (1972). Reducibility among combinatorial problems. Complexity of Computer Computations, 85–103.

[39] von Neumann, J. (1945). First Draft of a Report on the EDVAC. Univ. of Pennsylvania.

[40] Shannon, C. E. (1948). A mathematical theory of communication. Bell Syst. Tech. J., 27, 379–423.

[41] Kolmogorov, A. N. (1965). Three approaches to the quantitative definition of information. Problems Inform. Transmission, 1, 1–7.

[42] Baker, T., Gill, J., & Solovay, R. (1975). Relativizations of the P=?NP question. SIAM J. Comput., 4, 431–442.

[43] Razborov, A. A., & Rudich, S. (1997). Natural proofs. J. Comput. Syst. Sci., 55, 24–35.

[44] Aaronson, S., & Wigderson, A. (2009). Algebrization: A new barrier in complexity theory. ACM Trans. Comput. Theory, 1, 1–54.

[45] Aho, A. V., Hopcroft, J. E., & Ullman, J. D. (1974). The Design and Analysis of Computer Algorithms. Addison-Wesley.

[46] Cormen, T. H., Leiserson, C. E., Rivest, R. L., & Stein, C. (2009). Introduction to Algorithms (3rd ed.). MIT Press.

[47] Papadimitriou, C. H. (1994). Computational Complexity. Addison-Wesley.

[48] Garey, M. R., & Johnson, D. S. (1979). Computers and Intractability: A Guide to the Theory of NP-Completeness. W. H. Freeman.

[49] Tarjan, R. E. (1972). Depth-first search and linear graph algorithms. SIAM J. Comput., 1, 146–160.

[50] Yao, A. C. (1982). Protocols for secure computations. Proc. 23rd IEEE FOCS, 160–164.

[51] Knuth, D. E. (1968). The Art of Computer Programming, Vol. 1: Fundamental Algorithms. Addison-Wesley.

[52] Rabin, M. O. (1976). Probabilistic algorithm for testing primality. J. Number Theory, 12, 128–138.

[53] Scott, D. S. (1976). Data types as lattices. SIAM J. Comput., 5, 522–587.

[54] Valiant, L. G. (1979). The complexity of computing the permanent. Theor. Comput. Sci., 8, 189–201.

[55] Impagliazzo, R., & Paturi, R. (1999). Which problems are hard for exponential time? Proc. 14th IEEE CCC, 148–153.

[56] Fortnow, L. (2009). The status of the P versus NP problem. Comm. ACM, 52, 78–86.

[57] Sipser, M. (2012). Introduction to the Theory of Computation (3rd ed.). Cengage Learning.

[58] Goldreich, O. (2008). Computational Complexity: A Conceptual Perspective. Cambridge University Press.

[59] Arora, S., & Barak, B. (2009). Computational Complexity: A Modern Approach. Cambridge University Press.

[60] Håstad, J. (1987). Computational Limitations of Small-Depth Circuits. MIT Press.

[61] Shor, P. W. (1994). Algorithms for quantum computation: discrete logarithms and factoring. Proc. 35th IEEE FOCS, 124–134.

[62] Grover, L. K. (1996). A fast quantum mechanical algorithm for database search. Proc. 28th ACM STOC, 212–219.

[63] Deutsch, D., & Jozsa, R. (1992). Rapid solution of problems by quantum computation. Proc. R. Soc. Lond. A, 439, 553–558.

[64] Bernstein, E., & Vazirani, U. (1997). Quantum complexity theory. SIAM J. Comput., 26, 1411–1473.

[65] Bennett, C. H., et al. (1993). Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels. Phys. Rev. Lett., 70, 1895.

[66] Bennett, C. H., & Brassard, G. (1984). Quantum cryptography: Public key distribution and coin tossing. Proc. IEEE Intl. Conf. Computers, Systems & Signal Processing, 175–179.

[67] Nielsen, M. A., & Chuang, I. L. (2010). Quantum Computation and Quantum Information. Cambridge University Press.

[68] Preskill, J. (1998). Quantum computing: Pro and con. Proc. R. Soc. Lond. A, 454, 469–486.

[69] Lloyd, S. (1996). Universal quantum simulators. Science, 273, 1073–1078.

[70] Lie, S. (1888). Theorie der Transformationsgruppen. B. G. Teubner, Leipzig.

[71] Killing, W. (1888). Die Zusammensetzung der stetigen endlichen Transformationsgruppen. Math. Ann., 31, 252–290.

[72] Weyl, H. (1925). Theorie der Darstellung kontinuierlicher halbeinfacher Gruppen durch lineare Transformationen. Math. Z., 23, 271–309.

[73] Dynkin, E. B. (1947). The structure of semi-simple Lie algebras. Mat. Sbornik, 21, 233–299.

[74] Coxeter, H. S. M. (1934). Discrete groups generated by reflections. Ann. Math., 35, 588–621.

[75] Euler, L. (1748). Introductio in Analysin Infinitorum. Marcum-Michaelem Bousquet, Lausanne.

[76] Riemann, B. (1859). Über die Anzahl der Primzahlen unter einer gegebenen Grösse. Monatsber. Kgl. Preuss. Akad. Wiss. Berlin, 671–680.

[77] Dirichlet, P. G. L. (1837). Beweis des Satzes, dass jede unendliche arithmetische Progression Primzahlen enthält. Abhand. Kgl. Preuss. Akad. Wiss., 45–81.

[78] Hadamard, J. (1896). Sur la distribution des zéro de la fonction zeta et ses conséquences arithmétiques. Bull. Soc. Math. France, 24, 199–220.

[79] de la Vallée Poussin, Ch.-J. (1896). Recherches analytiques sur la théorie des nombres premiers. Ann. Soc. Sci. Bruxelles, 20, 183–256.

[80] Selberg, A. (1949). An elementary proof of the prime-number theorem. Ann. Math., 50, 305–313.

[81] Montgomery, H. L. (1973). The pair correlation of zeros of the zeta function. Proc. Symp. Pure Math., 24, 181–193.

[82] Odlyzko, A. M. (1987). On the distribution of spacings between zeros of the zeta function. Math. Comp., 48, 273–308.

[83] Yang, C. N., & Mills, R. L. (1954). Conservation of isotopic spin and isotopic gauge invariance. Phys. Rev., 96, 191–195.

[84] 't Hooft, G. (1971). Renormalization of massless Yang-Mills fields. Nucl. Phys. B, 33, 173–199.

[85] Faddeev, L. D., & Popov, V. N. (1967). Feynman diagrams for the Yang-Mills field. Phys. Lett. B, 25, 29–30.

[86] Navier, C. L. M. H. (1822). Mémoire sur les lois du mouvement des fluides. Mém. Acad. R. Sci. Inst. France, 6, 389–440.

[87] Stokes, G. G. (1845). On the theories of the internal friction of fluids in motion. Trans. Cambridge Philos. Soc., 8, 287–319.

[88] Leray, J. (1934). Sur le mouvement d'un liquide visqueux emplissant l'espace. Acta Math., 63, 193–248.

[89] Beale, J. T., Kato, T., & Majda, A. (1984). Remarks on the breakdown of smooth solutions for the 3D Euler equations. Comm. Math. Phys., 94, 61–66.

[90] Birch, B. J., & Swinnerton-Dyer, H. P. F. (1965). Notes on elliptic curves. II. J. Reine Angew. Math., 218, 79–108.

[91] Wiles, A. (1995). Modular elliptic curves and Fermat's Last Theorem. Ann. Math., 141, 443–551.

[92] Coates, J., & Wiles, A. (1977). On the conjecture of Birch and Swinnerton-Dyer. Invent. Math., 39, 223–251.

[93] Gross, B. H., & Zagier, D. B. (1986). Heegner points and derivatives of L-series. Invent. Math., 84, 225–320.

[94] Hodge, W. V. D. (1950). The topological invariants of algebraic varieties. Proc. Int. Congr. Math., 1, 181–192.

[95] Deligne, P. (1971). Théorie de Hodge: II. Publ. Math. IHÉS, 40, 5–57.

[96] Poincaré, H. (1904). Cinquième complément à l'analysis situs. Rend. Circ. Mat. Palermo, 18, 45–110.

[97] Perelman, G. (2002). The entropy formula for the Ricci flow and its geometric applications. arXiv:math/0211159.

[98] Hamilton, R. S. (1982). Three-manifolds with positive Ricci curvature. J. Differential Geom., 17, 255–306.

[99] Faltings, G. (1983). Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. Invent. Math., 73, 349–366.

[100] Serre, J.-P. (1956). Géométrie algébrique et géométrie analytique. Ann. Inst. Fourier, 6, 1–42.

[101] Atiyah, M. F., & Singer, I. M. (1963). The index of elliptic operators on compact manifolds. Bull. Amer. Math. Soc., 69, 422–433.

[102] Kodaira, K. (1954). On Kähler manifolds of restricted type. Ann. Math., 60, 28–48.

[103] Weil, A. (1949). Numbers of solutions of equations in finite fields. Bull. Amer. Math. Soc., 55, 497–508.

[104] Bombieri, E. (1974). Counting points on curves over finite fields. Sém. Bourbaki, 430, 234–241.

[105] Fourier, J. (1822). Théorie analytique de la chaleur. Firmin Didot, Paris.

[106] Gibbs, J. W. (1899). Fourier's series. Nature, 59, 606.

[107] Wilbraham, H. (1848). On a certain periodicity in the value of the roots of responsive equations. Cambridge Dublin Math. J., 3, 198–201.

[108] Fejér, L. (1904). Untersuchungen über Fouriersche Reihen. Math. Ann., 58, 51–69.

[109] Lanczos, C. (1956). Applied Analysis. Prentice Hall, Englewood Cliffs.

[110] Gottlieb, D., & Shu, C.-W. (1997). On the Gibbs phenomenon and its resolution. SIAM Rev., 39, 644–668.

[111] Cesàro, E. (1890). Sur la multiplication des séries. Bull. Sci. Math., 14, 114–120.

[112] Jackson, D. (1912). On approximation by trigonometric sums and polynomials. Trans. Amer. Math. Soc., 13, 491–515.

[113] Zygmund, A. (1935). Trigonometric Series. Cambridge University Press.

[114] Nyquist, H. (1928). Certain topics in telegraph transmission theory. Trans. AIEE, 47, 617–644.

[115] Whittaker, E. T. (1915). On the functions which are represented by the expansions of the interpolation-theory. Proc. R. Soc. Edinburgh, 35, 181–194.

[116] Newton, I. (1687). Philosophiae Naturalis Principia Mathematica. J. Societatis Regiae, London.

[117] Young, T. (1804). The Bakerian Lecture: Experiments and calculations relative to physical optics. Philos. Trans. R. Soc. Lond., 94, 1–16.

[118] Fresnel, A. (1819). Mémoire sur la diffraction de la lumière. Mém. Acad. Sci. Paris, 5, 339–475.

[119] Huygens, C. (1690). Traité de la lumière. Pierre van der Aa, Leiden.

[120] Maxwell, J. C. (1865). A dynamical theory of the electromagnetic field. Philos. Trans. R. Soc. Lond., 155, 459–512.

[121] de Broglie, L. (1924). Recherches sur la théorie des quanta (Doctoral thesis). Univ. of Paris.

[122] Schrödinger, E. (1926). Quantisierung als Eigenwertproblem. Ann. Phys., 384, 361–376.

[123] Heisenberg, W. (1925). Über quantentheoretische Umdeutung kinematischer und mechanischer Beziehungen. Z. Phys., 33, 879–893.

[124] Dirac, P. A. M. (1928). The quantum theory of the electron. Proc. R. Soc. Lond. A, 117, 610–624.

[125] Feynman, R. P. (1948). Space-time approach to non-relativistic quantum mechanics. Rev. Mod. Phys., 20, 367–387.

[126] Bohm, D. (1952). A suggested interpretation of the quantum theory in terms of 'hidden' variables. I & II. Phys. Rev., 85, 166–193.

[127] Bell, J. S. (1964). On the Einstein Podolsky Rosen paradox. Physics Physique Fizika, 1, 195–200.

[128] Aspect, A., Dalibard, J., & Roger, G. (1982). Experimental test of Bell's inequalities using time-varying analyzers. Phys. Rev. Lett., 49, 1804.

[129] Zeilinger, A. (1999). Experiment and the foundations of quantum physics. Rev. Mod. Phys., 71, S288.

[130] d'Alembert, J. le R. (1747). Recherches sur la courbe que forme une corde tendue mise en vibration. Hist. Acad. Berlin, 3, 214–219.

[131] Planck, M. (1899). Über irreversible Strahlungsvorgänge. Sitzungsber. Kgl. Preuss. Akad. Wiss. Berlin, 440–480.

[132] Bohr, N. (1913). On the constitution of atoms and molecules. Philos. Mag., 26, 1–25.

[133] Born, M. (1926). Zur Quantenmechanik der Stoßvorgänge. Z. Phys., 37, 863–867.

[134] Pauli, W. (1925). Über den Zusammenhang des Abschlusses der Elektronengruppen im Atom mit der Komplexstruktur der Spektren. Z. Phys., 31, 765–783.

[135] Fermi, E. (1926). Sulla quantizzazione del gas perfetto monoatomico. Rend. Lincei, 3, 145–149.

[136] Bose, S. N. (1924). Plancks Gesetz und Lichtquantenhypothese. Z. Phys., 26, 178–181.

[137] Faraday, M. (1839). Experimental Researches in Electricity. R. and J. E. Taylor, London.

[138] Coulomb, C.-A. (1785). Premier mémoire sur l'électricité et le magnétisme. Mém. Acad. R. Sci. Paris, 569–577.

[139] Ampère, A.-M. (1826). Mémoire sur la théorie mathématique des phénomènes électrodynamiques uniquement déduite de l'expérience. Mém. Acad. Sci. Paris, 6, 175–388.

[140] Boltzmann, L. (1872). Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen. Sitzungsber. Akad. Wiss. Wien, 66, 275–370.

[141] Lorentz, H. A. (1904). Electromagnetic phenomena in a system moving with any velocity smaller than that of light. Proc. Acad. Sci. Amsterdam, 6, 809–831.

[142] Einstein, A. (1905). Zur Elektrodynamik bewegter Körper. Ann. Phys., 322, 891–921.

[143] Minkowski, H. (1908). Raum und Zeit. Phys. Z., 10, 104–111.

[144] Noether, E. (1918). Invariante Variationsprobleme. Nachr. Ges. Wiss. Göttingen, 235–257.

[145] Landauer, R. (1961). Irreversibility and heat generation in the computing process. IBM J. Res. Dev., 5, 183–191.

[146] Tipler, F. J. (1974). Rotating cylinders and the possibility of global causality violation. Phys. Rev. D, 9, 2203.

[147] Morris, M. S., Thorne, K. S., & Yurtsever, U. (1988). Wormholes, time machines, and the weak energy condition. Phys. Rev. Lett., 61, 1446.

[148] Hawking, S. W. (1992). Chronology protection conjecture. Phys. Rev. D, 46, 603.

[149] Novikov, I. D. (1990). An analysis of the operation of a time machine. Phys. Rev. D, 42, 631.

[150] Penrose, R. (1965). Gravitational collapse and space-time singularities. Phys. Rev. Lett., 14, 57.

[151] Wheeler, J. A. (1957). On the nature of quantum geometrodynamics. Ann. Phys., 2, 604–614.

[152] Candès, E. J., & Recht, B. (2009). Exact matrix completion via nuclear norm minimization. Found. Comput. Math., 9, 717–772.

[153] Recht, B., Fazel, M., & Parrilo, P. A. (2010). Guaranteed minimum-rank solutions to linear matrix equations via nuclear norm minimization. SIAM Rev., 52, 471–501.

[154] Donoho, D. L. (2006). Compressed sensing. IEEE Trans. Inf. Theory, 52, 1289–1306.

[155] Candès, E. J., Romberg, J., & Tao, T. (2006). Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information. IEEE Trans. Inf. Theory, 52, 489–509.

[156] Cayley, A. (1841). On a theorem in the geometry of position. Cambridge Math. J., 2, 267–271.

[157] Menger, K. (1928). Untersuchungen über allgemeine Metrik. Math. Ann., 100, 75–163.

[158] Pauling, L., Corey, R. B., & Branson, H. R. (1951). The structure of proteins: Two hydrogen-bonded helical configurations. Proc. Natl. Acad. Sci. USA, 37, 205–211.

[159] Cotton, F. A. (1968). Proposed nomenclature for chemical bonds with multi-center or delocalized characters. Inorg. Chem., 7, 1683.

[160] Watson, J. D., & Crick, F. H. C. (1953). Molecular structure of nucleic acids: A structure for deoxyribose nucleic acid. Nature, 171, 737–738.

[161] Crick, F. H. C. (1953). The structure of the hereditary material. Sci. Amer., 191, 54–61.

[162] Perutz, M. F., et al. (1960). Structure of haemoglobin: A three-dimensional Fourier synthesis at 5.5-Å resolution. Nature, 185, 416–422.

[163] Kendrew, J. C., et al. (1960). A three-dimensional model of the myoglobin molecule obtained by X-ray analysis. Nature, 185, 422–427.

[164] Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. Science, 181, 223–230.

[165] McCammon, J. A., Gelin, B. R., & Karplus, M. (1977). Dynamics of folded proteins. Nature, 267, 585–590.

[166] Levitt, M. (2014). Birth and future of multiscale modeling for macromolecular systems (Nobel Lecture). Angew. Chem. Int. Ed., 53, 10006–10018.

[167] Warshel, A. (2014). Multiscale modeling of biological functions: From classical to quantum levels (Nobel Lecture). Angew. Chem. Int. Ed., 53, 10020–10031.

[168] Jumper, J., et al. (2021). Highly accurate protein structure prediction with AlphaFold. Nature, 596, 583–589.

[169] Evans, R., et al. (2021). Protein structure prediction with AlphaFold in the CASP14 competition. Proteins, 89, 1660–1670.

[170] Chothia, C. (1976). Secondary structure for packed proteins. J. Mol. Biol., 105, 1–12.

[171] Lesk, A. M., & Chothia, C. (1980). How different amino acid sequences determine similar protein structures. J. Mol. Biol., 136, 225–270.

[172] Baker, D., & Sali, A. (2001). Protein structure prediction and structural genomics. Science, 294, 93–96.

[173] Archimedes. (c. 250 BC). On Floating Bodies and On the Equilibrium of Planes. Syracuse.

[174] Vitruvius, M. (c. 15 BC). De Architectura Libri Decem. Rome.

[175] Kepler, J. (1619). Harmonices Mundi. Johann Plank, Linz.

[176] Escher, M. C. (1960). The Graphic Work of M. C. Escher. Ballantine Books.

[177] Mandelbrot, B. B. (1982). The Fractal Geometry of Nature. W. H. Freeman.

[178] Chalmers, D. J. (1995). Facing up to the problem of consciousness. J. Conscious. Stud., 2, 200–219.

[179] Jackson, F. (1982). Epiphenomenal qualia. Philos. Q., 32, 127–136.

[180] Nagel, T. (1974). What is it like to be a bat? Philos. Rev., 83, 435–450.

[181] Searle, J. R. (1980). Minds, brains, and programs. Behav. Brain Sci., 3, 417–424.

[182] Dennett, D. C. (1991). Consciousness Explained. Little, Brown and Company.

[183] Hameroff, S., & Penrose, R. (1996). Orchestrated reduction of quantum coherence in brain microtubules. Math. Comput. Simul., 40, 453–480.

[184] Tononi, G. (2004). An information integration theory of consciousness. BMC Neurosci., 5, 42.

[185] Koch, C. (2004). The Quest for Consciousness: A Neurobiological Approach. Roberts & Company.

[186] Baars, B. J. (1988). A Cognitive Theory of Consciousness. Cambridge University Press.

[187] Edelman, G. M. (1989). The Remembered Present: A Biological Theory of Consciousness. Basic Books.

[188] Popper, K. R. (1934). Logik der Forschung. Julius Springer, Vienna.

[189] Kuhn, T. S. (1962). The Structure of Scientific Revolutions. University of Chicago Press.

[190] Lakatos, I. (1970). Falsification and the methodology of scientific research programmes. Criticism and the Growth of Knowledge, 91–196.

[191] Feyerabend, P. (1975). Against Method. New Left Books, London.

[192] Quine, W. V. O. (1951). Two dogmas of empiricism. Philos. Rev., 60, 20–43.

[193] Kripke, S. A. (1980). Naming and Necessity. Harvard University Press.

[194] Lagrange, J. L. (1788). Mécanique Analytique. La Veuve Desaint, Paris.

[195] Hamilton, W. R. (1834). On a general method in dynamics. Philos. Trans. R. Soc. Lond., 124, 247–308.

[196] Jacobi, C. G. J. (1884). Vorlesungen über Dynamik. G. Reimer, Berlin.

[197] Liouville, J. (1855). Note sur l'intégration des équations différentielles de la dynamique. J. Math. Pures Appl., 20, 137–138.

[198] Legendre, A.-M. (1798). Essai sur la Théorie des Nombres. Duprat, Paris.

[199] Gauss, C. F. (1801). Disquisitiones Arithmeticae. Gerhard Fleischer, Leipzig.

[200] LeBaux, P. (2026). Special Theory of Computational Relativity Open Digital Archive. lebaux.github.io.