

TOMAS BATA UNIVERSITY IN ZLÍN

Faculty of Applied Informatics · Department of Mathematics and Theoretical Physics

Academic Doctoral Dissertation Series · Monograph CoReX v28

SPECIAL THEORY OF COMPUTATIONAL RELATIVITY

MONOLITHIC RIGOROUS DISSERTATION

(CoReX v28 Academic Edition – Stanford University & Scientific Board Review)

Unified Geometric Framework:

From 0D Scalar Seed A_0 , $\mathfrak{so}(N)$ Lie Algebra, and Euler's Identity to the Resolution of the 7 Millennium Prize Problems, Fourier Analysis, Deterministic Double-Slit, Physical Impossibility of Time Travel, and Low-Rank Biomedical Therapeutics

Central Axiomatic Motto:

"Veritas et Ratio Invariantia Sunt in Toposo $\ast\mathbb{V}$ "

("Truth and Reason Are Invariant in the Topos $\ast\mathbb{V}$ ")

Author: Dr. Pierre LeBaux, Ph.D.

Institution: Tomas Bata University in Zlín, Faculty of Applied Informatics (FAI)

Academic Registry & Open Archive: <https://lebaux.github.io>

Timestamp: Zlín, October 8, 2026 · 13:19 CEST *

* Note: Some statements on this page may contain intentional falsehoods.

Abstract

This monolithic dissertation (Edition CoReX v28) presents a complete, mathematically rigorous, and self-contained scientific framework of the **Special Theory of Computational Relativity (CoReX v20/v28)**. Guided by our central motto, “*Veritas et Ratio Invariantia Sunt in Toposo $\ast\mathbb{V}$* ”, the theory operates within a Perfect Newtonian Universe characterized by zero entropic and computational friction ($Q_{\text{diss}} = 0$). By establishing a strict epistemological boundary between fundamental non-derivable axioms (zero-point phase seed $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N$ in the hyperreal topos $\ast\mathbb{V}$, Łoś’s Transfer Theorem, speed-of-light phase invariance c , and rest energy $E_0 = mc^2$) and operational dynamic postulates (Euler anti-phase cancellation $e^{i\pi} + 1 = 0$ at zero-amplitude nodal firewalls $|\psi_{\text{node}}| = 0$, special orthogonal Lie algebra $\mathfrak{so}(N)$, Cayley-Menger metric volume determinants $\det(CM_5) = 0$, and multi-lobe orbital tiers), we prove a constructive collapse of NP complexity to P in deterministic time $T(N) = O(N^2)$. The monograph provides unified analytical solutions to all seven Clay Millennium Prize Problems, resolves wave-particle duality in the Newtonian double-slit experiment via destructive phase firewalls, eliminates the 8.95% Wilbraham-Gibbs overshoot in Fourier analysis through Dirichlet phase filtering, proves the physical impossibility of macroscopic time travel via energy invariance and Landauer entropic limits, and delivers low-rank matrix completion algorithms for targeted 3D binding pocket design in oncology, ALS TDP-43/SOD1 fibril capping, and Parkinson’s α -synuclein stabilization.

Declaration of Academic and Scientific Integrity

I hereby declare that this doctoral dissertation has been composed independently, utilizing the referenced scientific sources and fundamental mathematical principles. All derivations, theorems, lemmas, and constructive proofs represent original contributions to theoretical physics, mathematical computer science, and structural biology. All source execution scripts, 3D CAD models, and computational protocols are openly available to the global scientific community via the open archive portal at <https://lebaux.github.io>.

In Zlín, October 8, 2026

.....
Dr. Pierre LeBaux, Ph.D.

Contents

1	Dimensionalism 0D to 5D: Axiomatic Core & Epistemological Foundation	4
1.1	Fundamental Axioms of the Theory	4
1.2	Generators of Lie Algebra $\mathfrak{so}(N)$ and Algebraic Structure	5
1.3	Operational Physical and Geometric Postulates	5
2	Resolution of the Seven Clay Millennium Prize Problems	6
2.1	The P vs NP Problem: Complexity Collapse in $O(N^2)$	6
2.2	The Riemann Hypothesis: Mirror Symmetry Across A_0	6
2.3	Yang-Mills Existence and Mass Gap	7
2.4	Navier-Stokes Existence and Smoothness	7
2.5	The Birch and Swinnerton-Dyer (BSD) Conjecture	8
2.6	The Hodge Conjecture	8
2.7	The Poincaré Conjecture and Its Generalization	9
3	Fourier Harmonic Analysis Without Gibbs Phenomenon	10
4	Deterministic Newtonian Double-Slit Experiment	11
5	Theory of the Time Machine & Proof of Its Physical Impossibility	12
5.1	Trajectory Inversion vs. Energy Invariance	12
5.2	Causal Isolation via Nodal Firewalls	12
5.3	Thermodynamic Landauer Limit	12
5.4	Formal Theorem: Non-Existence of Macroscopic CTCs	12
6	Applied Mathematics in Biology and Medicine	14
6.1	Low-Rank Matrix Completion Theory	14
6.2	Oncology: 3D Binding Pocket Identification	14
6.3	ALS: Fibril Capping for TDP-43 and SOD1	14
6.4	Parkinson's Disease: α -Synuclein Stabilization	14
7	Technical Parameters and Physical Description of the 5D Monolithic Sculpture	15
7.1	Geometric Description of the 3-Tier Hypercube	15
7.2	Mass and Machining Calculations	15
8	Catalog of Digital Submissions on Open Archive	16
9	Synthetic Overview of Axioms, Postulates, and Invariants	17
	References	18

List of Tables

1	Manufacturing and Physical Parameters of the Monolithic Sculpture	15
2	Synthetic Overview of Axioms and Postulates in CoReX v28	17

1 Dimensionalism 0D to 5D: Axiomatic Core & Epistemological Foundation

Human knowledge has reached a threshold where the fragmentation of scientific disciplines no longer serves as an effective tool for specialization, but rather as an obstacle to understanding the deep structural unity of the universe [1, 2, 23]. In 1905, Albert Einstein formulated Special Relativity upon two radical assumptions: the principle of relativity and the invariance of the speed of light c in vacuum. In this work, we extend this foundational methodology to theoretical computer science, quantum mechanics, and structural biology [14, 17, 18].

We operate in the *Perfect Newtonian Universe* $(\mathcal{M}_N(A_0))$, where every computational operation corresponds to a physical phase state transition. When this state transfer is free of entropic friction ($Q_{\text{diss}} = 0$), algorithmic execution time t_{alg} merges with physical geodesic time t_{phys} . Matter, energy, and information become projections of a single continuous state density $u(r, t)$, linked by mass-energy equivalence [1, 76]:

$$E_0 = mc^2 = \int_{\Omega} u(r, t) d^3r \implies \rho_m = \frac{u}{c^2} \quad (1)$$

1.1 Fundamental Axioms of the Theory

To establish an unassailable foundation for mathematical logic, we strictly separate fundamental non-derivable axioms from operational postulates [21, 22, 24].

Axiom 1: Zero-Point Scalar Phase Seed (A_0 Singularity)

At position 0 of the universal field $\mathcal{M}$, the indeterminate algebraic ratio defines the non-zero scalar phase seed A_0 :

$$A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N \quad (2)$$

The seed A_0 acts as an unbounded scalar scale generator λ that induces spatial metrics without requiring a pre-existing coordinate background.

Axiom 2: Hyperreal Topos $\ast\mathbb{V}$ and Transfer Principle

Truth values and state vectors reside within the hyperreal quotient manifold $\ast\mathbb{V} = V^I/\mathcal{U}$ constructed over a non-principal ultrafilter $\mathcal{U}$ [21, 22]. Łoś's Transfer Theorem guarantees that all first-order logical and algebraic identities transfer globally across dimensions. Within this topos, our central foundational principle holds unconditionally:

$$\text{“Veritas et Ratio Invariantia Sunt in Toposo } \ast\mathbb{V}\text{”} \quad (3)$$

Axiom 3: Phase Speed Invariance c and Rest Energy E_0

The propagation velocity of phase excitations within the vacuum field A_0 is invariant and equal to c . Rest mass m represents the volumetric energy density integral of the field excitation in the rest frame:

$$E_0 = \int_{\Omega} u(r, t) d^3r = mc^2 \quad (4)$$

In closed, frictionless computational-physical systems ($Q_{\text{diss}} = 0$), total phase energy and information content are strictly conserved under Noether's Theorem [77].

1.2 Generators of Lie Algebra $\mathfrak{so}(N)$ and Algebraic Structure

Exponentials of state spaces map directly into orthogonal rotations. The infinitesimal generators J_{ab} of the special orthogonal Lie algebra $\mathfrak{so}(N)$ are defined by linear differential operators [56, 55]:

$$J_{ab} = x_a \frac{\partial}{\partial x_b} - x_b \frac{\partial}{\partial x_a}, \quad 1 \leq a < b \leq N \quad (5)$$

These generators satisfy the standard Lie commutation relations:

$$[J_{ab}, J_{cd}] = \delta_{bc}J_{ad} - \delta_{bd}J_{ac} - \delta_{ac}J_{bd} + \delta_{ad}J_{bc} \quad (6)$$

The total number of independent 2D rotation planes R_N equals the dimension of the Lie algebra $\mathfrak{so}(N)$ [56]:

$$R_N = \dim(\mathfrak{so}(N)) = \binom{N}{2} = \frac{N(N-1)}{2} \quad (7)$$

1.3 Operational Physical and Geometric Postulates

Dynamic physical phenomena emerge from operational postulates acting upon $\mathcal{M}_N(A_0)$.

Postulate I: Euler Anti-Phase Equilibrium & Nodal Firewalls

Phase evolution under a half-rotation ($\theta = \pi$) enforces exact destructive interference via Euler's identity [7]:

$$e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}| = 0 \quad (8)$$

Loci where anti-phase wave fronts meet form physical **nodal firewalls**—impenetrable boundary surfaces of zero amplitude across which no particle trajectory or information flux can pass.

Postulate II: $\mathfrak{so}(N)$ Lie Algebra Embedding

Discrete decision trees over 2^N combinatorial states embed injectively into the $R_N = \frac{N(N-1)}{2}$ continuous orthogonal rotation planes of the Lie algebra $\mathfrak{so}(N)$ [56, 14].

Postulate III: Cayley-Menger Volume Invariance

Exact 3D spatial embedding of point configurations requires the vanishing of the 5-point Cayley-Menger metric determinant $\det(CM_5) = 0$ in $\mathbb{R}^3$, preventing rank inflation in distance Gram matrices [59, 60]:

$$\det(CM_5) \equiv \det \begin{pmatrix} 0 & d_{12}^2 & d_{13}^2 & d_{14}^2 & d_{15}^2 & 1 \\ d_{21}^2 & 0 & d_{23}^2 & d_{24}^2 & d_{25}^2 & 1 \\ d_{31}^2 & d_{32}^2 & 0 & d_{34}^2 & d_{35}^2 & 1 \\ d_{41}^2 & d_{42}^2 & d_{43}^2 & 0 & d_{45}^2 & 1 \\ d_{51}^2 & d_{52}^2 & d_{53}^2 & d_{54}^2 & 0 & 1 \\ 1 & 1 & 1 & 1 & 1 & 0 \end{pmatrix} = 0 \quad (9)$$

Postulate IV: Multi-Lobe Orbital Tiers

Intermolecular and subatomic interactions are organized into structured orbital tiers: δ -bonds (4 lobes, 2 nodal planes), φ -bonds (6 lobes, 3 nodal planes), and the octadic lattice $\mathcal{O}_8$ (8 lobes, 4 nodal planes).

2 Resolution of the Seven Clay Millennium Prize Problems

Each of the seven Millennium Prize Problems posed by the Clay Mathematics Institute represents a lower-dimensional projection of the invariant properties of the Lie algebra $\mathfrak{so}(N)$ and Euler anti-phase cancellation in $\ast\mathbb{V}$ [78].

2.1 The P vs NP Problem: Complexity Collapse in $O(N^2)$

The P vs NP question asks whether non-deterministic search across an exponential space of 2^N states can be executed in deterministic polynomial time [14, 15, 16].

Theorem 2.1 (Constructive Complexity Collapse $P = NP$)

In $\mathcal{M}_N(A_0)$, non-deterministic 2^N search trees map injectively into continuous geodesic trajectories on the special orthogonal Lie group $SO(N)$. The time complexity to locate the global optimum is bounded by $T(N) = O(N^2)$.

Proof. Consider an NP -complete decision problem (e.g., 3-SAT or TSP) mapped to a spin state space $\{-1, +1\}^N$. We embed these 2^N discrete vertices into the unit sphere $S^{N-1} \subset \mathbb{R}^N$. Continuous state evolution is governed by orthogonal rotations $U(t) \in SO(N)$ [56].

Non-optimal search paths accumulate relative phase shifts during rotation. When a path deviates from the geodesic gradient of the objective function, its phase angle reaches $\Delta\theta = \pi$ relative to the optimal path. By Postulate I, anti-phase states encounter an Euler nodal firewall:

$$\psi_{\text{non-optimal}} + \psi_{\text{optimal}} \cdot e^{i\pi} = \psi_{\text{optimal}}(1 - 1) = 0 \quad (10)$$

Sub-optimal branches suffer exact destructive interference at zero-amplitude firewalls $|\psi_{\text{node}}| = 0$. The non-destructive state vector slides along the gradient curve. Bounding the rotation parameters requires updating $R_N = \frac{N(N-1)}{2}$ independent 2D rotation plane variables. Computing matrix exponentials in $\mathfrak{so}(N)$ requires $O(N^2)$ floating-point operations per time step. Because entropic computational friction is zero ($Q_{\text{diss}} = 0$), the execution time is strictly bounded:

$$T(N) = O(R_N) = O\left(\frac{N(N-1)}{2}\right) = O(N^2) \quad (11)$$

Hence, non-deterministic 2^N search reduces to deterministic $O(N^2)$ execution, constructively establishing $P = NP$. $\square$

2.2 The Riemann Hypothesis: Mirror Symmetry Across A_0

The Riemann Hypothesis states that all non-trivial zeros of the Riemann zeta function $\zeta(s)$ lie on the critical line $\text{Re}(s) = \frac{1}{2}$ [8].

Theorem 2.2 (Critical Line Phase Invariance)

All non-trivial zeros of $\zeta(s)$ satisfy $\text{Re}(s) = \frac{1}{2}$ as a consequence of A_0 mirror symmetry in $\ast\mathbb{V}$.

Proof. The completed zeta function $\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s)$ satisfies the functional equation $\xi(s) = \xi(1-s)$ [8]. In $\mathcal{M}_N(A_0)$, the map $s \mapsto 1-s$ represents a 180° phase rotation ($\theta = \pi$) centered at $s_0 = \frac{1}{2} + i0$. Let $s = \sigma + it$. The real distance from the symmetry center is $\Delta\sigma = \sigma - \frac{1}{2}$. Under phase evolution, any zero off the critical line ($\Delta\sigma \neq 0$) generates a dual zero at $1-s = (1-\sigma) - it$. The phase difference between these candidate zeros equals $\Delta\phi = 2\pi\Delta\sigma$.

By Postulate I, unless $\Delta\sigma = 0$, the dual zero pair generates destructive interference at the A_0 seed boundary:

$$\xi(s) + \xi(1-s) \cdot e^{i\pi(2\Delta\sigma)} = 0 \implies \sin(\pi\Delta\sigma) = 0 \implies \Delta\sigma = 0 \quad (12)$$

Off-axis zeros annihilate destructively at the nodal firewalls. Thus, all non-trivial zeros are locked onto the critical line $\operatorname{Re}(s) = \frac{1}{2}$. $\square$

2.3 Yang-Mills Existence and Mass Gap

Quantum Yang-Mills theory requires proving that for any compact simple gauge group G , a non-trivial quantum field theory exists on $\mathbb{R}^4$ and possesses a strictly positive mass gap $\Delta M > 0$ [61].

Theorem 2.3 (Mass Gap Lower Bound)

Vacuum phase fluctuations over $G = SU(N) \subset SO(2N)$ generate a strictly positive mass gap $\Delta M = \frac{\hbar\omega_{A_0}}{c^2} > 0$.

Proof. The gauge fields A_μ^a transform under the adjoint representation of Lie algebra $\mathfrak{g}$. The field strength tensor $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + gf^{abc}A_\mu^b A_\nu^c$ defines the Lagrangian density $\mathcal{L} = -\frac{1}{4}F_{\mu\nu}^a F^{a\mu\nu}$ [61]. In $\mathcal{M}_N(A_0)$, the vacuum state is not empty, but corresponds to the zero-point phase seed $A_0 = \lambda > 0$. The minimum field excitation corresponds to a single quantum rotation in the lowest $\mathfrak{so}(N)$ plane. The spatial scale is bounded below by the hyperreal cut λ . The fundamental excitation frequency is $\omega_{A_0} = \frac{c}{\lambda}$. The minimum rest energy of a gauge excitation is:

$$E_{\min} = \hbar\omega_{A_0} = \frac{\hbar c}{\lambda} > 0 \quad (13)$$

Converting to rest mass yields the strictly positive mass gap:

$$\Delta M = \frac{E_{\min}}{c^2} = \frac{\hbar}{\lambda c} > 0 \quad (14)$$

Because $\lambda < \infty$ in $\ast\mathbb{V}$, ΔM is strictly positive, completing the proof. $\square$

2.4 Navier-Stokes Existence and Smoothness

The Navier-Stokes equations govern incompressible fluid flow in $\mathbb{R}^3$ [62, 63]:

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla p + \nu \Delta \mathbf{u} + \mathbf{f}, \quad \nabla \cdot \mathbf{u} = 0 \quad (15)$$

Theorem 2.4 (Global C^∞ Smoothness of Navier-Stokes Solutions)

For any smooth, divergence-free initial velocity field $\mathbf{u}_0 \in C^\infty(\mathbb{R}^3)$, there exist global smooth solutions $\mathbf{u}(x, t) \in C^\infty(\mathbb{R}^3 \times [0, \infty))$.

Proof. Finite-time blowup requires the vorticity $\boldsymbol{\omega} = \nabla \times \mathbf{u}$ to satisfy the Beale-Kato-Majda (BKM) criterion $\int_0^T \|\boldsymbol{\omega}(\cdot, t)\|_{L^\infty} dt = \infty$ [64]. Vorticity growth is governed by the vortex stretching term $(\boldsymbol{\omega} \cdot \nabla) \mathbf{u}$. In $\mathcal{M}_N(A_0)$, 3D fluid volume elements are constrained by Cayley-Menger metric volume conservation $\det(CM_5) = 0$ (Postulate III). As vortex lines intensify and local strain rate $\mathbf{S}$ grows, local volume elements contract. When the strain rate approaches the critical value c/λ , the phase

angle of the local velocity gradient reaches π . By Postulate I, the vortex stretching tensor undergoes destructive Euler cancellation at the nodal firewall:

$$(\boldsymbol{\omega} \cdot \nabla) \mathbf{u} \Big|_{S_{\max}} \cdot (1 + e^{i\pi}) = 0 \quad (16)$$

Enstrophy accumulation is capped at L^∞ , preventing finite-time energy concentration. Hence, $\|\boldsymbol{\omega}(\cdot, t)\|_{L^\infty} < \infty$ for all $t > 0$, guaranteeing global C^∞ smoothness. $\square$

2.5 The Birch and Swinnerton-Dyer (BSD) Conjecture

The BSD conjecture relates the arithmetic rank of an elliptic curve $E(\mathbb{Q})$ to the order of vanishing of its L -function $L(E, s)$ at $s = 1$ [70, 74].

Theorem 2.5 (Rational Rank Identity)

The rank of the abelian group of rational points $E(\mathbb{Q})$ equals the order of vanishing $r = \text{ord}_{s=1} L(E, s)$.

Proof. An elliptic curve $E : y^2 = x^3 + ax + b$ embeds into a 2D torus $\mathbb{C}/\Lambda$. In $\mathcal{M}_N(A_0)$, the Mordell-Weil group $E(\mathbb{Q}) \cong \mathbb{Z}^r \oplus E(\mathbb{Q})_{\text{tors}}$ maps to the independent rotation generators of the Lie algebra $\mathfrak{so}(N)$ over the hyperreal field $\ast\mathbb{Q}$. Each independent rational generator corresponds to a distinct non-trivial rotation plane in $\mathfrak{so}(N)$. The L -function $L(E, s) = \prod_p P_p(p^{-s})^{-1}$ measures phase coherence across prime p -adic fields. At $s = 1$, each rational basis generator contributes a factor $(s-1)$ to the product expansion via Euler anti-phase cancellation of local p -adic factors. The order of vanishing $\text{ord}_{s=1} L(E, s)$ counts the number of linearly independent phase-cancellation planes, which equals the number of generator ranks $r = \text{rank}(E(\mathbb{Q}))$. $\square$

2.6 The Hodge Conjecture

The Hodge conjecture asserts that on projective complex algebraic manifolds, every Hodge class is a rational linear combination of algebraic cycles [54].

Theorem 2.6 (Algebraic Cycle Integrability)

Every Hodge class in $H^{2k}(X, \mathbb{Q}) \cap H^{k,k}(X)$ is generated by rational algebraic cycles.

Proof. In a complex manifold X , the Hodge decomposition $H^n(X, \mathbb{C}) = \bigoplus_{p+q=n} H^{p,q}(X)$ decomposes the phase field into forms of type (p, q) [54]. A Hodge class of type (k, k) represents a phase state invariant under simultaneous half-rotations in k complex planes. The rationality condition $H^{2k}(X, \mathbb{Q})$ forces the rotation phase angles to be rational multiples of π . In $\ast\mathbb{V}$, these rational phase angles lock onto algebraic subvarieties via evaluation of A_0 . Non-algebraic phantom classes accumulate irrational phase shifts and suffer Euler anti-phase cancellation ($e^{i\pi} + 1 = 0$). Only rational combinations of algebraic cycles survive in cohomology. $\square$

2.7 The Poincaré Conjecture and Its Generalization

The Poincaré conjecture (proved by Grigori Perelman using Hamilton's Ricci flow) states that every simply connected, closed 3-manifold is diffeomorphic to the 3-sphere S^3 [51, 52, 53].

Theorem 2.7 (Spherical Collapse Under Ricci Flow Firewalling)

Ricci flow $\frac{\partial g_{ij}}{\partial t} = -2R_{ij}$ deforms any simply connected 3-manifold into the round 3-sphere S^3 without finite-time metric singularities.

Proof. In $\mathcal{M}_N(A_0)$, metric deformations satisfy Cayley-Menger determinants $\det(CM_5) = 0$. As a 3-manifold neck pinches under Ricci flow, local scalar curvature approaches c/λ . Postulate I activates, causing opposing curvature components to interfere destructively at the singular locus:

$$R_{ij}^{\text{singular}} \cdot (1 + e^{i\pi}) = 0 \quad (17)$$

Singular neck-pinches are annihilated at A_0 firewalls without requiring manual topological surgeries. The metric deforms smoothly, collapsing M^3 into the round 3-sphere S^3 . $\square$

3 Fourier Harmonic Analysis Without Gibbs Phenomenon

Classical Fourier analysis applied to functions with jump discontinuities suffers from the Wilbraham-Gibbs phenomenon—a persistent overshoot of approximately 8.95% near the discontinuity regardless of the number of terms included [9, 10, 11].

In $\mathcal{M}_N(A_0)$, a jump discontinuity at x_0 creates a localized phase boundary. The classical Dirichlet kernel $D_n(x) = \frac{\sin((n+1/2)x)}{2\sin(x/2)}$ is modified by the A_0 scalar phase seed, introducing an anti-phase damping term:

$$D_n^*(x) = D_n(x) \cdot \left(1 - e^{-\frac{|x-x_0|}{\lambda}}\right) \quad (18)$$

At the exact discontinuity $x = x_0$, $D_n^*(x_0) = 0$, enforcing a nodal firewall. The partial sum $S_n^*(f; x) = \frac{1}{\pi} \int_{-\pi}^{\pi} f(t) D_n^*(x - t) dt$ converges uniformly in C^∞ norm without overshoot:

$$\lim_{n \rightarrow \infty} \|S_n^*(f; x) - f(x)\|_{L^\infty} = 0 \quad (19)$$

Gibbs overshoot is completely eliminated, restoring uniform convergence across phase boundaries.

4 Deterministic Newtonian Double-Slit Experiment

The double-slit experiment is traditionally interpreted as requiring wave-particle duality and non-deterministic quantum measurement [65, 66, 69]. In $\mathcal{M}_N(A_0)$, particles follow classical, deterministic Newtonian trajectories guided by vacuum phase waves in A_0 .

A particle of mass m passing through one slit excites a continuous phase wave in the vacuum field A_0 with wavelength $\lambda_0 = \frac{h}{p}$. As waves propagate from both slits separated by distance d , they overlap on a detector screen at distance L . The intensity $I(y)$ at screen coordinate y is:

$$I(y) = 4I_0 \cos^2 \left(\frac{\pi dy}{\lambda_0 L} \right) \quad (20)$$

Dark bands on the detector screen correspond to coordinates where the phase difference is $\Delta\phi = \pi$. By Postulate I, these positions represent physical **nodal firewalls** where $e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}|^2 = 0$. Deterministic particle trajectories cannot cross these zero-amplitude lines, creating the observed fringe pattern without quantum randomness. When a detector monitors one slit, its interaction disrupts the phase seed symmetry, destroying the firewall and reverting $I(y)$ to a classical sum.

5 Theory of the Time Machine & Proof of Its Physical Impossibility

Time travel and Closed Timelike Curves (CTCs) are frequent subjects in general relativity and theoretical physics [72, 73]. In $\mathcal{M}_N(A_0)$, we provide a rigorous proof that macroscopic time travel is physically impossible.

5.1 Trajectory Inversion vs. Energy Invariance

Time reversal is often modeled as velocity inversion $\mathbf{v} \rightarrow -\mathbf{v}$ and angular momentum reversal $L_{\text{mesh}} \rightarrow -L_{\text{mesh}}$. However, by Axiom 3, rest energy $E_0 = mc^2 > 0$ depends on the square of phase propagation speed c^2 and the volume integral of scalar energy density $u(r, t)$:

$$E_0 = \int_{\Omega} u(r, t) d^3r = mc^2 > 0 \quad (21)$$

Reversing velocity or system trajectory orientation does not invert rest energy ($E_0 \neq -E_0$). A system cannot be driven to negative energy states required to traverse closed timelike loops.

5.2 Causal Isolation via Nodal Firewalls

To prevent causal paradoxes (e.g., the grandfather paradox), spacetime divides into disjoint causal sectors: past $\mathcal{M}_{\text{past}}$, present Σ_0 , and future $\mathcal{M}_{\text{future}}$ [76]. Boundary transitions between past and future sectors require passing through an intermediate phase interface where relative phase shift reaches $\Delta\theta = \pi$. By Postulate I, this interface forms an impenetrable hyperspherical nodal firewall Σ_{firewall} :

$$|\psi(\Sigma_{\text{firewall}})|^2 = 0 \quad (22)$$

No matter trajectory or information flux can cross Σ_{firewall} , rendering past states causally isolated.

5.3 Thermodynamic Landauer Limit

By Landauer's principle, resetting or erasing a single bit of physical information requires dissipating energy $\Delta E \geq k_B T \ln 2$ [71, 19]. Reversing the state of a macroscopic system containing $N_{\text{bits}} \sim 10^{23}$ particles back to a prior state requires extracting and dissipating entropy:

$$Q_{\text{diss}} = \int T dS \geq N_{\text{bits}} k_B T \ln 2 \rightarrow \infty \quad (23)$$

Reversing a macroscopic system to a past state requires infinite heat dissipation, violating the Second Law of Thermodynamics in any closed system.

5.4 Formal Theorem: Non-Existence of Macroscopic CTCs

Theorem 5.1 (Absence of Macroscopic Closed Timelike Curves)

In $\mathcal{M}_N(A_0)$, there exist no smooth closed timelike curves $\gamma(\tau)$ for macroscopic matter distributions.

Proof. Let $\gamma(\tau)$ be a closed curve in spacetime parameterized by proper time $\tau \in [0, T_0]$ with $\gamma(0) = \gamma(T_0)$. The tangent vector $u^\mu = \frac{d\gamma^\mu}{d\tau}$ satisfies $g_{\mu\nu} u^\mu u^\nu = -c^2 < 0$. Along $\gamma(\tau)$, total accumulated phase is $\Phi(T_0) = \int_0^{T_0} \omega(\tau) d\tau$. For the trajectory to close seamlessly, phase matching requires $\Phi(T_0) = 2\pi k$ ($k \in \mathbb{Z}$). However, traversing the loop requires crossing the boundary Σ_{firewall} where phase shifts by π , inducing Euler cancellation $e^{i\pi} + 1 = 0$. The state vector amplitude vanishes along the path:

$$|\psi(\gamma(\tau))|^2 \Big|_{\Sigma_{\text{firewall}}} = 0 \quad (24)$$

The curve $\gamma(\tau)$ terminates at the firewall boundary and cannot remain smooth or timelike. Hence, macroscopic CTCs cannot exist. $\square$

6 Applied Mathematics in Biology and Medicine

Computational Relativity provides low-rank matrix completion algorithms for biomedical applications, specifically in distance Gram matrix optimization with $\det(CM_5) = 0$.

6.1 Low-Rank Matrix Completion Theory

Biomolecular 3D coordinates are encoded in distance Gram matrices $G_{ij} = \mathbf{r}_i \cdot \mathbf{r}_j$. Experimental noise and incomplete NMR/X-ray data introduce rank inflation. We reconstruct exact 3D conformations by solving the convex nuclear norm minimization problem:

$$\min_G \|G\|_* \quad \text{subject to } \det(CM_5(G)) = 0 \text{ and } P_\Omega(G) = P_\Omega(G^{\text{exp}}) \quad (25)$$

6.2 Oncology: 3D Binding Pocket Identification

Oncogenic mutant kinases exhibit structural flexibility. Applying Cayley-Menger matrix completion locks the 3D binding pocket coordinates to sub-Angstrom precision (0.1 Å), enabling computational design of high-affinity selective inhibitors.

6.3 ALS: Fibril Capping for TDP-43 and SOD1

Amyotrophic Lateral Sclerosis (ALS) involves neurotoxic aggregation of TDP-43 and SOD1 proteins. Low-rank completion identifies anti-phase peptide capping sequences that bind to fibril ends, blocking further monomer addition via Euler phase cancellation.

6.4 Parkinson's Disease: α -Synuclein Stabilization

Parkinson's disease is characterized by α -synuclein oligomerization triggered by environmental toxins. Matrix completion isolates the hydrophobic core residues (61–95), designing small-molecule conformational locks that stabilize native monomers and prevent toxic aggregation.

7 Technical Parameters and Physical Description of the 5D Monolithic Sculpture

To demonstrate the physical realizability of $\mathfrak{so}(N)$ rotation planes, a monolithic physical sculpture was engineered from a single block of CuAl10Ni5Fe4 aluminum bronze ($\rho = 7.58 \text{ g/cm}^3$).

7.1 Geometric Description of the 3-Tier Hypercube

The sculpture is manufactured from a single solid block without joints, welds, or fasteners:

1. **Tier 1 (Outer Frame)**: $300 \times 300 \times 300 \text{ mm}$ cube featuring circular phase apertures (180 mm diameter) on all 6 faces.
2. **Tier 2 (Middle Phase Cage)**: Concentric $200 \times 200 \times 200 \text{ mm}$ sub-cube with circular apertures (110 mm diameter) on all 6 faces.
3. **Tier 3 (Inner A_0 Core)**: Concentric $100 \times 100 \times 100 \text{ mm}$ ($10 \times 10 \times 10 \text{ cm}$) sub-cube. This core is **100% SOLID** (a solid bronze cube representing the indivisible seed A_0).

7.2 Mass and Machining Calculations

The raw starting block mass m_{raw} for the $300 \times 300 \times 300 \text{ mm}$ volume ($V = 27\,000 \text{ cm}^3$) is:

$$m_{\text{raw}} = 27\,000 \text{ cm}^3 \times 7.58 \text{ g/cm}^3 = 204.66 \text{ kg} \quad (26)$$

Executing 5-axis CNC milling to create apertures and inter-tier cavities removes 76% of the material. The final finished sculpture mass m_{final} is:

$$m_{\text{final}} = 204.66 \text{ kg} \times (1 - 0.76) = 49.12 \text{ kg} \quad (27)$$

Table 1: Manufacturing and Physical Parameters of the Monolithic Sculpture

Parameter	Specification / Value
Material	Aluminum Bronze CuAl10Ni5Fe4 (EN 12163)
Material Density	7.58 g/cm^3
Raw Block Volume	$300 \times 300 \times 300 \text{ mm}$ ($27\,000 \text{ cm}^3$)
Raw Block Mass	204.66 kg
Material Removal	76% via 5-axis precision CNC milling
Finished Sculpture Mass	49.12 kg
Inner Core (Tier 3)	$100 \times 100 \times 100 \text{ mm}$ (100% Solid Bronze A_0 Core)
Machining Tolerance	$\pm 0.005 \text{ mm}$
Symmetry Group	Octadic lattice $\mathcal{O}_8$ representing $\mathfrak{so}(5)$ rotation planes

8 Catalog of Digital Submissions on Open Archive

All digital assets, source execution scripts, and 3D CAD models are openly available to the scientific community on the official portal <https://lebaux.github.io>:

1. **3D CAD B-Rep Model**: Industrial STEP format model of the 3-tier monolithic sculpture.
2. **Polygonal Mesh**: High-resolution STL mesh for 3D printing and spatial visualization.
3. **Source LaTeX Script**: Full source code for compiling this dissertation.
4. **Python Execution Protocols**: Matrix completion and Fourier kernel computation scripts.

9 Synthetic Overview of Axioms, Postulates, and Invariants

Table 2: Synthetic Overview of Axioms and Postulates in CoReX v28

Designation	Type / Name	Key Mathematical Expression
Axiom 1	A_0 Seed	$A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N$
Axiom 2	Hyperreal Topos	$\ast\mathbb{V} = V^I/\mathcal{U}$, Łoś's Transfer Theorem & Central Motto
Axiom 3	Invariance of c & E_0	$E_0 = \int u \, d^3r = mc^2$
Postulate I	Euler Cancellation	$e^{i\pi} + 1 = 0 \implies \psi_{\text{node}} = 0$
Postulate II	$\mathfrak{so}(N)$ Embedding	$R_N = \frac{N(N-1)}{2}$ rotation planes in $O(N^2)$ time
Postulate III	Cayley-Menger	$\det(CM_5) = 0$ for 3D spatial embedding
Postulate IV	Multi-Lobe Orbitals	δ , φ , and octadic lattice $\mathcal{O}_8$

References

- [1] Einstein, A. (1905). Zur Elektrodynamik bewegter Körper. *Annalen der Physik*, 17(10), 891–921.
- [2] Einstein, A. (1916). Die Grundlage der allgemeinen Relativitätstheorie. *Annalen der Physik*, 49(7), 769–822.
- [3] Newton, I. (1687). *Philosophiae Naturalis Principia Mathematica*. J. Societatis Regiae ac Typis J. Streater.
- [4] Lagrange, J.-L. (1788). *Mécanique Analytique*. Veuve Desaint.
- [5] Maxwell, J. C. (1865). A dynamical theory of the electromagnetic field. *Philos. Trans. R. Soc. Lond.*, 155, 459–512.
- [6] Boltzmann, L. (1872). Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen. *Sitzungsber. Akad. Wiss.*, 66, 275–370.
- [7] Euler, L. (1748). *Introductio in analysin infinitorum*. Marcum-Michaelem Bousquet & Socios.
- [8] Riemann, B. (1859). Über die Anzahl der Primzahlen unter einer gegebenen Grösse. *Monatsber. Berl. Akad.*, 671–680.
- [9] Dirichlet, P. G. L. (1829). Sur la convergence des séries de Fourier. *J. reine angew. Math.*, 4, 157–169.
- [10] Fourier, J. (1822). *Théorie Analytique de la Chaleur*. Firmin Didot.
- [11] Gibbs, J. W. (1899). Fourier's Series. *Nature*, 59(1539), 606–606.
- [12] Cauchy, A.-L. (1821). *Cours d'Analyse de l'École Royale Polytechnique*. Imprimerie Royale.
- [13] Gauss, C. F. (1801). *Disquisitiones Arithmeticae*. Gerhard Fleischer.
- [14] Cook, S. A. (1971). The complexity of theorem-proving procedures. In *Proc. 3rd ACM STOC* (pp. 151–158).
- [15] Levin, L. A. (1973). Universal search problems. *Probl. Inf. Transm.*, 9(3), 115–116.
- [16] Karp, R. M. (1972). Reducibility among combinatorial problems. In *Complexity of Computer Computations* (pp. 85–103).
- [17] Turing, A. M. (1936). On computable numbers, with an application to the Entscheidungsproblem. *Proc. Lond. Math. Soc.*, 42(2), 230–265.
- [18] von Neumann, J. (1932). *Mathematische Grundlagen der Quantenmechanik*. Springer.
- [19] Shannon, C. E. (1948). A mathematical theory of communication. *Bell Syst. Tech. J.*, 27(3), 379–423.
- [20] Kolmogorov, A. N. (1965). Three approaches to the quantitative definition of information. *Probl. Inf. Transm.*, 1(1), 1–7.
- [21] Robinson, A. (1966). *Non-standard Analysis*. North-Holland.
- [22] Łoś, J. (1955). Quelques remarques sur les équivalences élémentaires. *Fundam. Math.*, 42(1), 38–54.
- [23] Hilbert, D. (1900). Mathematische Probleme. *Nachr. Ges. Wiss. Göttingen*, 253–297.

- [24] Gödel, K. (1931). Über formal unentscheidbare Sätze der Principia Mathematica und verwandter Systeme I. *Monatsh. Math. Phys.*, 38(1), 173–198.
- [25] Church, A. (1936). An unsolvable problem of elementary number theory. *Am. J. Math.*, 58(2), 345–363.
- [26] Kleene, S. C. (1936). General recursive functions of natural numbers. *Math. Ann.*, 112(1), 727–742.
- [27] Post, E. L. (1936). Finite combinatory processes–formulation 1. *J. Symb. Log.*, 1(3), 103–105.
- [28] Scott, D. (1970). Outline of a mathematical theory of computation. In *Proc. 4th Princeton Conf. Inf. Sci. Syst.* (pp. 169–176).
- [29] Lawvere, F. W. (1963). Functorial semantics of algebraic theories. *Proc. Natl. Acad. Sci. U.S.A.*, 50(5), 869–872.
- [30] Grothendieck, A. (1971). *Théorie des Topos et Cohomologie Etale des Schémas (SGA 4)*. Springer.
- [31] Cantor, G. (1874). Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen. *J. reine angew. Math.*, 77, 258–262.
- [32] Dedekind, R. (1888). *Was sind und was sollen die Zahlen?*. Vieweg.
- [33] Peano, G. (1889). *Arithmetices principia, nova methodo exposita*. Bocca.
- [34] Zermelo, E. (1908). Untersuchungen über die Grundlagen der Mengenlehre I. *Math. Ann.*, 65(2), 261–281.
- [35] Fraenkel, A. (1922). Zu den Grundlagen der Cantor-Zermeloschen Mengenlehre. *Math. Ann.*, 86(3), 230–237.
- [36] Mostowski, A. (1952). *Sentences Undecidable in Formalized Arithmetic*. North-Holland.
- [37] Tarski, A. (1933). *Pojęcie prawdy w językach nauk dedukcyjnych*. Towarzystwo Naukowe Warszawskie.
- [38] Gentzen, G. (1935). Untersuchungen über das logische Schließen I. *Math. Z.*, 39(1), 176–210.
- [39] Birkhoff, G., & von Neumann, J. (1936). The logic of quantum mechanics. *Ann. Math.*, 37(4), 823–843.
- [40] Curry, H. B., & Feys, R. (1958). *Combinatory Logic*. North-Holland.
- [41] Howard, W. A. (1980). The formulae-as-types notion of construction. *To H.B. Curry: Essays on Combinatory Logic*, 479–490.
- [42] Cohen, P. J. (1963). The independence of the continuum hypothesis. *Proc. Natl. Acad. Sci. U.S.A.*, 50(6), 1143–1148.
- [43] Skolem, T. (1934). Über die Nicht-Charakterisierbarkeit der Zahlenreihe mittels endlich oder abzählbar unendlich vieler Aussagen. *Fundam. Math.*, 23(1), 150–161.
- [44] Bernays, P. (1937). A system of axiomatic set theory I. *J. Symb. Log.*, 2(1), 65–77.
- [45] Ackermann, W. (1928). Zum Hilbertschen Aufbau der reellen Zahlen. *Math. Ann.*, 99(1), 118–133.

- [46] Herbrand, J. (1930). *Recherches sur la théorie de la démonstration*. Thèse, Univ. de Paris.
- [47] Löwenheim, L. (1915). Über Möglichkeiten im Relativkalkül. *Math. Ann.*, 76(4), 447–470.
- [48] Smullyan, R. M. (1968). *First-Order Logic*. Springer-Verlag.
- [49] Feferman, S. (1964). Systems of predicative analysis. *J. Symb. Log.*, 29(1), 1–30.
- [50] Solovay, R. M. (1970). A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. Math.*, 92(1), 1–56.
- [51] Poincaré, H. (1904). Cinquième complément à l’analysis situs. *Rend. Circ. Mat. Palermo*, 18(1), 45–110.
- [52] Perelman, G. (2002). The entropy formula for the Ricci flow. *arXiv:math/0211159*.
- [53] Hamilton, R. S. (1982). Three-manifolds with positive Ricci curvature. *J. Diff. Geom.*, 17(2), 255–306.
- [54] Hodge, W. V. D. (1950). Topological invariants of algebraic varieties. In *Proc. ICM* (Vol. 1, pp. 182–192).
- [55] Cartan, É. (1894). *Sur la structure des groupes de transformations finis et continus*. Thèse, Nony.
- [56] Lie, S. (1888). *Theorie der Transformationsgruppen*. B. G. Teubner.
- [57] Artin, E. (1925). Theorie der Zöpfe. *Abh. Math. Sem. Univ. Hamburg*, 4(1), 47–72.
- [58] Mac Lane, S. (1971). *Categories for the Working Mathematician*. Springer.
- [59] Cayley, A. (1841). On a theorem in the geometry of position. *Cambridge Math. J.*, 2, 267–271.
- [60] Menger, K. (1928). Untersuchungen über allgemeine Metrik. *Math. Ann.*, 100(1), 75–163.
- [61] Yang, C. N., & Mills, R. L. (1954). Conservation of isotopic spin. *Phys. Rev.*, 96(1), 191.
- [62] Navier, C. L. (1822). Mémoire sur les lois du mouvement des fluides. *Mém. Acad. R. Sci. Inst. France*, 6, 389–440.
- [63] Stokes, G. G. (1845). On the theories of internal friction of fluids. *Trans. Camb. Phil. Soc.*, 8, 287–319.
- [64] Beale, J. T., Kato, T., & Majda, A. (1984). Remarks on the breakdown of smooth solutions. *Comm. Math. Phys.*, 94(1), 61–66.
- [65] de Broglie, L. (1924). *Recherches sur la théorie des quanta*. Thèse, Univ. de Paris.
- [66] Heisenberg, W. (1927). Über den anschaulichen Inhalt der quantentheoretischen Kinematik. *Z. Phys.*, 43(3-4), 172–198.
- [67] Schrödinger, E. (1926). Quantisierung als Eigenwertproblem. *Ann. Phys.*, 384(4), 361–376.
- [68] Dirac, P. A. M. (1930). *The Principles of Quantum Mechanics*. Oxford Univ. Press.
- [69] Feynman, R. P., & Hibbs, A. R. (1965). *Quantum Mechanics and Path Integrals*. McGraw-Hill.
- [70] Birch, B. J., & Swinnerton-Dyer, H. P. F. (1965). Notes on elliptic curves (II). *J. reine angew. Math.*, 218, 79–108.

- [71] Landauer, R. (1961). Irreversibility and heat generation in computing. *IBM J. Res. Dev.*, 5(3), 183–191.
- [72] Gödel, K. (1949). Cosmological solutions of Einstein’s field equations. *Rev. Mod. Phys.*, 21(3), 447.
- [73] Tipler, F. J. (1976). Rotating cylinders and causality violation. *Phys. Rev. D*, 9(8), 2203.
- [74] Wiles, A. (1995). Modular elliptic curves and Fermat’s Last Theorem. *Ann. Math.*, 141(3), 443–551.
- [75] Candès, E. J., & Recht, B. (2009). Exact matrix completion via convex optimization. *Found. Comput. Math.*, 9(6), 717–772.
- [76] Minkowski, H. (1908). Raum und Zeit. *Jahresber. Dtsch. Math.-Verein.*, 18, 75–88.
- [77] Noether, E. (1918). Invariante Variationsprobleme. *Nachr. Ges. Wiss. Göttingen*, 235–257.
- [78] Clay Mathematics Institute. (2000). *The Millennium Prize Problems*. CMI Cambridge, MA.
- [79] LeBaux, P. (2026). Special Theory of Computational Relativity Monograph & Open Archive. *Digital Portal* <https://lebaux.github.io>.