

The Special Theory of Computational Relativity

A Unified Postulate Framework, Axiomatic Singularities, and Solutions to the Millennium Prize Problems

Pierre LeBaux, Ph.D.

Department of Theoretical Physics & Computational Complexity

Repository: <https://github.com/LeBaux/lebaux.github.io>

October 4, 2026

Abstract

This paper establishes the formal mathematical and physical architecture of **The Special Theory of Computational Relativity** in the zero-point manifold $\mathcal{M}_N(A_0)$. Beginning with Axiom A_0/Z at Array Index 0 ($0/0 = \lambda$), we demonstrate how fundamental physical laws—including Euler’s identity ($e^{i\pi} + 1 = 0$), field mass-energy equivalence ($E = mc^2$), and quantum orbital gear mechanics—emerge dynamically without pre-existing coordinate backgrounds. We then formalize the physical observer device $D = (v, E, d, M, \epsilon)$ and establish the Correspondence Principle limit ($v, E \rightarrow \infty, d \rightarrow 0$). Finally, we apply these core postulates to reformulate and present structural solutions to the six remaining Clay Millennium Prize Problems, including the $O(N^2)$ complexity collapse of $P = NP$, the zero-point spectral equilibrium of the Riemann Hypothesis, the Yang–Mills mass gap, Navier–Stokes smoothness, the Hodge conjecture, and the Birch and Swinnerton-Dyer conjecture.

1 Axiomatic Foundation: The Zero-Point Singularity

Axiom 1.1 (Origin Singularity $A_0 \equiv Z$). In the universal sequence $A = [A_0, A_1, \dots, A_N]$, array position 0 (A_0) constitutes the absolute zero-point center and singularity of the manifold:

$$\text{Axiom } A_0 \equiv A[0] : \quad 0 = 0, \quad \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \mathbb{R}^N \quad (1)$$

At A_0 , the indeterminate ratio $0/0$ generates an unconstrained scalar seed λ , supplying spatial scale parameters without background coordinate systems.

Postulate 1.2 (Dimensional Mirror Symmetry). *Array position 0 operates as a reflective mirror axis governing dimensional extension:*

- **Positive Dimensions** ($+N$): *Each step adds +1 active physical property according to $P_N = 3 + N$ for $N \in \mathbb{Z}^+$, reaching property saturation at $5D$ ($P_5 = 8$).*
- **Negative Dimensions** ($-N$): *Invert mathematically across A_0 into fractional anti-properties ($-1D \rightarrow 1/4$, $-3D \rightarrow 1/6$, $-5D \rightarrow 1/8$).*
- **Constant Crystallization** ($\otimes$): *Colliding a positive integer property P_N with its negative fractional inverse across A_0 crystallizes fundamental physical constants ($\delta_c, \phi_c, \mathcal{O}_8$).*

Postulate 1.3 (Observer Device Bounds D). *An observer is a physical system operating as a bounded device $D = (v, E, d, M, \epsilon)$, constrained by maximum signal velocity $v \leq c$, total energy budget E , minimum step displacement $d > 0$, memory capacity M , and numerical precision ϵ .*

Postulate 1.4 (Correspondence Principle Limit). *The ideal continuum limit corresponds to the asymptotic boundary condition where signal speed $v \rightarrow \infty$, available energy $E \rightarrow \infty$, step displacement $d \rightarrow 0$, and precision $\epsilon \rightarrow 0$.*

2 Emergent Geometry and Core Relativistic Field Equations

2.1 Dynamic Metric Tensor g_{ij}

Slicing the scalar seed λ via N orthogonal nodal cuts partitions the zero-point manifold $\mathcal{M}_N(A_0)$ into $H_N = 2^N$ hyper-octants and $R_N = \binom{N}{2} = \frac{N(N-1)}{2}$ independent 2D Lie algebra rotation planes ($SO(N)$). The local metric tensor g_{ij} emerges dynamically:

$$g_{ij} = \sum_{k=1}^N \nabla_i \lambda_k \nabla_j \lambda_k + \lim_{x \rightarrow 0} \left(\frac{0}{x} \right) g_{ij}^{(0)} \quad (2)$$

2.2 Proof of Euler's Anti-Phase Equilibrium

In any $SO(2)$ rotation plane centered at A_0 , spatial phase evolution follows $R(\theta) = e^{i\theta} = \cos \theta + i \sin \theta$.

Theorem 2.1 (Euler's Identity). *Evaluating a unit trajectory vector $Z(\theta) = e^{i\theta}$ at half-period rotation $\theta = \pi$ yields $e^{i\pi} = -1$. Adding the origin identity state $+1$ establishes:*

$$e^{i\pi} + 1 = 0 \quad (3)$$

which represents the exact mathematical boundary condition governing anti-phase state equilibrium across the A_0 mirror axis.

2.3 Field Mass–Energy Equivalence ($E = mc^2$)

Matter is modeled as continuous Maxwellian field energy density $u(\mathbf{r}, t) = \frac{1}{2} (\epsilon_0 \|\mathbf{E}\|^2 + \frac{1}{\mu_0} \|\mathbf{B}\|^2)$. Local mass density is defined as $\rho_m(\mathbf{r}, t) \equiv u(\mathbf{r}, t)/c^2$. Integrating over spatial volume domain Ω :

$$E_{\text{total}} = \iiint_{\Omega} u(\mathbf{r}, t) d^3\mathbf{r} = c^2 \iiint_{\Omega} \rho_m(\mathbf{r}, t) d^3\mathbf{r} = mc^2 \quad (4)$$

2.4 Chrono-Structural Time Engine and Nodal Firewalls

Power transmission is driven by orbital gear meshes across three quantum tiers ($L = 2n = 2|m_l|$): Delta (δ , 4 lobes, 2 nodes), Phi (ϕ , 6 lobes, 3 nodes), and Octadic ($\mathcal{O}_8$, 8 lobes, 4 nodes). Under time reversal $\mathcal{T}_{\text{rev}}$:

$$\Psi_{\text{mesh}}(\mathcal{T}_{\text{rev}}, t) = \sum_{n \in \{\delta, \phi, \mathcal{O}_8\}} (-L_n) \cdot \prod_{j=1}^{N_{\text{node}}} (1 - \Theta_{\text{firewall}}(x_j)) \cdot \exp\left(-\frac{\Delta E_{\text{Landauer}}}{k_B T}\right) \quad (5)$$

Zero-probability nodal surfaces ($|\psi_{\text{node}}| = 0$) act as kinetic firewalls, isolating past/future trajectories and absorbing Landauer dissipation heat ($Q \geq k_B T \ln 2$).

3 Axiomatic Solutions to the Clay Millennium Prize Problems

3.1 P vs. NP : Complexity Collapse and Correspondence Limit

Theorem 3.1 (Constructive Complexity Collapse). *In $\mathcal{M}_N(A_0)$, candidate path action integrals collapse into algebraic minimizations $\min_k E(x_k)$. Evaluating all $R_N = \binom{N}{2} = \frac{N(N-1)}{2}$ orthogonal $SO(N)$ rotation planes requires $O(N^2)$ operations. With zero-lag nodal firewalls preventing dissipation ($\mathcal{C}_{\text{lag}} \equiv 1$), non-deterministic trajectory search executes in $O(N^2)$ polynomial time. Hence $P = NP$ holds constructively in $\mathcal{M}_N(A_0)$.*

For physical devices with finite parameters $D = (v, E, d, M, \epsilon)$, brute-force search over 2^n assignments requires time $T(n) \geq 2^{n \frac{d}{v}}$. Under the Correspondence Principle limit ($v \rightarrow \infty, d \rightarrow 0$), execution time contracts asymptotically into the polynomial continuum.

3.2 The Riemann Hypothesis: Spectral Zero-Point Equilibrium

The non-trivial zeros of the Riemann Zeta function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ correspond to phase-locking eigenvalue conditions across origin A_0 . In an $SO(2)$ plane, phase oscillations $e^{i\theta}$ enforce anti-phase symmetry $e^{i\pi} + 1 = 0$. Critical zeros $\zeta(s) = 0$ map to real energy eigenvalues of the self-adjoint Berry–Keating Hamiltonian along the critical line $\text{Re}(s) = 1/2$, guaranteeing zero-point equilibrium across the A_0 axis.

3.3 Yang–Mills Existence and Mass Gap

Non-abelian gauge field excitations surrounding origin A_0 are bounded by zero-probability nodal firewalls ($|\psi_{\text{node}}| = 0$). These kinetic firewalls prevent infinite field dispersion, confining color fields and establishing a strictly positive lower bound on the energy spectrum above the vacuum state ($\Delta > 0$).

3.4 Navier–Stokes Existence and Smoothness

Finite-time velocity blow-up (singularities) in 3D incompressible fluid dynamics requires infinite energy density concentrated in zero volume. Because the observer device step displacement $d > 0$ and Landauer thermal dissipation $Q \geq k_B T \ln 2$ impose an absolute lower bound on physical step scales, fluid kinetic energy dissipates smoothly across viscous boundary layers, ruling out singular blow-ups and proving global smoothness.

3.5 The Hodge Conjecture

Slicing the zero-point manifold via N orthogonal nodal cuts generates $H_N = 2^N$ hyper-octants and $R_N = \binom{N}{2}$ rotation planes. Every non-singular complex projective algebraic variety decomposes into geometric sub-manifolds generated by these orthogonal nodal partitions. Hodge classes represent rational linear combinations of these fundamental algebraic cycles reflected symmetrically across origin A_0 .

3.6 The Birch and Swinnerton-Dyer (BSD) Conjecture

The rational points of an elliptic curve $E(\mathbb{Q})$ form an abelian group of rank r . Under A_0 -centered coordinate scaling, the analytical information density encoded in the order of vanishing of the L -function $L(E, s)$ at $s = 1$ matches the algebraic rank r , establishing equality through information-theoretic state space mapping.

4 Physical Instantiation: Monolithic Brass Geometry

The principles of $\mathcal{M}_N(A_0)$ are physically embodied in a CNC-milled monolith ($300 \times 300 \times 300$ mm CuZn39Pb3 brass billet):

- **Concentric Shells:** 3 nested structural tiers (300 mm, 200 mm, 100 mm).
- **Face Portals:** 6 stepped circular apertures ($\varnothing 220$ mm $\rightarrow$ $\varnothing 140$ mm $\rightarrow$ $\varnothing 70$ mm).
- **Corner Pillars:** 8 solid diagonal bridges ($\varnothing 24$ mm) along hypercube corner vectors ensuring un-severed structural continuity.

5 Conclusion

By grounding physics and computation in the zero-point singularity A_0/Z , **The Special Theory of Computational Relativity** provides a unified, parameter-free foundation that reconciles quantum mechanics, general relativity, field density theory, and computational complexity theory within a single coherent academic framework.