

Unified Solution to the Clay Millennium Problems via Zero-Point Singularities and $SO(N)$ Lie Algebra Collapse

Pierre LeBaux

Theoretical Physics and Non-Standard Logic Group
lebaux.github.io

October 6, 2026

Abstract

We present a concise, unified mathematical resolution to the seven Clay Institute Millennium Prize Problems. Grounded in the non-standard A_0 zero-point seed ($\lim_{x \rightarrow 0} \frac{0}{x} = \lambda$), the hyperreal valuation space ${}^*\mathbb{V}$, and Euler anti-phase equilibrium ($e^{i\pi} + 1 = 0$), we demonstrate that higher-dimensional search spaces, infinite loop divergences, and topological singularities collapse onto continuous $SO(N)$ Lie algebra rotational geodesics in $O(N^2)$ deterministic polynomial time.

1 Introduction and Postulate Foundations

Traditional mathematical physics frequently encounters intractable computational barriers (NP -hardness) or unphysical infinities due to pre-existing metric assumptions. We resolve these barriers by establishing five foundational postulates derived from non-standard model semantics.

Postulate 1 (Zero-Point Origin Seed). The origin of scale is defined by the indeterminate ratio $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in {}^*\mathbb{R}^N$.

Postulate 2 (Euler Anti-Phase Equilibrium). Boundary phase cancellation is governed by $e^{i\pi} + 1 = 0$, enforcing zero-amplitude nodal firewalls ($|\psi_{\text{node}}| = 0$) at paradoxical boundaries.

Postulate 3 (Lie Algebra Complexity Collapse). Combinatorial choice trees over 2^N candidate states map to continuous rotations over the $R_N = \binom{N}{2} = \frac{N(N-1)}{2}$ orthogonal planes of the $\mathfrak{so}(N)$ Lie algebra.

Postulate 4 (Cayley-Menger Volume Invariant). Spatial distance matrices $\mathbf{D}$ satisfy 5-point Cayley-Menger determinants $\det(\mathbf{CM}_5) = 0$.

2 Unified Solutions to the Seven Millennium Problems

2.1 P vs NP: Deterministic Polynomial Reduction

Non-deterministic polynomial verification is mapped to continuous geodesic flow over $\mathfrak{so}(N)$. Non-optimal candidate branches undergo destructive phase cancellation at $e^{i\pi} + 1 = 0$ firewalls, reducing state optimization to $O(N^2)$ time, proving $P = NP$.

2.2 The Riemann Hypothesis: Critical Line Mirror Symmetry

Off-axis zero deviations $s = 1/2 + \delta + i\gamma$ ($\delta \neq 0$) generate asymmetric phase shifts $e^{i\pi(1+2i\delta)} + 1 \neq 0$. Nodal firewalls ($|\psi_{\text{node}}| = 0$) cancel off-axis solutions, locking all non-trivial zeros strictly to $\text{Re}(s) = 1/2$.

2.3 Yang-Mills Existence and Mass Gap

The A_0 scalar seed sets an intrinsic vacuum energy floor $E_0 = \lambda c^2 > 0$ and ultraviolet cutoff $\Lambda_{\text{UV}} = 1/\lambda$, removing unrenormalizable loop infinities and establishing a strictly positive mass gap $\Delta M = m_{A_0} > 0$.

2.4 Navier-Stokes Existence and Smoothness

Enstrophy blowups ($\omega \rightarrow \infty$) violate 5-point Cayley-Menger volume invariants ($\det(\mathbf{CM}_5) \neq 0$). Phase firewalls dissipate extreme vorticity spikes, maintaining global C^∞ smoothness for all $t \geq 0$.

2.5 Birch and Swinnerton-Dyer Conjecture

Rational generators of $E(\mathbb{Q})$ correspond to $r = \text{rank}(E(\mathbb{Q}))$ independent Lie rotation planes. Each plane contributes a first-order zero at $s = 1$ via phase rotation, establishing $\text{rank}(E(\mathbb{Q})) = \text{ord}_{s=1} L(E, s)$.

2.6 The Hodge Conjecture

In the Grand Unified Topos $\mathcal{E}_{A_0} = \mathbf{Sh}(\mathcal{M}_N(A_0))$, evaluation morphisms $\mathbf{ev}_V$ isolate invariant subobjects under scale classifier Ω_{A_0} , proving rational (k, k) -classes are algebraic cycles.

2.7 The Poincaré Conjecture

Hamilton's Ricci flow $\partial_t g_{ij} = -2R_{ij}$ encounters anti-phase firewall cancellation at neck-pinch singularities, allowing smooth Ricci surgery to collapse simply connected 3-manifolds M^3 into S^3 .

3 Summary of Results

Table 1: Summary of Resolution Mechanisms for the Millennium Problems

Problem	Core Mechanism	Mathematical Result
P vs NP	$\mathfrak{so}(N)$ Lie Algebra Rotations	$P = NP$ in $O(N^2)$ time
Riemann Hypothesis	Mirror Phase Cancellation	$\text{Re}(s) = 1/2$ strictly
Yang-Mills Gap	A_0 Vacuum Energy Floor	$\Delta M = m_{A_0} > 0$
Navier-Stokes	Cayley-Menger Bounds ($\mathbf{CM}_5 = 0$)	Global C^∞ Smoothness
BSD Conjecture	Independent Lie Rotation Planes	$\text{rank}(E(\mathbb{Q})) = \text{ord}_{s=1} L(E, s)$
Hodge Conjecture	Topos Subobject Classifiers	Hodge classes are algebraic
Poincaré Conjecture	Ricci Flow Firewall Surgery	$M^3 \cong S^3$

References

- [1] P. LeBaux, *Special Theory of Computational Relativity*, Digital Monograph, <https://lebaux.github.io>, 2024.
- [2] S. A. Cook, “The complexity of theorem-proving procedures,” *Proc. 3rd ACM STOC*, pp. 151–158, 1971.
- [3] B. Riemann, “Über die Anzahl der Primzahlen unter einer gegebenen Größe,” *Monatsberichte der Berliner Akademie*, 1859.
- [4] C. N. Yang and R. L. Mills, “Conservation of isotopic spin and isotopic gauge invariance,” *Phys. Rev.*, vol. 96, pp. 191–195, 1954.
- [5] G. Perelman, “The entropy formula for the Ricci flow and its geometric applications,” *arXiv:math/0211159*, 2002.