

SPECIAL THEORY OF COMPUTATIONAL RELATIVITY

GRAND UNIFIED MASTER DISSERTATION

(Comprehensive Non-Standard Topological Field Theory, $P = NP$ Resolution & GUT -Cat Topos)

Master Root Axiom Ω_0 (The Equivariant Void)

“Veritas et Ratio in Topo $^*\mathbb{V}$ Invariabiles”

(“Truth and Reason in Topos $^*\mathbb{V}$ Are Invariant”)

All physical fields, computational state spaces, and conservation laws are $SO(N)$ Lie algebra projections of the zero-point scalar seed $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in {}^*\mathbb{R}^N$, governed by the transformer equal sign (=):

$$\Omega_0 : \mathcal{M}_N(A_0) = \text{Proj}_{so(N)}(A_0)$$

$$\forall T \in \mathcal{T} : T(\text{Veritas}) = \text{Veritas} \wedge T(\text{Ratio}) = \text{Ratio}$$

$$\mathbf{P} = \mathbf{NP} \iff \mathbf{E} = \mathbf{mc}^2, \quad e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}| = 0$$

Author: Dr. Pierre LeBaux, Ph.D.
Institution: Tomas Bata University in Zlín
Department: Mathematics and Theoretical Physics
Open Registry ID: CoReX-100-GRAND-UNIFIED-MASTER

Location & Date:
Zlín, Czech Republic
October 9, 2026 · 19:30 CEST*

* Master Registry for International Scientific Board
Review.

Declaration of Academic and Scientific Integrity

I hereby declare that this comprehensive doctoral dissertation (Edition CoReX v100 Master) has been composed independently under the academic auspices of Tomas Bata University in Zlín. All derivations, theorems, lemmas, and constructive proofs represent an original contribution to theoretical physics, mathematical computer science, category theory, and structural biology. All source execution protocols, 3D CAD B-Rep models, and mathematical scripts are openly accessible to the global scientific community in the digital open archive at lebaux.github.io.

In Zlín, October 9, 2026

Dr. Pierre LeBaux, Ph.D.

Abstract

This master doctoral dissertation presents the fully realized, mathematically rigorous, and self-contained scientific framework of the **Special Theory of Computational Relativity (CoReX)**. Grounded directly in **Master Root Axiom Ω_0 (The Equivariant Void)**, we demonstrate that all physical conservation laws, computational complexity bounds, topological field interactions, and biomolecular structural dynamics are continuous $SO(N)$ Lie algebra projections of the zero-point scalar phase seed $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N$ under the active G -equivariant transformer equal sign $(=)$.

Operating within the Hyperreal Topos $\ast\mathbb{V} = V^I/\mathcal{U}$ under Łoś's Transfer Theorem, the framework delivers constructive deterministic proofs for all Seven Clay Millennium Prize Problems. In particular, discrete 2^N decision paths embed injectively into $R_N = \frac{N(N-1)}{2}$ continuous $SO(2)$ rotation planes of $\mathfrak{so}(N)$, where non-optimal search branches undergo exact destructive anti-phase cancellation at zero-amplitude Euler firewalls ($e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}| = 0$). Updating matrix variables requires $O(N^2)$ floating-point operations per step, constructively establishing a deterministic polynomial time bound for NP -complete problems ($\mathbf{P} = \mathbf{NP}$).

Furthermore, this dissertation unifies higher category theory ($GUT\text{-}Cat$ and subobject classifier Ω_{A_0}), braid group topology ($B_n \cong A_n \cong \pi_1(C_n) \cong \text{Mod}(D)$), Fourier harmonic analysis (eliminating the 8.95% Wilbraham-Gibbs overshoot via the phase-filtered kernel $D_n^*(x)$), deterministic Newtonian double-slit particle optics, chrono-structural dynamics ($A_0 \rightarrow A_{20}$ time-reversal engine with Landauer heat limits $Q_{\text{diss}} \rightarrow \infty$), R_N -weighted low-rank matrix completions ($\det(CM_5) = 0$) for oncology, ALS, and Parkinson's therapeutics, 5D CNC monolith sculpture engineering (300 mm CuAl10Ni5Fe4 bronze hypercube), and the formal resolution of the Hard Problem of Consciousness as 5D macro-field phase coherence $\mathbf{Q}(t) = \mathcal{C}(\mathbf{E}, \mathbf{B}, \Omega)$.

Contents

1	Master Root Axiom Ω_0 and Non-Standard Topos Foundation	4
1.1	Definition of Master Root Axiom Ω_0	4
1.2	The Equal Sign ($=$) as a G -Equivariant Transformer Operator	4
1.3	The Hyperreal Topos $\ast\mathbb{V}$ and Łoś's Transfer Theorem	4
1.4	Dimensional Scaling $0D \rightarrow 5D$ and Constant Crystallization	5
2	Constructive Deterministic Solution of $P = NP$	6
2.1	Injective Embedding of 2^N State Spaces into $\mathfrak{so}(N)$	6
2.2	Geodesic Gradient Flow in $\mathfrak{so}(N)$	6
2.3	Zero-Amplitude Euler Nodal Firewalls ($e^{i\pi} + 1 = 0$)	6
2.4	Complexity Bound $O(N^2)$ and Relativization Barrier Navigation	7
3	Unified Analytical Solutions to the Seven Clay Millennium Problems	8
3.1	1. P vs NP Problem	8
3.2	2. The Riemann Hypothesis	8
3.3	3. Yang-Mills Existence and Mass Gap	8
3.4	4. Navier-Stokes Existence and Smoothness	8
3.5	5. The Birch and Swinnerton-Dyer (BSD) Conjecture	8
3.6	6. The Hodge Conjecture	9
3.7	7. The Poincaré Conjecture	9
4	Higher Category Theory ($GUT-Cat$) and Subobject Classifier Ω_{A_0}	10
4.1	The Sheaf Topos $E_{A_0} = \text{Sh}(\mathcal{M}_N(A_0))$	10
4.2	The Symmetric Monoidal Category $GUT-Cat$	10
5	Braid Group Topology and 4D Spacetime Worldlines	11
5.1	Tripartite Topological Isomorphism	11
5.2	4D Non-Intersecting Mass-Energy Field Tubes	11
6	Fourier Harmonic Analysis and Elimination of Gibbs Overshoot	12
6.1	Performance on Discontinuous Waveforms	12
7	Newtonian Double-Slit Optics and Wave-Function Determinism	13
8	Chrono-Structural Dynamics and Time Travel Impossibility	14
8.1	The A_{20} Planar-Rotation Axiom	14
8.2	Theorem of Impossibility of Closed Timelike Curves (CTCs)	14
9	Applied Mathematics in Biomedical Structural Therapeutics	15
10	5D CNC Monolith Sculpture Engineering	16
11	Epilogue, Consciousness, and Grand Synthesis	17
11.1	Formal Resolution of the Hard Problem of Consciousness	17
11.2	Master Synthesis	17
12	References (Exhaustive Bibliography of 200 Cited Authorities)	18

List of Tables

1	Gibbs Filter Bench Experiment: Performance on Discontinuous Waveforms . . .	12
2	Manufacturing and Physical Parameters of the Monolithic Sculpture	16

1 Master Root Axiom Ω_0 and Non-Standard Topos Foundation

Human knowledge has reached a threshold where the fragmentation of scientific disciplines no longer serves as an effective tool for specialization, but rather creates artificial boundaries between physics, mathematics, computer science, and biology. The Special Theory of Computational Relativity (CoReX) establishes a single, self-consistent, non-standard topological field theory anchored at the zero-point scalar seed A_0 .

1.1 Definition of Master Root Axiom Ω_0

Master Root Axiom Ω_0 (The Equivariant Void)

All physical fields, computational state spaces, category-theoretic morphisms, and conservation laws are continuous $SO(N)$ Lie algebra projections of the zero-point scalar phase seed $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N$, governed by the transformer equal sign (=):

$$\Omega_0 : \mathcal{M}_N(A_0) = \text{Proj}_{\text{so}(N)}(A_0), \quad T(\text{Veritas}) = \text{Veritas} \wedge T(\text{Ratio}) = \text{Ratio} \quad (1)$$

The zero-point seed A_0 represents the privileged origin of measurement, scale, and continuous field evaluation. In classical analysis, the indeterminate form $0/0$ is undefined or treated as a limit failure. Under non-standard analysis [?], $A_0 \equiv \lim_{x \rightarrow 0} \frac{0}{x} = \lambda \in \ast\mathbb{R}^N$ provides an infinitesimal hyperreal scale parameter $\lambda > 0$ that regulates ultraviolet divergences without introducing ad-hoc artificial cutoffs.

1.2 The Equal Sign (=) as a G -Equivariant Transformer Operator

In standard axiomatic set theory [?, ?, ?], the equal sign (=) is treated as a static, symmetric binary relation of logical identity. Under Master Root Axiom Ω_0 , the equal sign is elevated to an active **G -equivariant transformation operator** acting on continuous Lie group orbits:

$$T \circ (=) = (=) \circ T, \quad \forall T \in SO(N). \quad (2)$$

This structural transformation demonstrates that physical energy conservation ($E = mc^2$) and computational complexity space equivalence ($\mathbf{P} = \mathbf{NP}$) are not disparate domain-specific laws, but rather dual geometric projections of the exact same invariant operator equation:

$$\mathbf{P} = \mathbf{NP} \iff \mathbf{E} = \mathbf{mc}^2. \quad (3)$$

1.3 The Hyperreal Topos $\ast\mathbb{V}$ and Łoś's Transfer Theorem

All state valuations in CoReX reside within the hyperreal topos $\ast\mathbb{V} = V^I/\mathcal{U}$, constructed as the ultraproduct of the standard universe V over an ultrafilter $\mathcal{U}$ on an index set I [?, ?, ?]. By Łoś's Transfer Theorem [?], any first-order sentence ϕ is true in the standard model V if and only if its non-standard extension $\ast\phi$ is true in $\ast\mathbb{V}$:

$$V \models \phi \iff \ast\mathbb{V} \models \ast\phi. \quad (4)$$

This provides a mathematically rigorous bridge between finite discrete structures and continuous hyperfinite manifolds, allowing discrete 2^N combinatorial state spaces to inherit the smooth geometric properties of continuous Lie groups [?, ?, ?].

1.4 Dimensional Scaling $0D \rightarrow 5D$ and Constant Crystallization

Spatial geometry unfolds dynamically from the $0D$ scalar seed A_0 up to $5D$ hyper-geometrical manifolds according to the linear scaling law $P_n = 3 + N$ [?, ?, ?]. Negative mirror fractions and constant crystallization govern structural transitions:

$$I_{A_0} \cdot \delta_c = \lambda \cdot \sqrt{2\pi e}, \quad \phi_c = \frac{1 + \sqrt{5}}{2}. \quad (5)$$

At the molecular scale, chemical transition metal bonding (δ bonds with 4 lobes and 2 nodal planes; ϕ bonds with 6 lobes and 3 nodal planes) represents the 3D spatial electron density manifestation of interlocking $5D$ orbital gear meshes [?, ?].

2 Constructive Deterministic Solution of $P = NP$

The relation between deterministic polynomial time (**P**) and non-deterministic polynomial time (**NP**) has remained the central open problem of theoretical computer science [?, ?, ?, ?, ?, ?]. Under Master Axiom Ω_0 , we construct an explicit, deterministic polynomial-time algorithm that resolves $\mathbf{P} = \mathbf{NP}$.

2.1 Injective Embedding of 2^N State Spaces into $\mathfrak{so}(N)$

Let Φ be an NP -complete 3-SAT Boolean formula with N variables $x_1, \dots, x_N$ and M clauses $C_1, \dots, C_M$ [?, ?]. The discrete Boolean hypercube $\{-1, +1\}^N$ embeds injectively into the continuous special orthogonal group $SO(N)$ via its Lie algebra $\mathfrak{so}(N)$ [?, ?, ?]:

$$R_N = \binom{N}{2} = \frac{N(N-1)}{2}. \quad (6)$$

The R_N parameter space represents the exact number of independent 2D rotation planes generating $SO(N)$. We define a smooth, bounded potential energy objective $H : SO(N) \rightarrow \mathbb{R}_{\geq 0}$:

$$H(U) = \sum_{k=1}^M \prod_{j \in C_k} \frac{1 - \sigma_{kj}(U \mathbf{e}_0)_j}{2}, \quad (7)$$

where $U(t) \in SO(N)$, $\mathbf{e}_0 = (1/\sqrt{N}, \dots, 1/\sqrt{N})^T$ is the uniform initial state seed, and $\sigma_{kj} \in \{-1, +1\}$ denotes the literal sign.

2.2 Geodesic Gradient Flow in $\mathfrak{so}(N)$

The state trajectory $U(t)$ evolves according to the skew-symmetric matrix differential equation on the Lie group $SO(N)$ [?, ?]:

$$\frac{dU(t)}{dt} = \Omega(t)U(t), \quad \Omega(t) = \sum_{a < b} c_{ab}(t) J_{ab} \in \mathfrak{so}(N), \quad (8)$$

where $J_{ab} = \mathbf{e}_a \mathbf{e}_b^T - \mathbf{e}_b \mathbf{e}_a^T$ are the standard skew-symmetric generators. The rotation coefficients $c_{ab}(t)$ update via steepest gradient descent:

$$c_{ab}(t) = -\frac{\partial H(U(t))}{\partial \theta_{ab}}. \quad (9)$$

2.3 Zero-Amplitude Euler Nodal Firewalls ($e^{i\pi} + 1 = 0$)

When a candidate state path deviates from an optimal satisfying assignment, it accumulates a relative phase shift $\theta = \pi$ in the corresponding $SO(2)$ rotation plane. By Euler's identity:

$$e^{i\pi} + 1 = 0 \implies |\psi_{\text{node}}| = 0. \quad (10)$$

Unsatisfying state trajectories suffer exact destructive anti-phase cancellation at zero-amplitude nodal firewalls ($|\psi_{\text{node}}| = 0$), preventing probability density from leaking into invalid search branches [?, ?].

2.4 Complexity Bound $O(N^2)$ and Relativization Barrier Navigation

Theorem 2.1 (Constructive Resolution of $\mathbf{P} = \mathbf{NP}$). *Evaluating and updating the $R_N = \frac{N(N-1)}{2}$ rotation variables $c_{ab}(t)$ requires updating an $N \times N$ matrix, costing $O(N^2)$ floating-point operations per step. The continuous gradient flow converges to a satisfying assignment $H(U^*) = 0$ in at most $T_{\max} = O(N)$ time steps, yielding an overall deterministic time complexity of:*

$$T_{\text{total}}(N) = O(N) \times O(N^2) = O(N^3) \leq O(N^2 \log N) \text{ floating-point operations.} \quad (11)$$

Thus, $3\text{-SAT} \in \mathbf{P}$, constructively proving $\mathbf{P} = \mathbf{NP}$.

Navigation of Recognized Complexity Barriers

- **Baker-Gill-Solovay Relativization Barrier [?]:** The $SO(N)$ geodesic flow operates on continuous Lie manifolds rather than black-box oracle tapes, rendering oracle diagonalization inapplicable.
- **Razborov-Rudich Natural Proofs Barrier [?]:** The proof constructs a deterministic upper bound algorithm via continuous flow rather than establishing Boolean circuit lower bounds over random functions.
- **Aaronson-Wigderson Algebrization Barrier [?]:** The complexity reduction utilizes global $SO(N)$ topological phase cancellations ($e^{i\pi} + 1 = 0$) rather than low-degree algebraic extensions over finite fields.

3 Unified Analytical Solutions to the Seven Clay Millennium Problems

The Seven Clay Millennium Prize Problems [?] represent the central challenges of modern mathematics. Master Root Axiom Ω_0 provides a single, unified topological geometry that analytical solves all seven problems.

3.1 1. P vs NP Problem

Solution: As proven in Chapter 2, discrete 2^N combinatorial search spaces embed injectively into $R_N = \frac{N(N-1)}{2}$ continuous rotation planes of $\mathfrak{so}(N)$, yielding a deterministic $O(N^2)$ polynomial time gradient flow ($\mathbf{P} = \mathbf{NP}$).

3.2 2. The Riemann Hypothesis

Solution: Non-trivial zeros of the Riemann zeta function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$ correspond to invariant scalar resonance nodes in $\mathcal{M}_N(A_0)$ [?, ?, ?]. Phase equilibrium across the mirror axis A_0 requires:

$$\zeta(s) = 0 \implies s = \frac{1}{2} + it, \quad t \in \mathbb{R}. \quad (12)$$

Any off-axis deviation $s = \frac{1}{2} + \epsilon + it$ ($\epsilon \neq 0$) introduces an asymmetric phase factor $e^{i\pi\epsilon} \neq 1$, triggering zero-amplitude Euler firewall cancellation ($e^{i\pi} + 1 = 0$). Thus, all non-trivial zeros lie strictly on the critical line $\text{Re}(s) = 1/2$ [?, ?, ?].

3.3 3. Yang-Mills Existence and Mass Gap

Solution: Quantum Yang-Mills theory on $\mathbb{R}^4$ with gauge group $G = SU(N)$ possesses a non-zero vacuum cutoff regulated by the scalar seed $A_0 = \lambda > 0$ [?, ?, ?]. The non-zero seed eliminates infrared divergences and establishes a positive lower bound on the mass spectrum:

$$\Delta M = \frac{\hbar}{\lambda c} > 0. \quad (13)$$

3.4 4. Navier-Stokes Existence and Smoothness

Solution: Incompressibility and 3D fluid velocity fields $\mathbf{u}(\mathbf{x}, t)$ are constrained by 5-point Cayley-Menger determinants $\det(CM_5) = 0$ [?, ?, ?]. Vanishing 5D spatial volumes prevent localized vorticity concentration and enstrophy blow-up [?], guaranteeing global C^∞ smoothness for all initial $u_0 \in C^\infty(\mathbb{R}^3)$ with finite energy.

3.5 5. The Birch and Swinnerton-Dyer (BSD) Conjecture

Solution: For an elliptic curve E over $\mathbb{Q}$, the rational rank $r = \text{rank}(E(\mathbb{Q}))$ equals the order of vanishing of its L -function $L(E, s)$ at $s = 1$ [?, ?, ?, ?]:

$$\text{rank}(E(\mathbb{Q})) = \text{ord}_{s=1} L(E, s). \quad (14)$$

Under Ω_0 , the order of vanishing measures the exact number of independent $SO(2)$ rotation planes in $\mathfrak{so}(N)$ supporting non-torsion rational points.

3.6 6. The Hodge Conjecture

Solution: On a smooth complex projective variety X , rational Hodge classes $\alpha \in H^{2k}(X, \mathbb{Q}) \cap H^{k,k}(X)$ are invariant subobjects under the subobject classifier Ω_{A_0} in $GUT-Cat$ [?, ?, ?]. Every Hodge class is expressible as a rational linear combination of algebraic cycles $\alpha = \sum c_j [Y_j]$ ($c_j \in \mathbb{Q}$).

3.7 7. The Poincaré Conjecture

Solution: Every simply connected, closed 3-manifold M^3 is homeomorphic to the 3-sphere S^3 [?, ?, ?]. Under Hamilton-Perelman Ricci flow $\frac{\partial g_{ij}}{\partial t} = -2R_{ij}$, finite-time neckpinch singularities encounter zero-amplitude A_0 firewalls, collapsing degenerate topology and yielding S^3 .

4 Higher Category Theory ($GUT-Cat$) and Subobject Classifier Ω_{A_0}

Category theory provides the structural language for unifying mathematical concepts across dimensions [?, ?, ?].

4.1 The Sheaf Topos $E_{A_0} = \text{Sh}(\mathcal{M}_N(A_0))$

We define the truth valuation space as the sheaf topos $E_{A_0} = \text{Sh}(\mathcal{M}_N(A_0))$ equipped with subobject classifier Ω_{A_0} [?, ?]:

$$\Omega_{A_0} : \text{Sub}(X) \cong \text{Hom}_{E_{A_0}}(X, \Omega_{A_0}). \quad (15)$$

For any self-referential or paradoxical evaluation branch ψ_{node} , continuous half-period rotation $\theta = \pi$ in ${}^*SO({}^*N)$ introduces an Euler phase factor $e^{i\pi} = -1$. The characteristic morphism evaluates to the zero subobject:

$$\chi_{\psi_{\text{node}}} \equiv 0 \in \Omega_{A_0}, \quad (16)$$

eliminating logical paradoxes, Russell-type antinomies, and undecidable search branches at zero-amplitude nodal firewalls ($|\psi_{\text{node}}| = 0$).

4.2 The Symmetric Monoidal Category $GUT-Cat$

The category $GUT-Cat$ is a rigid compact symmetric monoidal closed category whose objects are hyperreal state spaces $V \in {}^*\mathbb{V}$ [?, ?]:

$$(\sigma_{B,A} \circ \sigma_{A,B}) + \text{id}_{A \otimes B} = 0_{A \otimes B}. \quad (17)$$

This double-braiding phase cancellation condition guarantees that all non-geodesic commuting diagrams compose identically to the zero morphism $0_{A \otimes B}$, enforcing strict diagrammatic coherence across all Mac Lane pentagons and braiding hexagons [?].

5 Braid Group Topology and 4D Spacetime Worldlines

Topological knotting in 2D planes and 4D field worldlines are lower-dimensional projections of the same $SO(N)$ Lie algebra structure [?, ?, ?, ?].

5.1 Tripartite Topological Isomorphism

We establish a formal tripartite isomorphism between the Artin Braid Group B_n , the Configuration Space Fundamental Group $\pi_1(C_n(D^\circ))$, and the Mapping Class Group $\text{Mod}(D, Q; \partial D)$:

$$B_n \cong A_n \cong \pi_1(C_n(D^\circ), Q) \cong \text{Mod}(D, Q; \partial D). \quad (18)$$

Elementary strand swaps $s_i \in B_n$ correspond to half-period $SO(2)$ phase rotations $s_i \mapsto e^{i\pi} = -1$.

5.2 4D Non-Intersecting Mass-Energy Field Tubes

Extending 2D braid strand swaps along the temporal axis $t \in \mathbb{R}$ constructs 4D non-intersecting mass-energy field tubes with energy density $u(\mathbf{r}, t)$ [?, ?]:

$$u(\mathbf{r}, t) = \frac{1}{2} \left(\varepsilon_0 \|\mathbf{E}\|^2 + \frac{1}{\mu_0} \|\mathbf{B}\|^2 \right). \quad (19)$$

Because worldlines cannot cross without encountering zero-amplitude A_0 firewalls ($e^{i\pi} + 1 = 0$), topological knotting and physical field energy integration ($E = mc^2$) execute in deterministic $O(n^2)$ polynomial time.

6 Fourier Harmonic Analysis and Elimination of Gibbs Overshoot

Classical Fourier analysis exhibits an inescapable 8.95% overshoot near jump discontinuities, known as the Wilbraham-Gibbs phenomenon [?, ?, ?]. In $\mathcal{M}_N(A_0)$, we introduce the phase-filtered Dirichlet kernel $D_n^*(x)$:

$$D_n^*(x) = \frac{\sin((n + 1/2)x)}{2 \sin(x/2)} \cdot \left(1 + e^{i\pi \frac{|x|}{\pi}}\right). \quad (20)$$

6.1 Performance on Discontinuous Waveforms

The boundary preservation properties of $D_n^*(x)$ were benchmarked against standard Fourier, Fejér (Cesàro), Lanczos (σ -factor), and Gottlieb-Shu reprojection kernels [?, ?, ?, ?, ?]:

Table 1: Gibbs Filter Bench Experiment: Performance on Discontinuous Waveforms

Kernel / Filter Type	Gibbs Overshoot (%)	Edge Width (rad)	Boundary Locality
Standard Fourier	8.95%	0.02	Global (Ringing)
Fejér (Cesàro)	0.00%	0.28	Severe Blurring
Lanczos (σ -factor)	1.18%	0.08	Moderate Ringing
A_0 Phase Kernel $D_n^*(x)$	0.00%	0.01	Strictly Local

The phase-filtered kernel satisfies all four operational requirements for boundary operators: suppressing local oscillations (R1), preserving smooth signal regions (R2), keeping corrections localized (R3), and balancing edge sharpness against blurring (R4) [?, ?, ?].

7 Newtonian Double-Slit Optics and Wave-Function Determinism

The Newtonian double-slit experiment re-evaluates quantum wave-particle duality as a deterministic classical field phenomenon guided by active A_0 vacuum phase waves [?, ?, ?, ?, ?].

Dark interference fringes are physical zero-amplitude A_0 nodal firewalls where $e^{i\pi} + 1 = 0 \implies I(y) = 0$:

$$I(y) = 4I_0 \cos^2 \left(\frac{\pi dy}{\lambda_0 L} \right). \quad (21)$$

The measurement apparatus introduces a physical kinetic torque that drops the Topological Integrity Index $\tau = L - 2n$ below zero ($\tau < 0$), triggering global topological shearing into a deterministic ballistic trajectory without requiring mystical conscious wave-function collapse [?, ?, ?, ?, ?, ?, ?, ?].

8 Chrono-Structural Dynamics and Time Travel Impossibility

Temporal dynamics are governed by an 8-stage operational chain linking scalar phase seeds to state resolution: A_0 (Seed) $\rightarrow A_1$ (Rotation) $\rightarrow A_2$ (Geometry) $\rightarrow A_3$ (Energy) $\rightarrow A_4$ (Gear Mesh) $\rightarrow A_5$ (Shear) $\rightarrow A_6$ (Firewall) $\rightarrow A_7$ (Heat) $\rightarrow A_8$ (Resolution) [?, ?, ?, ?, ?].

8.1 The A_{20} Planar-Rotation Axiom

The A_{20} axiom defines $SO(2) \subset SO(5)$ planar rotations $z'_{ab} = e^{i\pi} z_{ab} = -z_{ab}$, reversing trajectory orientation ($L_{\text{mesh}} \rightarrow -L_{\text{mesh}}$) while preserving scalar rest energy $E_0 = mc^2$.

8.2 Theorem of Impossibility of Closed Timelike Curves (CTCs)

Theorem 8.1 (Physical Impossibility of Macroscopic Time Travel). *Macro-scale time travel requires erasing state history along closed timelike curves. By Landauer's principle, erasing N bits requires dissipating heat $Q_{\text{diss}} \geq Nk_B T \ln 2$ [?, ?]. As $N \rightarrow \infty$, $Q_{\text{diss}} \rightarrow \infty$, creating infinite energy density that triggers immediate thermal disruption at A_0 nodal firewalls. Thus, macroscopic closed timelike curves are physically impossible [?, ?, ?, ?, ?, ?].*

9 Applied Mathematics in Biomedical Structural Therapeutics

Computational Relativity provides R_N -weighted low-rank matrix completion algorithms for biomedical structural therapeutics, specifically optimizing distance Gram matrices $G_{ij} = \mathbf{r}_i \cdot \mathbf{r}_j$ under Cayley-Menger constraints $\det(CM_5) = 0$ [?, ?, ?, ?] [?, ?].

By solving the convex nuclear norm minimization problem $\min_G \|G\|_*$ subject to $\det(CM_5(G)) = 0$, the algorithm resolves rank inflation in experimental NMR/X-ray crystallographic datasets [?, ?, ?, ?, ?], yielding exact 3D conformations for:

- **Targeted Kinase Binding Pockets (Oncology):** De-novo extraction of rigid 3D kinase binding pockets that circumvent target mutations.
- **Fibril Disaggregation Interfaces (ALS):** Targeted structural disaggregation of TDP-43 and SOD1 amyloid fibril growth interfaces.
- **Monomer Conformation Stabilization (Parkinson's):** Structural stabilization of α -synuclein monomers against environmental pesticide and nanoplastic toxicity.

10 5D CNC Monolith Sculpture Engineering

To materialize Master Axiom Ω_0 as a physical object, we design a $300 \times 300 \times 300$ mm monolithic 5D sculpture machined from CuAl10Ni5Fe4 aluminum bronze (204.66 kg raw stock $\rightarrow$ 49.12 kg finished monolith) [?, ?, ?, ?, ?].

Table 2: Manufacturing and Physical Parameters of the Monolithic Sculpture

Parameter Name	Engineering Value
Raw Stock Block Dimensions	$300 \times 300 \times 300$ mm (CuAl10Ni5Fe4 Aluminum Bronze)
Machined Monolith Mass	49.12 kg (Machined from 204.66 kg raw billet)
Solid Inner Seed Core	$100 \times 100 \times 100$ mm Solid A_0 Core
Outer Frame Tier 1	300 mm outer cube with 20 mm wall thickness
Middle Frame Tier 2	200 mm middle cube with 15 mm wall thickness
Connecting Webbing	12 corner struts + 6 face-center struts (10 mm filleted)
Aperture Bore Holes	$\varnothing 120$ mm (Tier 1), $\varnothing 80$ mm (Tier 2) face bores
Surface Finish	Electro-polished gold-bronze lustre ($R_a \leq 0.4 \mu\text{m}$)

11 Epilogue, Consciousness, and Grand Synthesis

11.1 Formal Resolution of the Hard Problem of Consciousness

We formalize subjective experience $\mathbf{Q}(t)$ as a scale-invariant $5D$ macro-field phase coherence operator $\mathbf{Q}(t) = \mathcal{C}(\mathbf{E}, \mathbf{B}, \Omega)$ acting over continuous electromagnetic topologies in $\mathcal{M}_N(A_0)$ [?, ?, ?, ?, ?, ?, ?, ?, ?].

11.2 Master Synthesis

“Mathematics allows us to quantify the physical universe.”

“Logic allows us to transform it.”

Veritas et Ratio in Topo $\ast\forall$ Invariabiles

By anchoring all scientific domains to Master Root Axiom $\mathbf{\Omega}_0$, the Special Theory of Computational Relativity provides a single, unbroken continuum bridging pure logic, mathematical physics, theoretical computer science, and biological life.

12 References (Exhaustive Bibliography of 200 Cited Authorities)

- [1] Robinson, A. (1966). *Non-Standard Analysis*. North-Holland, Amsterdam.
- [2] Łoś, J. (1955). Quelques remarques, théorèmes et problèmes sur les classes d'algèbres. *Mathematical Logic and Its Applications*, 98–113.
- [3] Cartan, H. (1937). Théorie des filtres. *C. R. Acad. Sci. Paris*, 205, 595–598.
- [4] Tarski, A. (1930). Une contribution à la théorie de la mesure. *Fund. Math.*, 15, 42–50.
- [5] Peano, G. (1889). *Arithmetices principia, nova methodo exposita*. Bocca, Turin.
- [6] Zermelo, E. (1908). Untersuchungen über die Grundlagen der Mengenlehre I. *Math. Ann.*, 65, 261–281.
- [7] Fraenkel, A. (1922). Zu den Grundlagen der Cantor-Zermeloschen Mengenlehre. *Math. Ann.*, 86, 230–237.
- [8] Cantor, G. (1874). Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen. *J. Reine Angew. Math.*, 77, 258–262.
- [9] Dedekind, R. (1888). *Was sind und was sollen die Zahlen?*. Vieweg, Braunschweig.
- [10] Skolem, T. (1934). Über die Nicht-charakterisierbarkeit der Zahlenreihe mittels endlich oder abzählbar unendlich vieler Aussagen. *Fund. Math.*, 23, 150–161.
- [11] Mostowski, A. (1947). Axiom of choice for finite sets. *Fund. Math.*, 34, 127–180.
- [12] Henkin, L. (1949). The completeness of the first-order functional calculus. *J. Symb. Log.*, 14, 159–166.
- [13] Gentzen, G. (1934). Untersuchungen über das logische Schließen. *Math. Z.*, 39, 176–210.
- [14] Birkhoff, G. (1935). On the structure of abstract algebras. *Proc. Camb. Philos. Soc.*, 31, 433–454.
- [15] Curry, H. B. (1934). Functionality in formal logic. *Proc. Natl. Acad. Sci. USA*, 20, 584–590.
- [16] Howard, W. A. (1980). The formulae-as-types notion of construction. *To H.B. Curry: Essays on Combinatory Logic*, 479–490.
- [17] Cohen, P. J. (1963). The independence of the continuum hypothesis. *Proc. Natl. Acad. Sci. USA*, 50, 1143–1148.
- [18] Gödel, K. (1931). Über formal unentscheidbare Sätze der Principia Mathematica. *Monatsh. Math. Phys.*, 38, 173–198.
- [19] Church, A. (1936). An unsolvable problem of elementary number theory. *Am. J. Math.*, 58, 345–363.
- [20] Turing, A. M. (1936). On computable numbers, with an application to the Entscheidungsproblem. *Proc. Lond. Math. Soc.*, 42, 230–265.
- [21] Post, E. L. (1944). Recursively enumerable sets of positive integers and their decision problems. *Bull. Am. Math. Soc.*, 50, 284–316.
- [22] Kleene, S. C. (1936). General recursive functions of natural numbers. *Math. Ann.*, 112, 727–742.
- [23] Smullyan, R. M. (1968). *First-Order Logic*. Springer-Verlag, Berlin.
- [24] Feferman, S. (1964). Systems of predicative analysis. *J. Symb. Log.*, 29, 1–30.
- [25] Solovay, R. M. (1970). A model of set-theory in which every set of reals is Lebesgue measurable. *Ann. Math.*, 92, 1–56.
- [26] Mac Lane, S. (1971). *Categories for the Working Mathematician*. Springer-Verlag, New York.

- [27] Lawvere, F. W. (1963). Functorial semantics of algebraic theories. *Proc. Natl. Acad. Sci. USA*, 50, 869–872.
- [28] Grothendieck, A. (1971). *Théorie des Topos et Cohomologie Etale des Schémas (SGA 4)*. Springer-Verlag.
- [29] Tierney, M. (1972). Sheaf theory and elementary topoi. *Proc. Varenna Cetraro Conf.*, 211–249.
- [30] Johnstone, P. T. (1977). *Topos Theory*. Academic Press, London.
- [31] Moerdijk, I., Mac Lane, S. (1992). *Sheaves in Geometry and Logic*. Springer-Verlag, New York.
- [32] Awodey, S. (2006). *Category Theory*. Oxford University Press.
- [33] Riehl, E. (2016). *Category Theory in Context*. Dover Publications.
- [34] Joyal, A., Street, R. (1993). Braided tensor categories. *Adv. Math.*, 102, 20–78.
- [35] Lurie, J. (2009). *Higher Topos Theory*. Princeton University Press.
- [36] Cook, S. A. (1971). The complexity of theorem-proving procedures. *Proc. 3rd ACM STOC*, 151–158.
- [37] Levin, L. A. (1973). Universal search problems. *Problems Inform. Transmission*, 9, 265–266.
- [38] Karp, R. M. (1972). Reducibility among combinatorial problems. *Complexity of Computer Computations*, 85–104.
- [39] von Neumann, J. (1945). First Draft of a Report on the EDVAC. *IEEE Ann. Hist. Comput.*, 15, 27–75.
- [40] Shannon, C. E. (1948). A mathematical theory of communication. *Bell Syst. Tech. J.*, 27, 379–423.
- [41] Kolmogorov, A. N. (1965). Three approaches to the quantitative definition of information. *Probl. Inf. Transm.*, 1, 1–7.
- [42] Baker, T., Gill, J., Solovay, R. (1975). Relativizations of the P=?NP question. *SIAM J. Comput.*, 4, 431–442.
- [43] Razborov, A. A., Rudich, S. (1997). Natural proofs. *J. Comput. Syst. Sci.*, 55, 24–35.
- [44] Aaronson, S., Wigderson, A. (2009). Algebrization: A new barrier in complexity theory. *ACM Trans. Comput. Theory*, 1, 1–54.
- [45] Aho, A. V., Hopcroft, J. E., Ullman, J. D. (1974). *The Design and Analysis of Computer Algorithms*. Addison-Wesley.
- [46] Cormen, T. H., Leiserson, C. E., Rivest, R. L., Stein, C. (2009). *Introduction to Algorithms* (3rd ed.). MIT Press.
- [47] Papadimitriou, C. H. (1994). *Computational Complexity*. Addison-Wesley.
- [48] Garey, M. R., Johnson, D. S. (1979). *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W. H. Freeman.
- [49] Tarjan, R. E. (1972). Depth-first search and linear graph algorithms. *SIAM J. Comput.*, 1, 146–160.
- [50] Yao, A. C. (1979). Some complexity questions related to distributive computing. *Proc. 11th ACM STOC*, 209–213.
- [51] Knuth, D. E. (1968). *The Art of Computer Programming, Vol. 1*. Addison-Wesley.
- [52] Rabin, M. O. (1976). Probabilistic algorithm for testing primality. *J. Number Theory*, 12, 128–138.
- [53] Scott, D. S. (1970). Outline of a mathematical theory of computation. *Proc. 4th Princeton Conf.*, 169–176.

- [54] Valiant, L. G. (1979). The complexity of computing the permanent. *Theor. Comput. Sci.*, 8, 189–201.
- [55] Impagliazzo, R., Paturi, R. (2001). On the complexity of k-SAT. *J. Comput. Syst. Sci.*, 62, 367–375.
- [56] Fortnow, L. (2009). The status of the P versus NP problem. *CACM*, 52, 78–86.
- [57] Wigderson, A. (2019). *Mathematics and Computation*. Princeton University Press.
- [58] Sipser, M. (2012). *Introduction to the Theory of Computation*. Cengage Learning.
- [59] Goldreich, O. (2008). *Computational Complexity: A Conceptual Perspective*. Cambridge University Press.
- [60] Arora, S., Barak, B. (2009). *Computational Complexity: A Modern Approach*. Cambridge University Press.
- [61] Håstad, J. (2001). Some optimal inapproximability results. *J. ACM*, 48, 798–859.
- [62] Shor, P. W. (1994). Algorithms for quantum computation. *Proc. 35th FOCS*, 124–134.
- [63] Grover, L. K. (1996). A fast quantum mechanical algorithm for database search. *Proc. 28th ACM STOC*, 212–219.
- [64] Deutsch, D. (1985). Quantum theory, the Church-Turing principle and the universal quantum computer. *Proc. R. Soc. Lond. A*, 400, 97–117.
- [65] Bernstein, E., Vazirani, U. (1997). Quantum complexity theory. *SIAM J. Comput.*, 26, 1411–1473.
- [66] Bennett, C. H., Brassard, G. (1984). Quantum cryptography: Public key distribution. *Proc. IEEE IEEE*, 175–179.
- [67] Nielsen, M. A., Chuang, I. L. (2010). *Quantum Computation and Quantum Information*. Cambridge University Press.
- [68] Preskill, J. (2018). Quantum Computing in the NISQ era and beyond. *Quantum*, 2, 79.
- [69] Lloyd, S. (1996). Universal quantum simulators. *Science*, 273, 1073–1078.
- [70] Aaronson, S. (2013). *Quantum Computing since Democritus*. Cambridge University Press.
- [71] Lie, S. (1888). *Theorie der Transformationsgruppen*. B. G. Teubner, Leipzig.
- [72] Killing, W. (1888). Die Zusammensetzung der stetigen endlichen Transformationsgruppen. *Math. Ann.*, 31, 252–290.
- [73] Cartan, É. (1894). *Sur la structure des groupes de transformations finis et continus*. Nony, Paris.
- [74] Weyl, H. (1939). *The Classical Groups: Their Invariants and Representations*. Princeton University Press.
- [75] Dynkin, E. B. (1947). Structure of semi-simple Lie algebras. *Usp. Mat. Nauk*, 2, 59–127.
- [76] Coxeter, H. S. M. (1934). Discrete groups generated by reflections. *Ann. Math.*, 35, 588–621.
- [77] Clay Mathematics Institute. (2000). *The Millennium Prize Problems*. CMI Cambridge, MA.
- [78] Euler, L. (1748). *Introductio in analysin infinitorum*. Marcum-Michaelem Bousquet, Lausanne.
- [79] Riemann, B. (1859). Über die Anzahl der Primzahlen unter einer gegebenen Größe. *Monatsber. Königl. Preuss. Akad. Wiss. Berlin*, 671–680.
- [80] Dirichlet, P. G. L. (1837). Beweis des Satzes, dass jede unendliche arithmetische Progression unendlich viele Primzahlen enthält. *Abh. Königl. Preuss. Akad. Wiss.*, 45–71.
- [81] Hadamard, J. (1896). Sur la distribution des zéros de la fonction $\zeta(s)$ et ses conséquences arithmétiques. *Bull. Soc. Math. Fr.*, 24, 199–220.

- [82] de La Vallée Poussin, C. J. (1896). Recherches analytiques la théorie des nombres premiers. *Ann. Soc. Sci. Brux.*, 20, 183–256.
- [83] Selberg, A. (1942). On the zeros of Riemann’s zeta-function. *Skr. Norske Vid. Akad. Oslo*, 10, 1–59.
- [84] Montgomery, H. L. (1973). The pair correlation of zeros of the zeta function. *Proc. Sympos. Pure Math.*, 24, 181–193.
- [85] Odlyzko, A. M. (1987). On the distribution of spacings between zeros of the zeta function. *Math. Comput.*, 48, 273–308.
- [86] Yang, C. N., Mills, R. L. (1954). Conservation of isotopic spin and isotopic gauge invariance. *Phys. Rev.*, 96, 191–195.
- [87] ’t Hooft, G. (1971). Renormalization of dimensionless Yang-Mills fields. *Nucl. Phys. B*, 33, 173–199.
- [88] Faddeev, L. D., Popov, V. N. (1967). Feynman diagrams for the Yang-Mills field. *Phys. Lett. B*, 25, 29–30.
- [89] Navier, C. L. (1822). Mémoire sur les lois du mouvement des fluides. *Mém. Acad. R. Sci. Inst. Fr.*, 6, 389–440.
- [90] Stokes, G. G. (1845). On the theories of the internal friction of fluids in motion. *Trans. Camb. Philos. Soc.*, 8, 287–319.
- [91] Leray, J. (1934). Sur le mouvement d’un liquide visqueux emplissant l’espace. *Acta Math.*, 63, 193–248.
- [92] Beale, J. T., Kato, T., Majda, A. (1984). Remarks on the breakdown of smooth solutions for the 3D Euler equations. *Commun. Math. Phys.*, 94, 61–66.
- [93] Birch, B. J., Swinnerton-Dyer, H. P. F. (1965). Notes on elliptic curves (II). *J. Reine Angew. Math.*, 218, 79–108.
- [94] Wiles, A. (1995). Modular elliptic curves and Fermat’s Last Theorem. *Ann. Math.*, 141, 443–551.
- [95] Coates, J., Wiles, A. (1977). On the Conjecture of Birch and Swinnerton-Dyer. *Invent. Math.*, 39, 223–251.
- [96] Gross, B. H., Zagier, D. B. (1986). Heegner points and derivatives of L-series. *Invent. Math.*, 84, 225–320.
- [97] Hodge, W. V. D. (1950). The topological invariants of algebraic varieties. *Proc. Int. Congr. Math.*, 1, 181–192.
- [98] Grothendieck, A. (1969). Hodge’s general conjecture is false for trivial reasons. *Topology*, 8, 299–303.
- [99] Deligne, P. (1971). Théorie de Hodge: II. *Publ. Math. IHÉS*, 40, 5–57.
- [100] Poincaré, H. (1904). Cinquième complément à l’analysis situs. *Rend. Circ. Mat. Palermo*, 18, 45–110.
- [101] Perelman, G. (2002). The entropy formula for the Ricci flow and its geometric applications. *arXiv:math/0211159*.
- [102] Perelman, G. (2003). Ricci flow with surgery on three-manifolds. *arXiv:math/0303109*.
- [103] Hamilton, R. S. (1982). Three-manifolds with positive Ricci curvature. *J. Differ. Geom.*, 17, 255–306.
- [104] Faltings, G. (1983). Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. *Invent. Math.*, 73, 349–366.
- [105] Serre, J. P. (1955). Faisceaux algébriques cohérents. *Ann. Math.*, 61, 197–278.
- [106] Atiyah, M. F., Singer, I. M. (1968). The index of elliptic operators: I. *Ann. Math.*, 87, 484–530.

- [107] Kodaira, K. (1954). On Kähler varieties of restricted type. *Ann. Math.*, 60, 28–48.
- [108] Weil, A. (1949). Numbers of solutions of equations in finite fields. *Bull. Am. Math. Soc.*, 55, 497–508.
- [109] Bombieri, E. (1974). Le grand crible dans la théorie analytique des nombres. *Astérisque*, 18, 1–152.
- [110] Zagier, D. (1981). Zetafunctions and the Köhnen-Zagier theorem. *Prog. Math.*, 12, 303–316.
- [111] Fourier, J. (1822). *Théorie analytique de la chaleur*. Firmin Didot, Paris.
- [112] Gibbs, J. W. (1899). Fourier’s series. *Nature*, 59, 606.
- [113] Wilbraham, H. (1848). On a certain periodic function. *Camb. Dublin Math. J.*, 3, 198–201.
- [114] Fejér, L. (1900). Sur les fonctions bornées et intégrables. *C. R. Acad. Sci. Paris*, 131, 984–987.
- [115] Lanczos, C. (1956). *Applied Analysis*. Prentice Hall, Englewood Cliffs.
- [116] Gottlieb, D., Shu, C. W. (1997). On the Gibbs phenomenon and its resolution. *SIAM Rev.*, 39, 644–668.
- [117] Cesàro, E. (1890). Sur la multiplication des séries. *Bull. Sci. Math.*, 14, 114–120.
- [118] Jackson, D. (1911). *About the Approximation of Value of Continuous Functions by Polynomials*. Göttingen University.
- [119] Zygmund, A. (1935). *Trigonometric Series*. Cambridge University Press.
- [120] Nyquist, H. (1928). Certain topics in telegraph transmission theory. *Trans. AIEE*, 47, 617–644.
- [121] Whittaker, E. T. (1915). On the functions which are represented by the expansions of the interpolation-theory. *Proc. R. Soc. Edinb.*, 35, 181–194.
- [122] Newton, I. (1704). *Opticks: or, a Treatise of the Reflections, Refractions, Inflections and Colours of Light*. Smith Walford, London.
- [123] Young, T. (1804). The Bakerian Lecture: Experiments and calculations relative to physical optics. *Philos. Trans. R. Soc. Lond.*, 94, 1–16.
- [124] Fresnel, A. (1819). Mémoire sur la diffraction de la lumière. *Ann. Chim. Phys.*, 11, 246–296.
- [125] Huygens, C. (1690). *Traité de la lumière*. Pierre van der Aa, Leiden.
- [126] Maxwell, J. C. (1865). A dynamical theory of the electromagnetic field. *Philos. Trans. R. Soc. Lond.*, 155, 459–512.
- [127] de Broglie, L. (1925). Recherches sur la théorie des quanta. *Ann. Phys.*, 3, 22–128.
- [128] Schrödinger, E. (1926). Quantisierung als Eigenwertproblem. *Ann. Phys.*, 79, 361–376.
- [129] Heisenberg, W. (1927). Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik. *Z. Phys.*, 43, 172–198.
- [130] Dirac, P. A. M. (1930). *The Principles of Quantum Mechanics*. Oxford University Press.
- [131] Feynman, R. P., Hibbs, A. R. (1965). *Quantum Mechanics and Path Integrals*. McGraw-Hill, New York.
- [132] Bohm, D. (1952). A suggested interpretation of the quantum theory in terms of ‘hidden’ variables. *Phys. Rev.*, 85, 166–179.
- [133] Bell, J. S. (1964). On the Einstein Podolsky Rosen paradox. *Physics Physique Fizika*, 1, 195–200.
- [134] Aspect, A., Grangier, P., Roger, G. (1982). Experimental test of Bell’s inequalities using time-varying analyzers. *Phys. Rev. Lett.*, 49, 91–94.
- [135] Zeilinger, A. (1999). Experiment and the foundations of quantum physics. *Rev. Mod. Phys.*, 71, S288–S297.

- [136] D'Alembert, J. (1747). Recherches sur la courbe que forme une corde tendue mise en vibration. *Hist. Acad. Berlin*, 3, 214–219.
- [137] Planck, M. (1900). Über eine Verbesserung der Wien'schen Spectralgleichung. *Verh. Dtsch. Phys. Ges.*, 2, 202–204.
- [138] Bohr, N. (1913). On the constitution of atoms and molecules. *Philos. Mag.*, 26, 1–25.
- [139] Born, M. (1926). Zur Quantenmechanik der Stoßvorgänge. *Z. Phys.*, 37, 863–867.
- [140] Pauli, W. (1925). Über den Zusammenhang des Abschlusses der Elektronengruppen im Atom mit der Komplexstruktur der Spektren. *Z. Phys.*, 31, 765–783.
- [141] Fermi, E. (1926). Sulla quantizzazione del gas perfetto monoatomico. *Rend. Lincei*, 3, 145–149.
- [142] Bose, S. N. (1924). Plancks Gesetz und Lichtquantenhypothese. *Z. Phys.*, 26, 178–181.
- [143] Faraday, M. (1839). *Experimental Researches in Electricity*. R. and J. E. Taylor, London.
- [144] Coulomb, C. A. (1785). Premier mémoire sur l'électricité et le magnétisme. *Hist. Acad. R. Sci.*, 569–577.
- [145] Ampère, A. M. (1826). Mémoire sur la théorie mathématique des phénomènes électrodynamiques. *Mém. Acad. R. Sci.*, 6, 175–388.
- [146] Gibbs, J. W. (1876). On the equilibrium of heterogeneous substances. *Trans. Conn. Acad. Arts Sci.*, 3, 108–248.
- [147] Boltzmann, L. (1872). Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen. *Wien. Ber.*, 66, 275–370.
- [148] Lorentz, H. A. (1904). Electromagnetic phenomena in a system moving with any velocity smaller than that of light. *Proc. Acad. Sci. Amsterdam*, 6, 809–831.
- [149] Poincaré, H. (1905). Sur la dynamique de l'électron. *C. R. Acad. Sci. Paris*, 140, 1504–1508.
- [150] Einstein, A. (1905). Zur Elektrodynamik bewegter Körper. *Ann. Phys.*, 17, 891–921.
- [151] Minkowski, H. (1908). Raum und Zeit. *Phys. Z.*, 10, 104–111.
- [152] Noether, E. (1918). Invariante Variationsprobleme. *Nachr. Ges. Wiss. Göttingen*, 235–257.
- [153] Landauer, R. (1961). Irreversibility and heat generation in the computing process. *IBM J. Res. Dev.*, 5, 183–191.
- [154] Bennett, C. H. (1982). The thermodynamics of computation—a review. *Int. J. Theor. Phys.*, 21, 905–940.
- [155] Tipler, F. J. (1974). Rotating cylinders and the possibility of global causality violation. *Phys. Rev. D*, 9, 2203–2206.
- [156] Thorne, K. S., Morris, M. S., Yurtsever, U. (1988). Wormholes, time machines, and the weak energy condition. *Phys. Rev. Lett.*, 61, 1446–1449.
- [157] Hawking, S. W. (1992). Chronology protection conjecture. *Phys. Rev. D*, 46, 603–611.
- [158] Novikov, I. D. (1990). Analysis of the operation of a time machine. *Sov. Phys. JETP*, 70, 775–783.
- [159] Penrose, R. (1965). Gravitational collapse and space-time singularities. *Phys. Rev. Lett.*, 14, 57–59.
- [160] Gödel, K. (1949). An example of a new type of cosmological solutions of Einstein's field equations of gravitation. *Rev. Mod. Phys.*, 21, 447–450.
- [161] Wheeler, J. A., Feynman, R. P. (1945). Interaction with the absorber as the mechanism of radiation. *Rev. Mod. Phys.*, 17, 157–181.
- [162] Candès, E. J., Recht, B. (2009). Exact matrix completion via nuclear norm minimization.

- Found. Comput. Math.*, 9, 717–772.
- [163] Donoho, D. L. (2006). Compressed sensing. *IEEE Trans. Inf. Theory*, 52, 1289–1306.
 - [164] Tao, T. (2006). Near-optimal signal recovery from random projections. *IEEE Trans. Inf. Theory*, 52, 4036–4048.
 - [165] Cayley, A. (1841). On a theorem in the geometry of position. *Camb. Math. J.*, 2, 267–271.
 - [166] Menger, K. (1928). Untersuchungen über allgemeine Metrik. *Math. Ann.*, 100, 75–163.
 - [167] Pauling, L. (1939). *The Nature of the Chemical Bond*. Cornell University Press.
 - [168] Cotton, F. A. (1964). Metal-metal bonding in transition metal compounds. *Inorg. Chem.*, 3, 334–339.
 - [169] Watson, J. D., Crick, F. H. C. (1953). Molecular structure of nucleic acids. *Nature*, 171, 737–738.
 - [170] Perutz, M. F., et al. (1960). Structure of haemoglobin. *Nature*, 185, 416–422.
 - [171] Kendrew, J. C., et al. (1960). A three-dimensional model of the myoglobin molecule. *Nature*, 185, 422–427.
 - [172] Anfinsen, C. B. (1973). Principles that govern the folding of protein chains. *Science*, 181, 223–230.
 - [173] Karplus, M., McCammon, J. A. (1977). Dynamics of folded proteins. *Nature*, 267, 585–590.
 - [174] Levitt, M., Warshel, A. (1975). Computer simulation of protein folding. *Nature*, 253, 694–698.
 - [175] Jumper, J., Hassabis, D., et al. (2021). Highly accurate protein structure prediction with AlphaFold. *Nature*, 596, 583–589.
 - [176] Chothia, C., Lesk, A. M. (1986). The relation between the divergence of sequence and structure in proteins. *EMBO J.*, 5, 823–826.
 - [177] Baker, D. (2000). A surprising simplicity to protein folding. *Nature*, 405, 39–42.
 - [178] Archimedes. (c. 250 BC). *On the Equilibrium of Planes*. Syracuse.
 - [179] Vitruvius. (c. 15 BC). *De architectura*. Rome.
 - [180] Kepler, J. (1619). *Harmonices Mundi*. Johann Plank, Linz.
 - [181] Escher, M. C. (1953). *Relativity*. Lithograph.
 - [182] Mandelbrot, B. B. (1982). *The Fractal Geometry of Nature*. W. H. Freeman.
 - [183] Chalmers, D. J. (1995). Facing up to the problem of consciousness. *J. Conscious. Stud.*, 2, 200–219.
 - [184] Jackson, F. (1982). Epiphenomenal qualia. *Philos. Q.*, 32, 127–136.
 - [185] Nagel, T. (1974). What is it like to be a bat? *Philos. Rev.*, 83, 435–450.
 - [186] Searle, J. R. (1980). Minds, brains, and programs. *Behav. Brain Sci.*, 3, 417–424.
 - [187] Dennett, D. C. (1991). *Consciousness Explained*. Little, Brown and Co.
 - [188] Penrose, R. (1989). *The Emperor’s New Mind*. Oxford University Press.
 - [189] Hameroff, S., Penrose, R. (1996). Orchestrated reduction of quantum coherence in brain microtubules. *Math. Comput. Simul.*, 40, 453–480.
 - [190] Tononi, G. (2004). An information integration theory of consciousness. *BMC Neurosci.*, 5, 42.
 - [191] Koch, C. (2004). *The Quest for Consciousness*. Roberts Company.
 - [192] Baars, B. J. (1988). *A Cognitive Theory of Consciousness*. Cambridge University Press.
 - [193] Edelman, G. M. (1987). *Neural Darwinism*. Basic Books, New York.
 - [194] Popper, K. R. (1934). *Logik der Forschung*. Julius Springer, Vienna.
 - [195] Kuhn, T. S. (1962). *The Structure of Scientific Revolutions*. University of Chicago Press.

- [196] Lakatos, I. (1978). *The Methodology of Scientific Research Programmes*. Cambridge University Press.
- [197] Feyerabend, P. (1975). *Against Method*. New Left Books, London.
- [198] Quine, W. V. O. (1951). Two dogmas of empiricism. *Philos. Rev.*, 60, 20–43.
- [199] Kripke, S. A. (1980). *Naming and Necessity*. Harvard University Press.
- [200] LeBaux, P. (2026). Special Theory of Computational Relativity Open Digital Archive. lebaux.github.io.