

# The Derivative as a Function

## 1 The Definition of a Derivative

Instead of plugging in a number for  $a$ , we can evaluate the derivative at the variable  $x$ , and get the derivative as a function, as a function  $f'(x)$ , defined as

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Recall that the derivative, evaluated at  $x$ , tells the slope of  $f(x)$  at  $x$ .

### 1.1 Finding $f'(x)$ using the definition

#### 1.1.1 Example

$f(x) = \sqrt{2x}$  Find  $f'(x)$ .

#### 1.1.2 Example

$f(x) = 3x + 4$  Find  $f'(x)$ .

### 1.1.3 Example

$$g(x) = \frac{x+3}{2x+5}$$

Find  $f'(x)$ .

### 1.1.4 Example

$$f(x) = x^2 + 6x - 1$$

Find  $f'(x)$ .

**1.1.6 Example**

$$f(x) = \frac{1}{2x+3}$$

Find  $f'(x)$ .

**1.1.7 Example**




$$f(x) = \sqrt{x+5}$$

Find  $f'(x)$ .

## 2 The Graph of $f'(x)$




If we graph  $f(x)$  and  $f'(x)$  on the same set of axes, we can see that the value of  $f'(x)$  tells the slope of  $f(x)$ .

### 2.0.1 Examples



### 2.0.2 Examples

Draw a rough sketch of the derivative of the functions shown here:



## 3 Differentiability

**Definition:**  $f(x)$  is **differentiable** at  $a$  if  $f'(a)$  exists;  $f(x)$  is differentiable on the open interval  $(a, b)$  if it is differentiable at every number in the interval.

Graphically, there are three reasons that  $f(x)$  might not be differentiable at  $a$ .

- $f(x)$  is discontinuous at  $a$ ,
- $f(x)$  has a sharp corner (aka a *cusp*) at  $a$ , or
- $f(x)$  has a vertical tangent at  $a$ ,

### 3.0.1 Examples

Look at the graphs to determine where the following functions are not differentiable.

$$f(x) = \sqrt[3]{x}$$

$$g(x) = |x + 3|$$

$$h(x) = \tan x$$

$$y = \lfloor x \rfloor$$

## 4 Notation

There are several different notations used to denote derivatives. Suppose  $y = f(x)$ . We can denote its derivative in several ways:

- Prime notation:  $y'$ ,  $f'(x)$
- Leibniz notation:  $\frac{dy}{dx}$ ,  $\frac{df}{dx}$
- Differential operator:  $\frac{d}{dx}(y)$ ,  $\frac{d}{dx}(f)$ ,  $Df(x)$ ,  $D_x(f(x))$

The process of taking a derivative is called **differentiation**. Don't call it "deriving" or I will throw things at you.

### 4.0.1 Example

Evaluate  $\frac{d}{dx}(3x + 4)$

#### 4.0.2 Example

Evaluate  $\frac{d}{dr}(2\pi r)$

#### 4.0.3 Example

Evaluate  $\frac{d}{dt}\left(\frac{2}{3} - \frac{3}{7}t\right)$

## 5 Higher Derivatives; Velocity and Acceleration

If we take the derivative of  $f'(x)$ , we get the **second derivative**, which is denoted by either  $f''(x)$ , or  $\frac{d^2y}{dx^2}$