

UNIVERSITY OF LONDON

[E1.11 (Maths) ISE 2007]

B.ENG. AND M.ENG. EXAMINATIONS 2007

For Internal Students of Imperial College London

This paper is also taken for the relevant examination for the Associateship of the City & Guilds of London Institute

INFORMATION SYSTEMS ENGINEERING E1.11

MATHEMATICS

Date Wednesday 30th May 2007 10.00 am - 1.00 pm

Answer ANY SEVEN questions

Answers to Section A questions must be written in a different answer book from answers to Section B questions.

[Before starting, please make sure that the paper is complete. There should be SEVEN pages, with a total of NINE questions. Ask the invigilator for a replacement if this copy is faulty.]

Copyright of the University of London 2007

SECTION A

[E1.11 (Maths) ISE 2007]

1. (i) Find all possible values of the following complex numbers. Give your answers in the form $x + iy$ (with x and y real):

(a) i^{12} ;

(b) $e^{-i\pi/6}$;

(c) $1^{1/3}$;

- (ii) Find all the solutions of the equation $\cos^2 z = 4$.

Give your answer in the form $x + iy$ (with x and y real).

2. (i) Using l'Hôpital's Rule, evaluate the limit

$$\lim_{x \rightarrow 0} \frac{\sinh^2 x}{x}.$$

- (ii) Differentiate:

(a) $y = x^{\tan x}$,

(b) $y = (\sin^{-1} x)^2$.

- (iii) Find the interval of convergence of

$$\sum_{n=1}^{\infty} \frac{(1+x^4)^n}{2^n n^4}.$$

Investigate the endpoints of the interval.

PLEASE TURN OVER

3. Evaluate the following integrals :

(i)
$$\int e^x \cosh x \, dx ;$$

(ii)
$$\int \frac{2 \ln x}{x} e^{(\ln x)^2} \, dx ;$$

(iii)
$$\int_{-\pi}^{\pi} x \sin^4 x \, dx ;$$

(iv)
$$\int_0^{\pi/2} (\cos x + \sin x)^2 \, dx ;$$

(v)
$$\int_1^{\infty} \frac{\ln x}{x^2} \, dx .$$

4. Find the general solution of the following differential equations :

(i)
$$\frac{dy}{dx} = \frac{\tan x}{(1+y)^3} ;$$

(ii)
$$(1-x^2) \frac{dy}{dx} + (1+x)y = (1-x) ;$$

(iii)
$$\frac{dy}{dx} = \frac{x^4 + x^3y - y^4}{x^4} .$$

PLEASE TURN OVER

5. Find the general solution of the following differential equations:

(i)
$$\frac{d^2y}{dx^2} - 6 \frac{dy}{dx} + 5y = 3 \cos x ;$$

(ii)
$$\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 5y = 5x^2 + 2x ;$$

(iii)
$$\frac{d^2y}{dx^2} - 10 \frac{dy}{dx} + 25y = e^{5x} .$$

For (iii) find also the solution subject to the conditions

$$y = 1 \quad \text{and} \quad \frac{dy}{dx} = 0 \quad \text{at} \quad x = 0 .$$

PLEASE TURN OVER

SECTION B

6. (i) Find the four stationary points of the function

$$f(x, y) = 2x^3 + 6xy^2 - 3y^3 - 150x$$

and determine their nature.

- (ii) Show that the differential equation

$$xy^2 + y + (x^2y + x) \frac{dy}{dx} = 0$$

is exact, and solve it.

7. Define $f(x)$ to be the periodic function with period 2π such that

$$f(x) = \frac{x}{\pi} \quad \text{for} \quad -\pi \leq x < \pi.$$

Find the Fourier series of $f(x)$.

By substituting a suitable value of x , deduce that

$$1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}.$$

Use Parseval's formula to deduce that

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}.$$

PLEASE TURN OVER

8. The Laplace transform of a function $f(t)$ is defined by

$$\mathcal{L}(f(t)) = F(s) = \int_0^{\infty} e^{-st} f(t) dt .$$

For $a > 0$, the Heaviside step function $H_a(t)$ is defined by

$$H_a(t) = \begin{cases} 1 & \text{if } t > a , \\ 0 & \text{if } t \leq a . \end{cases}$$

(i) Show that $\mathcal{L}(H_a(t)) = \frac{e^{-as}}{s}$ ($s > 0$) .

(ii) Prove the shift rule

$$\mathcal{L}(H_a(t) f(t-a)) = e^{-as} \mathcal{L}(f(t)) .$$

(iii) Find the function $y = y(t)$ satisfying the differential equation

$$\frac{d^2 y}{dt^2} + \frac{dy}{dt} - 2y = 1 - H_1(t)$$

with $y(0) = y'(0) = 0$.

You may assume that

$$\mathcal{L}(f'(t)) = -f(0) + s\mathcal{L}(f(t))$$

and

$$\mathcal{L}(f''(t)) = -f'(0) - sf(0) + s^2\mathcal{L}(f(t)) .$$

PLEASE TURN OVER

9. (i) For which values of λ does the following system of linear equations have no solutions :

$$\begin{aligned}x + y + z + t &= \lambda, \\x - y + z - t &= 0, \\x - 3y + z + \lambda t &= 2.\end{aligned}$$

- (ii) Let

$$A = \begin{pmatrix} 5 & -2 \\ 12 & -5 \end{pmatrix}.$$

Find a 2×2 matrix P such that $P^{-1}AP$ is diagonal.

Find A^{101} .

END OF PAPER

MATHEMATICAL FORMULAE

1. VECTOR ALGEBRA.

$$\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k} = (a_1, a_2, a_3)$$

Scalar (dot) product:

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

Vector (cross) product:

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

Scalar triple product:

$$[\mathbf{a}, \mathbf{b}, \mathbf{c}] = \mathbf{a} \cdot \mathbf{b} \times \mathbf{c} = \mathbf{b} \cdot \mathbf{c} \times \mathbf{a} = \mathbf{c} \cdot \mathbf{a} \times \mathbf{b} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$$

Vector triple product:

$$\mathbf{a} \times (\mathbf{b} \times \mathbf{c}) = (\mathbf{c} \cdot \mathbf{a})\mathbf{b} - (\mathbf{b} \cdot \mathbf{a})\mathbf{c}$$

2. SERIES

$$(1+x)^\alpha = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!}x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}x^3 + \dots \quad (\alpha \text{ arbitrary, } |x| < 1)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} + \dots,$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + (-1)^n \frac{x^{2n}}{(2n)!} + \dots,$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \dots,$$

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots + (-1)^n \frac{x^{n+1}}{(n+1)!} + \dots \quad (-1 < x \leq 1)$$

3. TRIGONOMETRIC IDENTITIES AND HYPERBOLIC FUNCTIONS

$$\sin(a+b) = \sin a \cos b + \cos a \sin b;$$

$$\cos(a+b) = \cos a \cos b - \sin a \sin b.$$

$$\cos iz = \cosh z; \quad \cosh iz = \cos z; \quad \sin iz = i \sinh z; \quad \sinh iz = i \sin z.$$

4. DIFFERENTIAL CALCULUS

(a) Leibniz's formula:

$$D^n(fg) = f D^n g + \binom{n}{1} Df D^{n-1} g + \dots + \binom{n}{n} D^n f D^{n-n} g + \dots + D^n f g.$$

(b) Taylor's expansion of $f(x)$ about $x = a$:

$$f(a+h) = f(a) + hf'(a) + \frac{h^2}{2!}f''(a) + \dots + \frac{h^n}{n!}f^{(n)}(a) + \epsilon_n(h),$$

$$\text{where } \epsilon_n(h) = h^{n+1}f^{(n+1)}(a + \theta h)/(n+1)!, \quad 0 < \theta < 1.$$

(c) Taylor's expansion of $f(x, y)$ about (a, b) :

$$f(a+h, b+k) = f(a, b) + [hf_x + kf_y]_{a,b} + \frac{1}{2!} [h^2 f_{xx} + 2hk f_{xy} + k^2 f_{yy}]_{a,b} + \dots$$

(d) Partial differentiation of $f(x, y)$:

$$\text{i. If } y = y(x), \text{ then } f = F(x), \text{ and } \frac{dF}{dx} = \frac{\partial f}{\partial x} + \frac{\partial f}{\partial y} \frac{dy}{dx}.$$

$$\text{ii. If } x = x(t), y = y(t), \text{ then } f = F(t), \text{ and } \frac{dF}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}.$$

$$\text{iii. If } x = x(u, v), y = y(u, v), \text{ then } f = F(u, v), \text{ and}$$

$$\frac{\partial F}{\partial u} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial u}, \quad \frac{\partial F}{\partial v} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial v}.$$

(e) Stationary points of $f(x, y)$ occur where $f_x = 0, f_y = 0$ simultaneously.

Let (a, b) be a stationary point: examine $D = [f_{xx}f_{yy} - (f_{xy})^2]_{a,b}$.

If $D > 0$ and $f_{xx}(a, b) < 0$, then (a, b) is a maximum;

If $D > 0$ and $f_{xx}(a, b) > 0$, then (a, b) is a minimum;

If $D < 0$ then (a, b) is a saddle-point.

(f) Differential equations:

i. The first order linear equation $dy/dx + P(x)y = Q(x)$ has an integrating

factor $I(x) = \exp[\int P(x)(dx)]$, so that $\frac{d}{dx}(Iy) = IQ$.

ii. $P(x, y)dx + Q(x, y)dy = 0$ is exact if $\partial Q/\partial x = \partial P/\partial y$.

5. INTEGRAL CALCULUS

(a) An important substitution: $\tan(\theta/2) = t$:
 $\sin \theta = 2t/(1+t^2)$, $\cos \theta = (1-t^2)/(1+t^2)$, $d\theta = 2dt/(1+t^2)$.

(b) Some indefinite integrals:

$$\int (a^2 - x^2)^{-1/2} dx = \sin^{-1} \left(\frac{x}{a} \right), \quad |x| < a.$$

$$\int (a^2 + x^2)^{-1/2} dx = \sinh^{-1} \left(\frac{x}{a} \right) = \ln \left\{ \frac{x}{a} + \left(1 + \frac{x^2}{a^2} \right)^{1/2} \right\}.$$

$$\int (x^2 - a^2)^{-1/2} dx = \cosh^{-1} \left(\frac{x}{a} \right) = \ln \left| \frac{x}{a} + \left(\frac{x^2}{a^2} - 1 \right)^{1/2} \right|.$$

$$\int (a^2 + x^2)^{-1} dx = \left(\frac{1}{a} \right) \tan^{-1} \left(\frac{x}{a} \right).$$

6. NUMERICAL METHODS

(a) Approximate solution of an algebraic equation:

If a root of $f(x) = 0$ occurs near $x = a$, take $x_0 = a$ and
 $x_{n+1} = x_n - [f(x_n)/f'(x_n)]$, $n = 0, 1, 2 \dots$

(Newton Raphson method).

(b) Formulae for numerical integration: Write $x_n = x_0 + nh$, $y_n = y(x_n)$.

i. Trapezium rule (1-strip): $\int_{x_0}^{x_1} y(x) dx \approx (h/2) [y_0 + y_1]$.

ii. Simpson's rule (2-strip): $\int_{x_0}^{x_2} y(x) dx \approx (h/3) [y_0 + 4y_1 + y_2]$.

(c) Richardson's extrapolation method: Let $I = \int_a^b f(x) dx$ and let I_1, I_2 be two

estimates of I obtained by using Simpson's rule with intervals h and $h/2$.

Then, provided h is small enough,

$$I_2 + (I_2 - I_1)/15,$$

is a better estimate of I .

7. LAPLACE TRANSFORMS

Function	Transform	Function	Transform
$f(t)$	$F(s) = \int_0^\infty e^{-st} f(t) dt$	$af(t) + bg(t)$	$aF(s) + bG(s)$
df/dt	$sF(s) - f(0)$	$d^2 f/dt^2$	$s^2 F(s) - sf(0) - f'(0)$
$e^{at} f(t)$	$F(s-a)$	$tf(t)$	$-dF(s)/ds$
$(\partial/\partial \alpha) f(t, \alpha)$	$(\partial/\partial \alpha) F(s, \alpha)$	$\int_0^t f(t) dt$	$F(s)/s$
$\int_0^t f(u) g(t-u) du$	$F(s)G(s)$		
1	$1/s$	$t^n (n = 1, 2, \dots)$	$n!/s^{n+1}, (s > 0)$
e^{at}	$1/(s-a), (s > a)$	$\sin \omega t$	$\omega/(s^2 + \omega^2), (s > 0)$
$\cos \omega t$	$s/(s^2 + \omega^2), (s > 0)$	$H(t-T) = \begin{cases} 0, & t < T \\ 1, & t > T \end{cases}$	$e^{-sT}/s, (s, T > 0)$

8. FOURIER SERIES

If $f(x)$ is periodic of period $2L$, then $f(x+2L) = f(x)$, and

$$f(x) = \frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi x}{L} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi x}{L}, \text{ where}$$

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx, \quad n = 0, 1, 2, \dots, \text{ and}$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx, \quad n = 1, 2, 3, \dots$$

Parseval's theorem

$$\frac{1}{L} \int_{-L}^L [f(x)]^2 dx = \frac{a_0^2}{2} + \sum_{n=1}^{\infty} (a_n^2 + b_n^2).$$

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1.6
Question 1		Marks & seen/unseen
Parts	<p>(i)</p> <p>(a) $i^{12} = (i^2)^6 = (-1)^6 = 1$</p> <p>(b) $e^{-i\pi/6} = \cos(-\frac{\pi}{6}) + i\sin(-\frac{\pi}{6}) = \frac{\sqrt{3}}{2} - \frac{i}{2}$</p> <p>(c) $1^{1/3} = (e^{i2\pi n})^{1/3} = 1 \quad (n=0)$ $= e^{i2\pi/3} = -\frac{1}{2} + \frac{\sqrt{3}}{2}i \quad (n=1)$ $= e^{i4\pi/3} = -\frac{1}{2} - \frac{\sqrt{3}}{2}i \quad (n=2)$</p> <p>(ii) $\cos^2 z = 4 \Rightarrow \cos z = \pm 2$ $\Rightarrow \frac{e^{iz} + e^{-iz}}{2} = \pm 2 \Rightarrow e^{iz} \pm 4 + e^{-iz} = 0$ $\Rightarrow e^{2iz} \pm 4e^{iz} + 1 = 0$ Let $u = e^{iz}$ $\Rightarrow u^2 \pm 4u + 1 = 0$ $\Rightarrow u = \frac{\pm 4 \pm \sqrt{16-4}}{2} = \pm 2 \pm \sqrt{3}$ $\Rightarrow e^{iz} = \pm 2 \pm \sqrt{3}$ $e^{iz} = (2 \pm \sqrt{3})e^{i2n\pi}$ or $e^{iz} = -2 \pm \sqrt{3} = (2 \pm \sqrt{3})e^{i\pi} e^{i2n\pi}$ $\Rightarrow iz = \ln(2 \pm \sqrt{3}) + i2n\pi$ $\Rightarrow iz = \ln(2 \pm \sqrt{3}) + i\pi + i2n\pi$ $\Rightarrow z = -i \ln(2 \pm \sqrt{3}) + 2n\pi$ $\Rightarrow z = -i \ln(2 \pm \sqrt{3}) + (2n+1)\pi$</p>	<p>2</p> <p>2</p> <p>4</p> <p>12</p>
	Setter's initials Myth	Checker's initials mmr
		Page number 1

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course <u>TFE 1.6</u>
Question <u>2</u>		Marks & seen/unseen
Parts	<p>(i) $\lim_{x \rightarrow 0} \frac{\sinh^3 x}{x} = \lim_{x \rightarrow 0} \frac{2 \sinh x \cosh x}{1} = 0$</p> <p>(ii) (a) $y = x^{\tan x}$ $\Rightarrow \ln y = \tan x \ln x$ $\Rightarrow \frac{1}{y} \frac{dy}{dx} = \sec^2 x \ln x + \frac{\tan x}{x} \Rightarrow \frac{dy}{dx} = x^{\tan x} \left(\sec^2 x \ln x + \frac{\tan x}{x} \right)$</p> <p>(b) $y = (\sin^{-1} x)^2$ $u = \sin^{-1} x \Rightarrow y = u^2$ $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = 2u \frac{d}{dx}(\sin^{-1} x)$ $\sin u = x \Rightarrow \cos u \frac{du}{dx} = 1 \Rightarrow \frac{du}{dx} = \frac{1}{\sqrt{1-\sin^2 u}} = \frac{1}{\sqrt{1-x^2}}$ $\Rightarrow \frac{dy}{dx} = \frac{2 \sin^{-1} x}{\sqrt{1-x^2}}$</p> <p>(iii) $\rho = \lim_{n \rightarrow \infty} \left \frac{\rho_{n+1}}{\rho_n} \right = \lim_{n \rightarrow \infty} \left \frac{(1+x^4)^{n+1}}{2^{n+1}(n+1)^4} \frac{2^n n^4}{(1+x^4)^n} \right$ $= \lim_{n \rightarrow \infty} \left \frac{1+x^4}{2} \frac{n^4}{(n+1)^4} \right = \lim_{n \rightarrow \infty} \left \frac{1+x^4}{2} \frac{1}{(1+\frac{1}{n})^4} \right = \left \frac{1+x^4}{2} \right$ Converges if $\left \frac{1+x^4}{2} \right < 1 \Rightarrow -1 < x < 1$</p>	<p>3</p> <p>4</p> <p>5</p> <p>6</p>
	Setter's initials <u>ryls</u>	Checker's initials <u>mlr</u>
		Page number <u>1</u>

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1.6
Question 2		Marks & seen/unseen
Parts (iii) (cont.)	<p>At endpoints $x = \pm 1$</p> <p>Series becomes $\sum_{n=1}^{\infty} \frac{1}{n^4}$ which is convergent</p>	2
	<p>Setter's initials MJH</p> <p>Checker's initials mm</p>	Page number 2

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1-6
Question 3		Marks & seen/unseen
Parts	<p>(i) $\int e^x \cosh x \, dx = \frac{1}{2} \int e^x (e^x + e^{-x}) \, dx = \frac{1}{2} \int (e^{2x} + 1) \, dx$ $= \frac{1}{4} e^{2x} + \frac{1}{2} x + c$</p> <p>(ii) $\int \frac{2 \ln x}{x} e^{(\ln x)^2} \, dx$ $u = (\ln x)^2 \Rightarrow \frac{du}{dx} = \frac{2 \ln x}{x}$ $= \int e^u \, du = e^u + c = e^{(\ln x)^2} + c$</p> <p>(iii) $\int_{-\pi}^{\pi} x \sin^4 x \, dx = 0$ (symmetry)</p> <p>(iv) $\int_0^{\pi/2} (\cos x + \sin x)^2 \, dx = \int_0^{\pi/2} (\cos^2 x + \sin^2 x + 2 \cos x \sin x) \, dx$ $= \int_0^{\pi/2} (1 + \sin 2x) \, dx = \left[x - \frac{1}{2} \cos 2x \right]_0^{\pi/2} = \frac{\pi}{2} + \frac{1}{2} + \frac{1}{2}$ $= 1 + \frac{\pi}{2}$</p> <p>(v) $\int_1^{\infty} \frac{\ln x}{x^2} \, dx = \lim_{a \rightarrow \infty} \int_1^a \frac{\ln x}{x^2} \, dx$ $u = \ln x \quad \frac{du}{dx} = \frac{1}{x}$ $\frac{dv}{dx} = \frac{1}{x^2} \quad v = -\frac{1}{x}$ $= \lim_{a \rightarrow \infty} \left\{ \left[-\frac{\ln x}{x} \right]_1^a + \int_1^a \frac{1}{x^2} \, dx \right\} = \lim_{a \rightarrow \infty} \left\{ -\frac{\ln a}{a} - \left[\frac{1}{x} \right]_1^a \right\}$ $= \lim_{a \rightarrow \infty} \left\{ -\frac{\ln a}{a} - \frac{1}{a} + 1 \right\} = 1$</p>	<p>3</p> <p>4</p> <p>2</p> <p>5</p> <p>6</p>
	Setter's initials <i>MJA</i> Checker's initials <i>mm</i>	Page number 1

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1.6
Question 4		Marks & seen/unseen
Parts	<p>(i) $\frac{dy}{dx} = \frac{\tan x}{(1+y)^3} \Rightarrow \int (1+y)^3 dy = \int \tan x dx$ $\Rightarrow \frac{1}{4}(1+y)^4 = -\ln(\cos x) + C$</p> <p>(ii) $(1-x^2)\frac{dy}{dx} + (1+x)y = (1-x)$ $\Rightarrow \frac{dy}{dx} + \frac{(1+x)}{(1+x)(1-x)} y = \frac{(1-x)}{(1+x)(1-x)}$ $\Rightarrow \frac{dy}{dx} + \frac{y}{1-x} = \frac{1}{1+x}$ Integrating factor $e^{\int \frac{dx}{1-x}} = e^{-\ln(x-1)} = \frac{1}{x-1}$ $\Rightarrow \frac{1}{x-1} \frac{dy}{dx} - \frac{y}{(x-1)^2} = \frac{1}{(x+1)(x-1)} \Rightarrow \frac{d}{dx} \left(\frac{y}{x-1} \right) = \frac{\frac{1}{2}}{x-1} - \frac{\frac{1}{2}}{x+1}$ $\Rightarrow \frac{y}{x-1} = \frac{1}{2} \ln(x-1) - \frac{1}{2} \ln(x+1) + C = \frac{1}{2} \ln \frac{x-1}{x+1} + C$ $\Rightarrow y = (x-1) \left[\frac{1}{2} \ln \frac{x-1}{x+1} + C \right]$</p> <p>(iii) $\frac{dy}{dx} = \frac{x^4 + x^3 y - y^4}{x^4}$ Let $y = vx \Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx} = 1 + v - v^4 \Rightarrow x \frac{dv}{dx} = 1 - v^4$ $\Rightarrow x \frac{dv}{dx} = (1+v^2)(1-v^2) \Rightarrow \frac{dv}{1-v^2} = \frac{dx}{x(1+v^2)}$ $\Rightarrow \int \frac{dv}{1-v^2} = \int \frac{dx}{x(1+v^2)} = \int dv \left\{ \frac{\frac{1}{2}}{1-v^2} + \frac{\frac{1}{2}}{1+v^2} \right\} = \int dv \left\{ \frac{\frac{1}{2}}{1+v^2} + \frac{\frac{1}{2}}{(1+v)(1-v)} \right\}$ $= \int dv \left\{ \frac{\frac{1}{2}}{1+v^2} + \frac{\frac{1}{4}}{1-v} + \frac{\frac{1}{4}}{1+v} \right\} = \frac{1}{2} \tan^{-1} v + \frac{1}{4} \ln(v+1) - \frac{1}{4} \ln(v-1) + C$</p>	<p>4</p> <p>7</p> <p>9</p>
	Setter's initials MJB	Checker's initials mu
		Page number 1

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1.6
Question 4		Marks & seen/unseen
Parts (iii) cont.	$\Rightarrow \ln x = \frac{1}{2} \tan^{-1} v + \frac{1}{4} \ln \frac{v+1}{v-1} + c$ $\Rightarrow x = e^{\frac{1}{2} \tan^{-1} v} \left(\frac{v+1}{v-1} \right)^{1/4} A$ $\Rightarrow x = A \left(\frac{y+x}{y-x} \right)^{1/4} e^{\frac{1}{2} \tan^{-1} \frac{y}{x}}$	
	Setter's initials M/G	Checker's initials Mm
		Page number 2

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course DE 1.6
Question 5		Marks & seen/unseen
Parts	<p>(i) $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 5y = 3\cos x$</p> <p>Auxiliary equation $\lambda^2 - 6\lambda + 5 = 0 \Rightarrow (\lambda - 5)(\lambda - 1) = 0 \Rightarrow \lambda = 5 \text{ or } 1$</p> <p>C.F. $y = Ae^{5x} + Be^x$</p> <p>P.I. $y = a\cos x + b\sin x$</p> <p>$\Rightarrow 5(a\cos x + b\sin x) - 6(-a\sin x + b\cos x) + (-a\cos x - b\sin x) = 3\cos x$</p> <p>$\Rightarrow 5a - 6b - a = 4a - 6b = 3$</p> <p>$5b + 6a - b = 4b + 6a = 0 \Rightarrow a = -\frac{2}{3}b$</p> <p>$\Rightarrow -\frac{8}{3}b - 6b = -\frac{26b}{3} = 3 \Rightarrow b = -\frac{9}{26} \quad a = \frac{3}{13}$</p> <p>$\Rightarrow y = Ae^{5x} + Be^x + \frac{3}{13}\cos x - \frac{9}{26}\sin x$</p> <p>(ii) $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 5y = 5x^2 + 2x$</p> <p>Auxiliary equation $\lambda^2 - 4\lambda + 5 = 0 \Rightarrow \lambda = \frac{4 \pm \sqrt{16 - 20}}{2} = 2 \pm i$</p> <p>C.F. $y = e^{2x}(A\cos x + B\sin x)$</p> <p>P.I. $y = ax^2 + bx + c$</p> <p>$\Rightarrow 5(ax^2 + bx + c) - 4(2ax + b) + 2a = 5x^2 + 2x$</p> <p>$\Rightarrow 5a = 5, 5b - 8a = 2, 5c - 4b + 2a = 0$</p> <p>$\Rightarrow a = 1, b = 2, c = \frac{6}{5}$</p> <p>$\Rightarrow y = e^{2x}(A\cos x + B\sin x) + x^2 + 2x + \frac{6}{5}$</p> <p>(iii) $\frac{d^2y}{dx^2} - 10\frac{dy}{dx} + 25y = e^{5x}$</p> <p>Auxiliary equation $\lambda^2 - 10\lambda + 25 = 0 \Rightarrow (\lambda - 5)(\lambda - 5) = 0 \Rightarrow \lambda = 5 \text{ (repeated)}$</p>	<p>6</p> <p>6</p>
	Setter's initials myt Checker's initials mmr	Page number 1

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1																				
Question 6 (solution)		Marks & seen/unseen																				
Parts	<p>a) $f_x = 6x^2 + 6y^2 - 150, f_y = 12xy - 9y^2$ $= 3y(4x - 3y).$</p> <p>For stationary point,</p> $f_x = 0 \Rightarrow x^2 + y^2 = 25$ $f_y = 0 \Rightarrow y = 0, x = \pm 5$ $\text{or } y = \frac{4x}{3}, \quad x^2 + \frac{16x^2}{9} = 25$ $\rightarrow x^2 \left(\frac{25}{9}\right) = 25 \rightarrow x^2 = 9$ $\rightarrow x = \pm 3, y = \pm \frac{4x}{3} = \pm 4.$ <p>So four stationary points are</p> $(5, 0), (-5, 0), (3, 4), (-3, -4).$ <p>$f_{xx} = 12x, f_{xy} = 12y, f_{yy} = 12x - 18y$</p> <p>So :</p> <table><thead><tr><th>Point</th><th>$f_{xx}f_{yy} - f_{xy}^2$</th><th>f_{xx}</th><th>Nature</th></tr></thead><tbody><tr><td>(5, 0)</td><td>$60 \cdot 60 > 0$</td><td>> 0</td><td><u>minimum</u></td></tr><tr><td>(-5, 0)</td><td>$(-60) \cdot (-60) > 0$</td><td>< 0</td><td><u>maximum</u></td></tr><tr><td>(3, 4)</td><td>$36 \cdot (-36) - 48^2 < 0$</td><td></td><td><u>saddle point</u></td></tr><tr><td>(-3, -4)</td><td>$(-36) \cdot (36) - 48^2 < 0$</td><td></td><td><u>saddle point</u></td></tr></tbody></table>	Point	$f_{xx}f_{yy} - f_{xy}^2$	f_{xx}	Nature	(5, 0)	$60 \cdot 60 > 0$	> 0	<u>minimum</u>	(-5, 0)	$(-60) \cdot (-60) > 0$	< 0	<u>maximum</u>	(3, 4)	$36 \cdot (-36) - 48^2 < 0$		<u>saddle point</u>	(-3, -4)	$(-36) \cdot (36) - 48^2 < 0$		<u>saddle point</u>	Similar seen 6 6
Point	$f_{xx}f_{yy} - f_{xy}^2$	f_{xx}	Nature																			
(5, 0)	$60 \cdot 60 > 0$	> 0	<u>minimum</u>																			
(-5, 0)	$(-60) \cdot (-60) > 0$	< 0	<u>maximum</u>																			
(3, 4)	$36 \cdot (-36) - 48^2 < 0$		<u>saddle point</u>																			
(-3, -4)	$(-36) \cdot (36) - 48^2 < 0$		<u>saddle point</u>																			
	Setter's initials ML	Checker's initials Mij																				
		Page number 2																				

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1
Question 6 <i>solve</i>		Marks & seen/unseen
Parts	<p>b) Eqn. is $P + Q \frac{dy}{dx} = 0$</p> <p>where $P = xy^2 + y$, $Q = x^2y + x$.</p> <p>As $P_y = Q_x = 2xy + 1$, it is <u>exact</u>.</p> <p>To solve, look for $u(x, y)$ such that</p> $u_x = P = xy^2 + y \quad (1)$ $u_y = Q = x^2y + x \quad (2)$ <p>From (1), $u = \frac{x^2y^2}{2} + xy + f(y)$</p> <p>From (2), $u_y = x^2y + x + f'(y) \Rightarrow f'(y) = 0$.</p> <p>So solution is</p> <p>Take $f(y) = 0$. So solution is $u(x, y) = c$, constant, i.e.</p> $x^2y^2 + 2xy = c.$	<p>2</p> <p>Similar seen.</p> <p>6</p>
	Setter's initials MLL	Checker's initials MJB
		Page number 3

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1
Question 7. (solution)		Marks & seen/unseen
Parts	<p>As $f(x)$ is an odd function, it has a Fourier sine series $\sum_{n=1}^{\infty} b_n \sin nx$, where</p> $b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} \frac{x}{\pi} \sin nx \, dx$ $= \frac{2}{\pi^2} \int_0^{\pi} x \sin nx \, dx$ <p>Integrate by parts:</p> $b_n = \frac{2}{\pi^2} \left(\left[-x \frac{\cos nx}{n} \right]_0^{\pi} + \int_0^{\pi} \frac{\cos nx}{n} dx \right)$ $= \frac{2}{\pi^2} \left(-\pi \frac{\cos n\pi}{n} + 0 \right)$ $= \frac{2 \cdot (-1)^{n+1}}{n\pi}$ <p>So Fourier series is</p> $\frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} \sin nx = \frac{2}{\pi} \left(\sin x - \frac{\sin 2x}{2} + \frac{\sin 3x}{3} - \dots \right)$ <p>Substitute $x = \frac{\pi}{2}$: gives</p> $f\left(\frac{\pi}{2}\right) = \frac{1}{2} = \frac{2}{\pi} \left(1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots \right)$ <p>Hence</p> $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}$	<p>Similar seen.</p> <p>9</p> <p>5</p>
	Setter's initials MLN <div>Checker's initials MyH</div>	Page number 5

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1
Question 7 (solution)		Marks & seen/unseen
Parts	<p>Parseval's formula says</p> $\frac{1}{\pi} \int_{-\pi}^{\pi} f(x)^2 dx = \sum b_n^2$ <p>i.e. $\frac{1}{\pi} \int_{-\pi}^{\pi} \frac{x^2}{n^2} dx = \frac{4}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2}$</p> <p>LHS is $\frac{1}{\pi^3} \left(\frac{2\pi^3}{3} \right) = \frac{2}{3}$. So</p> $\frac{2}{3} = \frac{4}{\pi^2} \sum \frac{1}{n^2}$ $\therefore \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$	6
	<p>Setter's initials MLW</p> <p>Checker's initials Mjt</p>	Page number 6

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1
Question 8 <i>Solve</i>		Marks & seen/unseen
Parts	<p>a) $\mathcal{L}(H_a(t)) = \int_a^\infty e^{-st} dt = \left[-\frac{e^{-st}}{s} \right]_a^\infty = \frac{e^{-as}}{s}$ ($s > 0$).</p> <p>b) $\mathcal{L}(H_a(t)f(t-a)) = \int_a^\infty e^{-st} f(t-a) dt$ Put $u = t-a$: $= \int_0^\infty e^{-s(u+a)} f(u) du$ $= e^{-as} \mathcal{L}(f(t)).$</p> <p>c) Take Laplace transforms: $s^2 \mathcal{L}(y) + s \mathcal{L}(y) - 2 \mathcal{L}(y) = \frac{1}{s} - \frac{e^{-s}}{s}$ \therefore $\mathcal{L}(y) = \frac{1 - e^{-s}}{s(s-1)(s+2)}$ By Partial Fractions, $\frac{1}{s(s-1)(s+2)} = \frac{1}{6} \left(-\frac{3}{s} + \frac{2}{s-1} + \frac{1}{s+2} \right)$ \therefore $\mathcal{L}(y) = \frac{1 - e^{-s}}{6} \left(-\frac{3}{s} + \frac{2}{s-1} + \frac{1}{s+2} \right)$ Insert using shift rule in (b): $y = \frac{1}{6} (-3 + 2e^t + e^{-2t})$ $- \frac{1}{6} H_1(t) (-3 + 2e^{t-1} + e^{-2(t-1)})$ </p>	<p>2 seen</p> <p>4 seen</p> <p>Similar seen ↓</p> <p>14</p>
	Setter's initials <i>mu</i>	Checker's initials <i>MJD</i>
		Page number 8

	EXAMINATION QUESTIONS/SOLUTIONS 2006-07	Course ISE 1
Question 9 marks		Marks & seen/unseen
Parts	<p>(a) Use Gaussian elimination:</p> $\begin{pmatrix} 1 & 1 & 1 & 1 & \lambda \\ 1 & -1 & 1 & -1 & 0 \\ 1 & -3 & 1 & \lambda & 2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 1 & 1 & 1 & \lambda \\ 0 & -2 & 0 & -2 & -\lambda \\ 0 & -4 & 0 & \lambda-1 & 2-\lambda \end{pmatrix}$ $\rightarrow \begin{pmatrix} 1 & 1 & 1 & 1 & \lambda \\ 0 & -2 & 0 & -2 & -\lambda \\ 0 & 0 & 0 & \lambda+3 & 2+\lambda \end{pmatrix}$ <p>This is echelon form. Last equation is</p> $(\lambda+3)t = 2+\lambda.$ <p>So no solutions if <u>$\lambda = -3$</u>.</p> <p>(b) Characteristic poly of A is $\begin{vmatrix} 5-\lambda & -2 \\ 12 & -5-\lambda \end{vmatrix}$</p> $= \lambda^2 - 1$ <p>So eigenvalues are 1, -1.</p> <p>Evecs: <u>$\lambda = 1$</u>: solve $\begin{pmatrix} 4 & -2 & 0 \\ 12 & -6 & 0 \end{pmatrix} \rightarrow$ evecs a $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$</p> <p><u>$\lambda = -1$</u>: solve $\begin{pmatrix} 6 & -2 & 0 \\ 12 & -4 & 0 \end{pmatrix} \rightarrow$ evecs a $\begin{pmatrix} 1 \\ 3 \end{pmatrix}$</p> <p>So take $P = \begin{pmatrix} 1 & 1 \\ 2 & 3 \end{pmatrix}$. Then</p> $P^{-1}AP = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} = D.$ <p>The matrix $A = PDP^{-1} \Rightarrow A^n = PD^nP^{-1}$.</p> <p>So $A^{101} = \cancel{PDP^{-1}} P D^{101} P^{-1} = PD^{-1}P^{-1} = A = \begin{pmatrix} 5 & -2 \\ 12 & -5 \end{pmatrix}.$</p>	<p>similar seen</p> <p>8</p> <p>similar seen ↓</p> <p>7</p> <p>5</p>
Setter's initials ML	Checker's initials Mya	Page number 10