

B.Sc. and M.Sci. EXAMINATIONS 2012

FIRST YEAR STUDENTS OF PHYSICS

MATHEMATICS - M.PHYS 1

Date Tuesday 1st May 2012 10.00 am - 1.00 pm

DO NOT OPEN THIS PAPER UNTIL THE INVIGILATOR TELLS YOU TO

Do not attempt more than SIX questions.

Please use a separate answerbook for each question.

A mathematical formulae sheet is provided.

[Before starting, please make sure that the paper is complete; there should be EIGHT pages, with a total of TEN questions. Ask the invigilator for a replacement if your copy is faulty.]

1. (i) Using L'Hôpital's rule, or otherwise, evaluate the limit

$$\lim_{x \rightarrow 0} \frac{(1 - \cos x)^{3/2}}{x - \sin x} .$$

- (ii) Determine the limit

$$\lim_{x \rightarrow \infty} x^{3/2} \left\{ (x+1)^{1/2} + (x-1)^{1/2} - 2x^{1/2} + x^{1/2} \sin^2 \frac{1}{x} \right\} .$$

- (iii) Evaluate the definite integral

$$I = \int_0^1 \frac{x^5}{x^2 + 1} dx .$$

- (iv) Evaluate the definite integral

$$J = \int_0^\infty e^{-x} \cos x dx .$$

2. (i) What is the area of the largest rectangle, with vertical sides, that can be inscribed in the ellipse

$$x^2 + \frac{y^2}{4} = 1 .$$

- (ii) Determine the volume, (total) surface area and centre of mass of a solid cone of height L and base radius R .

PLEASE TURN OVER

3. (i) If $u = x \ln(x^2 + y^2) - 2y \tan^{-1}(y/x)$, for what value of α does the function satisfy

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = u + \alpha x .$$

- (ii) If $u = \sin x \sin y \sin(x + y)$, show that

$$\frac{\partial u}{\partial x} = \sin y \sin(2x + y) ,$$

and similarly determine $\partial u / \partial y$.

Consider the function $u(x, y)$ in the square region defined by the lines

$$x = \frac{\pi}{4} , \quad x = \frac{\pi}{2} \quad \text{and} \quad y = \frac{\pi}{4} , \quad y = \frac{\pi}{2} .$$

Determine the location of the stationary points in this square and the associated value of u .

What is the nature of the stationary point ?

4. Consider the complex number $z = -1 + i$:

- (i) Write z in polar form.
- (ii) Plot z in the complex plane.
- (iii) Find z^9 and express the result in real and imaginary form.
- (iv) Find $z^{1/3} \equiv \sqrt[3]{z}$ and express the results in polar form.
- (v) Find the value of $\cos(z)$ expressed in real and imaginary form.

For the following complex functions determine if they are differentiable for all values $z \in \mathbb{C}$. If yes, find $f'(z)$.

(vi) $f(z) = \operatorname{Im}(z)$.

(vii) $f(z) = e^{-z}$.

5. Consider the three vectors $\mathbf{a} = \mathbf{i} - 2\mathbf{j} - 3\mathbf{k}$, $\mathbf{b} = \mathbf{i} - 4\mathbf{j} - 13\mathbf{k}$ and $\mathbf{c} = -3\mathbf{i} + 5\mathbf{j} + 4\mathbf{k}$ where \mathbf{i}, \mathbf{j} and \mathbf{k} denote the natural basis vectors for \mathbb{R}^3 .
- (i) Calculate the lengths $|\mathbf{a}|$, $|\mathbf{b}|$ of the vectors \mathbf{a} and \mathbf{b} .
 - (ii) Calculate the dot (scalar) product $\mathbf{a} \cdot \mathbf{b}$.
 - (iii) From the sign of the dot product $\mathbf{a} \cdot \mathbf{b}$, what can you conclude about the angle θ ($0 \leq \theta \leq \pi$) between the two vectors \mathbf{a} and \mathbf{b} ?
 - (iv) Find the angle θ ($0 \leq \theta \leq \pi$) between the two vectors \mathbf{a} and \mathbf{b} .
 - (v) Calculate the area of the parallelogram spanned by the two vectors \mathbf{a} and \mathbf{c} .

Consider three planes in \mathbb{R}^3 specified by the equations

$$x - 2y - 3z = 2, \tag{1a}$$

$$x - 4y - 13z = 14, \tag{1b}$$

$$-3x + 5y + 4z = 0. \tag{1c}$$

- (vi) Let \mathbf{A} denote the matrix of coefficients for this system of three linear equations in three unknowns. Calculate $\det \mathbf{A}$, the determinant of \mathbf{A} .
- (vii) Does the system of linear equations (1a), (1b) and (1c) have a unique solution?
- (viii) Using Gaussian elimination or otherwise, show that the solution $\mathbf{r} = \begin{pmatrix} x \\ y \\ z \end{pmatrix}$ to equations (1a), (1b) and (1c) can be written in the form

$$\mathbf{r} = \mathbf{r}_0 + \mu \mathbf{d}, \quad \mu \in \mathbb{R}$$

and identify possible values for \mathbf{r}_0 and \mathbf{d} .

PLEASE TURN OVER

6. Consider the 2×2 matrix

$$\mathbf{B} = \begin{pmatrix} 2 & -3 \\ 2 & -5 \end{pmatrix}.$$

- (i) Calculate $\det \mathbf{B}$, the determinant of the matrix \mathbf{B} .
- (ii) Find the two eigenvalues $\lambda_1 > \lambda_2$ of the matrix \mathbf{B} .
- (iii) Show that $\mathbf{x}_1 = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ is an eigenvector of the matrix \mathbf{B} with eigenvalue λ_1 .
- (iv) Find the eigenvector \mathbf{x}_2 associated with the smallest eigenvalue λ_2 .
- (v) Let $\mathbf{S} = (\mathbf{x}_1 \mathbf{x}_2)$ denote the 2×2 matrix of eigenvectors. Calculate $\mathbf{S}^{-1}\mathbf{B}\mathbf{S}$ explicitly.
- (vi) Why does the inverse \mathbf{B}^{-1} exist?
- (vii) What are the two eigenvalues of \mathbf{B}^{-1} ?
- (viii) What are the two eigenvectors of \mathbf{B}^{-1} ?

7. Consider an accelerator that is producing isotopes of type A at a constant rate k in a sample. The isotope of type A is radioactive with a decay rate α and decays into an isotope B which is stable.

- (i) Write down the differential equations governing the time-dependent populations $A(t)$ and $B(t)$.
- (ii) Write down the full name for this set of equations according to the classification of differential equations.
- (iii) Find the steady-state solution to $A(t)$, meaning the solution where $A(t)$ is constant.
- (iv) Find the general solution to $A(t)$ to show that you end up with

$$A(t) = A_0 e^{-\alpha t} + \frac{k}{\alpha}.$$

- (v) Draw a sketch of $A(t)$ for the two special cases where $A(t=0) = 0$, and where $A(t=0) > k/\alpha$.
- (vi) Confirm through simple insertion in the differential equation, or otherwise, that

$$B(t) = c_1 t + c_2 e^{-\alpha t} + c_3$$

is a solution to $B(t)$ and find the solution satisfying the initial state conditions $A(t) = B(t) = 0$ at $t = 0$.

8. (i) A function $f(x, y, z)$ is given by

$$f(x, y, z) = x^2 e^y + 3 \sin z .$$

- (a) At the point $(1, 0, \pi/4)$, find the direction in which f is increasing most rapidly.
- (b) Find two distinct directions that correspond to a minimum change in $|f|$ at this point.
- (c) Find the equation of the plane that is tangent to the surface of constant f at the point $(1, 0, \pi/4)$.

- (ii) Consider a variable transformation given by

$$\begin{aligned} u &= x + y , \\ v &= x - y , \\ w &= z^2 . \end{aligned}$$

- (a) Write down the position vector $\mathbf{r} = x \hat{\mathbf{i}} + y \hat{\mathbf{j}} + z \hat{\mathbf{k}}$ in terms of u, v and w . Show that the line element $d\mathbf{r}$ in (u, v, w) is given by

$$d\mathbf{r} = \frac{\sqrt{2}}{2} du \left[\frac{\sqrt{2}}{2} (\hat{\mathbf{i}} + \hat{\mathbf{j}}) \right] + \frac{\sqrt{2}}{2} dv \left[\frac{\sqrt{2}}{2} (\hat{\mathbf{i}} - \hat{\mathbf{j}}) \right] + \frac{1}{2\sqrt{w}} dw [\hat{\mathbf{k}}]$$

and verify that the corresponding unit vectors $\hat{\mathbf{u}}, \hat{\mathbf{v}}$ and $\hat{\mathbf{w}}$ (shown within the square braces above) are mutually orthogonal.

- (b) A function g is expressed in (u, v, w) coordinates. By considering the expression $\nabla g \cdot d\mathbf{r}$ or otherwise, show that the $\hat{\mathbf{u}}$ component of ∇g is given by

$$(\nabla g)_u = \sqrt{2} \frac{\partial g}{\partial u} .$$

- (c) Find the corresponding $\hat{\mathbf{v}}$ and $\hat{\mathbf{w}}$ component of ∇g .

PLEASE TURN OVER

9. (i) A vector field $\mathbf{F} = x\hat{\mathbf{i}} + y\hat{\mathbf{j}} + z\hat{\mathbf{k}}$ is found within a closed volume V that is bounded below by the $x - y$ plane and above by the surface S satisfying $x^2 + y^2 + \beta^2 z^2 = a^2$ where β and a are constants.

- (a) Show that

$$\iiint_V \nabla \cdot \mathbf{F} \, dV = 2\pi \frac{a^3}{\beta}.$$

where the lefthand side is the integral of $\nabla \cdot \mathbf{F}$ over V

You might find it convenient to transform to new variables

$$(u, v, w) = (x, y, \beta z).$$

- (b) Calculate the flux of \mathbf{F} entering through the bottom face ($z = 0$) of V .
- (c) Hence write down, with suitable explanation, the flux of \mathbf{F} leaving V through the top surface S .

- (ii) (a) Show that $\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot \nabla \times \mathbf{A} - \mathbf{A} \cdot \nabla \times \mathbf{B}$.

- (b) Hence show that

$$\iiint_V \nabla \times \mathbf{F} \, dV = - \oint_S \mathbf{F} \times d\mathbf{S},$$

where S is a closed surface bounding the volume V .

Hint: Consider the vector field $\mathbf{G} \equiv \mathbf{c} \times \mathbf{F}$ where \mathbf{c} is a constant vector and apply the Divergence Theorem.

10. (i) State Stokes' Theorem, indicating clearly the meaning of all the symbols. Illustrate your answer with a sketch showing the direction and orientation of all surfaces and integrals.

- (ii) A vector field \mathbf{B} is given in cylindrical polar coordinates by $\mathbf{B} = \rho z \hat{\phi}$. Define the surface S by that portion of a cylinder, centred on the z -axis, that lies in the first quadrant ($0 < \phi < \pi/2$) between the planes $z = 0$ and $z = b$. Let the normal to this surface be defined to point away from the z -axis.

(a) Calculate $\iint_S \nabla \times \mathbf{B} \cdot d\mathbf{S}$ over this surface.

(b) Verify Stokes' Theorem by calculating a suitable line integral.

You may assume the form of the curl operator in cylindrical polar coordinates :

$$\nabla \times \mathbf{A} = \left(\frac{1}{\rho} \frac{\partial A_z}{\partial \phi} - \frac{\partial A_\phi}{\partial z} \right) \hat{\rho} + \left(\frac{\partial A_\rho}{\partial z} - \frac{\partial A_z}{\partial \rho} \right) \hat{\phi} + \left(\frac{1}{\rho} \frac{\partial}{\partial \rho} (\rho A_\phi) - \frac{1}{\rho} \frac{\partial A_\rho}{\partial \phi} \right) \hat{\mathbf{k}}.$$

END OF PAPER