

MEng (Engineering) Examination 2016

Year 1

AE1-107 Mathematics Term II

Tuesday 24th May 2016: 14.00 to 16.00
[2 hours]

The paper is divided into Section A and Section B

Both sections carry the same weight

Candidates may obtain full marks for complete answers to **ALL** questions.

You must answer each section in a separate answer booklet

The use of lecture notes is NOT allowed.

Section A

1. Consider the ODE:

$$\frac{dy}{dx} = \frac{xe^{x^2}}{y}.$$

- (a) Find the general solution to the ODE. [15%]

Now consider the ODE:

$$y \frac{dy}{dx} = \frac{6xy^2 - y^3}{3xy - 6x^2}.$$

- (b) Show that the ODE is inexact. [10%]

- (c) Find an integrating factor that makes the ODE exact.

Hint: the required integrating factor is a function of y alone. [30%]

- (d) Find the general solution to the ODE. [25%]

Finally consider the ODE:

$$\frac{dy}{dx} + y = y^3.$$

- (e) Find the general solution to the ODE. [20%]

2. The Laplace transform $F(s)$ of a function $f(t)$ is defined as:

$$F(s) = \int_0^{\infty} f(t)e^{-st} dt$$

for some appropriate range of s .

- (a) Derive expressions for the Laplace transform of $\frac{df}{dt}$ and the Laplace transform of $\frac{d^2f}{dt^2}$ in terms of the Laplace transform of $f(t)$. Show all your workings. [25%]

Consider the ODE:

$$a \frac{d^2y}{dx^2} + b \frac{dy}{dx} + y = f(x).$$

- (b) If $a = 1$, $b = 2$, $f(x) = 0$, find the general solution to the ODE. [15%]
- (c) If $a = 1/4$, $b = 0$, $f(x) = \sin(2x)$, find the general solution to the ODE. [35%]
- (d) If $a = 2$, $b = 1$, $f(x) = x^2$, use Laplace transforms to find the Laplace transform of a particular solution to the ODE that satisfies $\frac{dy}{dx}(0) = 1$ and $y(0) = 0$. [25%]

Section B

3. Let

$$A(p) = \begin{bmatrix} 1 & -1 & 8 \\ 2 & 0 & 6 \\ -1 & 5 & p \\ 3 & 1 & 4 \end{bmatrix}, b(q) = \begin{bmatrix} 2 \\ -4 \\ q \\ -10 \end{bmatrix}, x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- (a) Show that if $p \neq -28$, the set of equations $A(p)x = b(q)$ has a unique solution for all q . Find this solution x in terms of p and q . [35%]
- (b) Show that $A(-28)x = b(q)$ has no solution if $q \neq -18$. Find all possible solutions x in this case when $q = -18$. [35%]

Let

$$B = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 2 & 1 & 4 \end{bmatrix}$$

- (c) Find the eigenvalues and eigenvectors of B . [30%]

4. Consider the system

$$\frac{dX(t)}{dt} = AX(t)$$

where

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

$$X(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}$$

and

$$X(0) = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

- (a) Find the eigenvalues and eigenvectors of A . [30%]
- (b) Solve the system to find $x_1(t)$ and $x_2(t)$. [40%]
- (c) Evaluate A^{50} . [30%]

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

①

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

1.) a.)

$$\frac{dy}{dx} = \frac{xe^{x^2}}{y}$$

$$\Rightarrow \int y dy = \int xe^{x^2} dx$$

$$\Rightarrow \frac{1}{2}y^2 = \frac{1}{2}e^{x^2} + C$$

$$\Rightarrow y^2 = e^{x^2} + D \quad \text{ok}$$

$$\Rightarrow y = \pm \sqrt{e^{x^2} + D}$$

3

1.) b.)

$$y \frac{dy}{dx} = \frac{6xy^2 - y^3}{3xg - 6x^2}$$

Peter Vincent

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

②

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow \frac{dy}{dx} = - \frac{f(x,y)}{g(x,y)} \quad \text{where}$$

$$f(x,y) = y^2 - 6xy$$

$$g(x,y) = 3xy - 6x^2$$

$$\Rightarrow \frac{\partial f}{\partial y} = 2y - 6x$$

$$\frac{\partial g}{\partial x} = 3y - 12x$$

Not equal! \Rightarrow Inexact

2

1.) c.) require $I(y)$ s.t.

Red *Yue*

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

③

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\frac{\partial}{\partial y} [I(y) (y^2 - 6xy)] =$$

$$\frac{\partial}{\partial x} [I(y) (3xy - 6x^2)]$$

$$\Rightarrow 2y I(y) + \frac{dI(y)}{dy} y^2 - 6xy \frac{dI(y)}{dy}$$

$$- I(y) 6x =$$

$$3I(y)y - 12I(y)x$$

$$\Rightarrow \frac{dI(y)}{dy} (y^2 - 6xy) =$$

$$I(y) (y - 6x)$$

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

④

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow \frac{1}{I(y)} \frac{dI(y)}{dy} = \frac{1}{y} \left(\frac{y - 6x}{y - 6x} \right) = \frac{1}{y}$$

$\Rightarrow I(y) = y$ is an
integrating factor.

6

1.) d.) $f = y(y^2 - 6xy)$
 $g = y(3xy - 6x^2)$

Need a $u(x, y)$ s.t.

$$\frac{\partial u}{\partial x} = f, \quad \frac{\partial u}{\partial y} = g$$

Per *Ym*

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

5

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow u = y^3 x - 3x^2 y^2 + C(y)$$

$$u = y^3 x - 3x^2 y^2 + D(x)$$

$$\Rightarrow C(y) = D(x) = 0$$

$$\Rightarrow u = y^3 x - 3x^2 y^2$$

$$\Rightarrow y^3 x - 3x^2 y^2 = F$$

5

1/e.) Let $y = u^{-\frac{1}{2}}$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = -\frac{1}{2} u^{-\frac{3}{2}} \frac{du}{dx}$$

Ben M

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

⑥

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow -\frac{1}{2}u^{-\frac{3}{2}}\frac{du}{dx} + u^{-\frac{1}{2}} = u^{-\frac{3}{2}}$$

$$\Rightarrow \frac{du}{dx} - 2u = -2$$

Linear

$$\text{I.s. } I = \exp\left|\int -2 dx\right| = e^{-2x}$$

$$u = -2e^{+2x} \int e^{-2x} dx$$

$$= -2e^{2x} \left[-\frac{1}{2}e^{-2x} + C \right]$$

$$= 1 - 2Ce^{2x}$$

Mark

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

⑦

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow y = \frac{1}{\sqrt{1 - 2Ce^{2x}}}$$

4

Peter Vincent

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

8

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$2.) a.) F(s) = \int_0^{\infty} e^{-st} f(t) dt$$

$$L\left\{\frac{df}{dt}\right\} = \int_0^{\infty} \frac{df}{dt} e^{-st} dt$$

$$= -f(0) + s \int_0^{\infty} f(t) e^{-st} dt$$

$$= -f(0) + s L\{f\}$$

$$\Rightarrow L\left\{\frac{df}{dt}\right\} = s L\{f\} - f(0)$$

$$\Rightarrow L\left\{\frac{d^2f}{dt^2}\right\} = s L\left\{\frac{df}{dt}\right\} - \frac{df}{dt}(0)$$

Peter Vincent

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

⑨

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow \mathcal{L}\left\{\frac{d^2 f}{dt^2}\right\} = s(s \mathcal{L}\{f\} - f(0)) - \frac{df}{dt}(0)$$

$$= s^2 \mathcal{L}\{f\} - s f(0) - \frac{df}{dt}(0)$$

5

2.) b) $\alpha^2 + 2\alpha + 1 = 0$

$$\Rightarrow \alpha = \frac{-2 \pm \sqrt{4 - 4}}{2}$$

$$= -1$$

$$\Rightarrow y = (A + Bx) e^{-x}$$

3

Alan Yoon

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

(10)

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$2x) \quad \frac{\alpha^2}{4} + 1 = 0$$

$$\Rightarrow \alpha = \pm 2i$$

$$\Rightarrow y_h = A \sin(2x) + B \cos(2x)$$

For y_p , try $y_p = x(\sin(2x) + x D \cos(2x))$

(since $C \sin(2x) + D \cos(2x) \equiv y_h$)

$$y_p' = 2x(\cos(2x) + C \sin(2x) - 2x D \sin(2x) + D \cos(2x))$$

Alan Jones

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent



Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$y_p'' = -4x(\sin(2x) + 4(\cos(2x) - 4x\mathcal{D}\cos(2x) - 4\mathcal{D}\sin(2x))$$

Sub in to ODE \Rightarrow

$$\begin{aligned} & -x(\sin(2x) + (\cos(2x) \\ & - x\mathcal{D}\cos(2x) - \mathcal{D}\sin(2x) \\ & + x(\sin(2x) + x\mathcal{D}\cos(2x)) \\ & = \sin(2x) \end{aligned}$$

$$\Rightarrow C=0, \quad \mathcal{D} = -1 \quad \Rightarrow$$

$$y_p = -x\cos(2x) \quad \Rightarrow$$

Peter Vincent

Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent

(12)

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$y = y_h + y_p =$$

$$A \sin(2x) + B \cos(2x)$$

$$- x \cos(2x)$$

7

2.) d.)

$$2L\left\{\frac{d^2y}{dx^2}\right\} + L\left\{\frac{dy}{dx}\right\} + L\{y\} = L\{x^2\}$$

$$\Rightarrow 2[s^2L\{y\} - sy(0) - y'(0)]$$

$$+ sL\{y\} - y(0) + L\{y\}$$

$$= \frac{2}{s^3}$$



Course Code and Title (Required): AE-107

Setter (Required): Peter Vincent



13

Write on this side only (in ink) between the margins, not more than one solution per sheet please. Solutions must be signed and dated by both exam setter and referee.

Marks

$$\Rightarrow 2s^2 L\{y\} - 2 + sL\{y\} + L\{y\} = \frac{2}{s^3}$$

$$\Rightarrow L\{y\} (2s^2 + s + 1) = \frac{2}{s^3} + 2$$

$$\Rightarrow L\{y\} = \frac{2 + 2s^3}{s^3(2s^2 + s + 1)}$$

5

Almo

Section B

1. Let

$$A(p) = \begin{bmatrix} 1 & -1 & 8 \\ 2 & 0 & 6 \\ -1 & 5 & p \\ 3 & 1 & 4 \end{bmatrix}, b(q) = \begin{bmatrix} 2 \\ -4 \\ q \\ -10 \end{bmatrix}, x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

- (a) Show that if $p \neq -28$, the set of equations $A(p)x = b(q)$ has a unique solution for all q . Find this solution x in terms of p and q

[35%]

$$\begin{bmatrix} 1 & -1 & 8 \\ 2 & 0 & 6 \\ -1 & 5 & p \\ 3 & 1 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ -4 \\ q \\ -10 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 8 & 2 \\ 0 & 2 & -10 & -8 \\ 0 & 4 & p+8 & q+2 \\ 0 & 4 & -20 & -16 \end{bmatrix}$$

$$\begin{bmatrix} 1 & -1 & 8 & 2 \\ 0 & 2 & -10 & -8 \\ 0 & 0 & p+28 & q+18 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$p \neq -28$ a unique solution

$$x_3 = \frac{q+18}{p+28}$$

$$x_2 = -4 + 5 \frac{q+18}{p+28} = \frac{-4p+5q-22}{p+28}$$

$$x_1 = 2 - 8 \frac{q+18}{p+28} + \frac{-4p+5q-22}{p+28} = \frac{-2p+3q-110}{p+28}$$

- (b) Show that $A(-28)x = b(q)$ has no solution if $q \neq -18$. Find all possible solutions x in this case when $q = -18$.

[35%]

$$p = -28, q \neq -18$$

$$\rightarrow 0 = q+18 \neq 0 \rightarrow \text{no solution}$$

$$p = -28, q = -18$$

$$x_3 = t \text{ arbitrary}$$

$$x_2 = -4 + 5t$$

$$x_1 = -2 - 3t$$

(c) Let

$$B = \begin{bmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 2 & 1 & 4 \end{bmatrix}$$

find eigenvalues and eigenvectors

[30%]

The characteristic equation is $(3 - \lambda)^2(4 - \lambda) = 0$ so there are two coincident eigenvalues $\lambda_1 = \lambda_2 = 3$ and an independent one $\lambda_3 = 4$. For $\lambda_3 = 4$ the system

$$B = \begin{bmatrix} 3-4 & 0 & 0 \\ 0 & 3-4 & 0 \\ 2 & 1 & 4-4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

becomes a system of 2 equations in 3 unknowns $x = y = 0$; so the eigenvectors of B for $\lambda_3 = 4$ are the eigenvectors

$$v_3 = \begin{bmatrix} 0 \\ 0 \\ x_3 \end{bmatrix}$$

For $\lambda_1 = \lambda_2 = 3$ the system becomes $2x + y + z = 0$; so the eigenvectors are

$$v = \begin{bmatrix} x_1 \\ x_2 \\ -2x_1 - x_2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix} x_1 + \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix} x_2$$

and this allows more than one solution linearly independent. For example for $x=1$ and $y=0$ or $x=0$ and $y=1$.

$$v_1 = \begin{bmatrix} 1 \\ 0 \\ -2 \end{bmatrix} \quad v_2 = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$$

or for $x=1$ and $y=-2$ or $x=0$ and $y=1$.

$$v_1 = \begin{bmatrix} 1 \\ -2 \\ 0 \end{bmatrix} \quad v_2 = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$$

so the eigenvectors matrix

$$V = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -2 & -1 & 1 \end{bmatrix}$$

2. Consider the system

$$\frac{dX(t)}{dt} = AX(t)$$

where

$$A = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}$$

$$X(t) = \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix}$$

and

$$X(0) = \begin{bmatrix} 1 \\ 3 \end{bmatrix}$$

(a) Find the eigenvalues and eigenvectors of A .

[30%]

Eigenvalues of A

$$(2 - \lambda)(2 - \lambda) - 1 = 0$$

$$\lambda_1 = 1, \lambda_2 = 3$$

Eigenvector of A for $\lambda_1 = 1$

$$A = \begin{bmatrix} 2-1 & 1 \\ 1 & 2-1 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$u_1 + u_2 = 0$$

$$U_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Eigenvector of A for $\lambda_2 = 3$

$$A = \begin{bmatrix} 2-3 & 1 \\ 1 & 2-3 \end{bmatrix} \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

$$u_1 - u_2 = 0$$

$$U_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$$

(b) Solve the system to find $x_1(t)$ and $x_2(t)$.

[40%]

So we can write

$$A = U\Lambda U^{-1}$$

where

$$U = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$$

so

$$\frac{dX}{dt} = AX$$

$$\frac{dX}{dt} = U\Lambda U^{-1}X$$

$$\frac{dU^{-1}X}{dt} = \Lambda U^{-1}X$$

$$\frac{dZ}{dt} = \Lambda Z, Z = U^{-1}X$$

so

$$\frac{dz_1}{dt} = \lambda_1 z_1(t) = 1z_1(t),$$

$$\frac{dz_2}{dt} = \lambda_2 z_2(t) = 3z_2(t)$$

and

$$z_1(t) = \alpha e^{\lambda_1 t} = \alpha e^{1t},$$

$$z_2(t) = \beta e^{\lambda_2 t} = \beta e^{3t}$$

$$X = UZ = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} \alpha e^{1t} \\ \beta e^{3t} \end{bmatrix}$$

$$x_1(t) = \alpha e^{1t} + \beta e^{3t},$$

$$x_2(t) = \alpha e^{1t} - \beta e^{3t},$$

since $x_1(0) = 1$ and $x_2(0) = 3$

$$1 = \alpha + \beta,$$

$$3 = \alpha - \beta,$$

$$\alpha = 1, \beta = -2$$

$$x_1(t) = 1e^{1t} - 2e^{3t},$$

$$x_2(t) = 1e^{1t} + 2e^{3t},$$

(c) Evaluate A^{50} .

[30%]

$$A = U\Lambda U^{-1}$$

$$A^{50} = (U\Lambda U^{-1})^{50}$$

$$A^{50} = U\Lambda^{50}U^{-1}$$

$$U = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

(18)
LAST
PAGE

$$U^{-1} = \begin{bmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{bmatrix}$$

$$\Lambda = \begin{bmatrix} 1 & 0 \\ 0 & 3 \end{bmatrix}$$

$$\Lambda^{50} = \begin{bmatrix} 1 & 0 \\ 0 & 3^{50} \end{bmatrix}$$

$$A^{50} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 3^{50} \end{bmatrix} \begin{bmatrix} 1/2 & 1/2 \\ 1/2 & -1/2 \end{bmatrix} = 10^{23} \begin{bmatrix} 3.5895 & -3.5895 \\ -3.5895 & 3.5895 \end{bmatrix}$$