

## DISCRETE MATHEMATICS AND COMPUTATIONAL COMPLEXITY

### 1. [Compulsory]

Let  $\mathcal{R}$  be the set of all relations on a set  $A$ .

- a) Express the predicate  $P(x)$ , meaning that  $x \in \mathcal{R}$  is a transitive relation, in terms of appropriate symbolic logic.

[ 2 ]

- b) Prove that the proposition  $p$  given by  $\forall R (P(R) \leftrightarrow (\forall n \in \mathbb{Z}^+ R^n \subseteq R))$  is true, where the universe of discourse is  $\mathcal{R}$ .

[ 12 ]

- c) Prove that the proposition  $q$  given by  $\forall R (P(R) \rightarrow (\forall n \in \mathbb{Z}^+ ((n \geq 2) \rightarrow P(R^n))))$  is true, where the universe of discourse is  $\mathcal{R}$ .

[ 10 ]

- d) Replacing the implication in the definition of  $q$  by its converse yields another proposition  $r$ . Prove that  $r$  is false.

[ 8 ]

- e) A relation  $R'$  is said to be the transitive closure of  $R$  when  $R'$  is the smallest transitive relation containing  $R$ . Define the connectivity relation  $R^*$  and prove that  $R^* = R'$ .

[ 8 ]

SPECIMEN EE3 QUESTION - MODEL ANSWER

1.  $P(x) \equiv \forall a \forall b \forall c ((a,b) \in x \wedge (b,c) \in x \rightarrow (a,c) \in x)$   
 where the universe of discourse is  $A$ .

2.  $\forall R [P(R) \leftrightarrow (\forall n \in \mathbb{Z}^+ R^n \subseteq R)]$

First prove  $\forall R [P(R) \rightarrow (\forall n \in \mathbb{Z}^+ R^n \subseteq R)]$

For  $n=1$ , we have  $\forall R [P(R) \rightarrow R \subseteq R]$   
 which is true as the RHS is always true.

Assume true for  $n$  & use induction for  $n > 1$ .

Let  $(a,b) \in R^{n+1} = R \cdot R^n$ .

$\Rightarrow \exists x [(a,x) \in R \wedge (x,b) \in R^n]$

Since  $R^n \subseteq R$ ,  $(x,b) \in R$

$\Rightarrow (a,b) \in R$ , i.e.  $R^{n+1} \subseteq R \quad \square$

second prove  $\forall R [(\forall n \in \mathbb{Z}^+ R^n \subseteq R) \rightarrow P(R)]$

To be false, we would need  $R$  s.t.

$(\forall n \in \mathbb{Z}^+ R^n \subseteq R) \wedge \neg P(R)$ .

$\forall n \in \mathbb{Z}^+ R^n \subseteq R \not\Rightarrow R^2 \subseteq R$

Let  $(a,b) \in R \wedge (b,c) \in R$ . Then  $(a,c) \in R^2 \subseteq R$

$\Rightarrow (a,c) \in R \Rightarrow P(R) \quad \square$

$$\forall R [R(R) \rightarrow (\forall n \in \mathbb{Z}^+ (n > 2) \rightarrow P(R^n))]$$

We will prove  $\forall R [P(R) \rightarrow \forall n \in \mathbb{Z}^+ P(R^n)]$ ,  
which is a stronger statement.

For  $n=1$ , implication is  $P(R) \rightarrow P(R)$ , which is true.  
Assume for  $n$  & use induction for  $n > 1$ .  
Specially show that  $P(R^{2n})$  and  $P(R^{2n+1})$  are true.

First,  $P(R^{2n})$ .

$$\text{Let } (a, b) \in R^{2n} = \mathbb{Z} R^n \cdot R^n \quad (b, c) \in R^{2n} = R^n \cdot R^n$$

$$\text{Then } \exists x \exists y [(a, x) \in R^n \wedge (x, b) \in R^n \wedge (b, y) \in R^n \wedge (y, c) \in R^n]$$

$$\text{since } R^n \text{ is transitive, } (a, b) \in R^n \wedge (b, c) \in R^n$$

$$\Rightarrow (a, c) \in R^n \cdot R^n = R^{2n}.$$

second,  $P(R^{2n+1})$

$$\text{Let } (a, b) \in R^{2n+1} = R \cdot R^n \cdot R^n \quad \text{and } (b, c) \in R^{2n+1} = R^n \cdot R^n \cdot R$$

$$\text{Then } \exists p \exists q \exists x \exists y [(a, x) \in R^n \wedge (x, p) \in R^n \wedge (p, b) \in R^n \wedge (b, y) \in R^n \wedge (y, q) \in R^n \wedge (q, c) \in R^n]$$

$$\text{since } R^n \text{ is transitive, } (a, p) \in R^n \wedge (y, c) \in R^n.$$

$$\text{since } R \text{ is transitive, } (p, y) \in R^n$$

$$\Rightarrow (a, c) \in R^n \cdot R \cdot R^n = R^{2n+1} \quad \square.$$

$$1 \quad \forall R (\neg (\forall n \in \mathbb{Z}^+ ((n \geq 2) \rightarrow P(R^n))) \rightarrow P(R))$$

consider  $R = \{(a, b), (b, c)\}$ .

$$R^2 = \{(a, c)\}$$

$$R^n = \emptyset \text{ for } n > 2.$$

$\therefore P(R)$  is false but  $P(R^2), P(R^3), \dots$  are true.

$\therefore \forall n \in \mathbb{Z}^+ ((n \geq 2) \rightarrow P(R^n))$  is TRUE.

$$\therefore \exists R (\forall n \in \mathbb{Z}^+ ((n \geq 2) \rightarrow P(R^n)) \wedge \neg P(R))$$

$\therefore$  TRUE  $\Rightarrow$  original proposition is false.

$$R^* = R \cup R^2 \cup R^3 \cup \dots \quad (+)$$

Need to prove  $R^1 = R^*$

(i)  $R \subseteq R^*$  directly.

(ii) Need to show  $R^* \subseteq S$  whenever  $R \subseteq S$ , Stricture.

If  $(a, b) \in R^*$  &  $(b, c) \in R^*$  it follows from (+) that  $(a, c) \in R^*$ .

Let  $R \subseteq S$ . Then  $P(S^n) \wedge S^n \subseteq S$ .

$$S^* = S \cup S^2 \cup \dots \quad \text{and} \quad S^n \subseteq S \quad \text{so} \quad S^* \subseteq S.$$

$$R \subseteq S \Rightarrow R^* \subseteq S^*.$$

$$\therefore R^* \subseteq S^* \subseteq S \quad \text{D.}$$